

Jos Manuel Fernandez-Real

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/11326211/jose-manuel-fernandez-real-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118
papers

9,150
citations

45
h-index

95
g-index

120
ext. papers

10,744
ext. citations

9
avg, IF

6.04
L-index

#	Paper	IF	Citations
118	Specific adipose tissue gene knockdown prevents diet-induced body weight gain, impacting fat accretion-related gene and protein expression.. <i>Molecular Therapy - Nucleic Acids</i> , 2022 , 27, 870-879	10.7	1
117	Downregulation of peripheral lipopolysaccharide binding protein impacts on perigonadal adipose tissue only in female mice. <i>Biomedicine and Pharmacotherapy</i> , 2022 , 151, 113156	7.5	0
116	Adipose tissue knockdown of lysozyme reduces local inflammation and improves adipogenesis in high-fat diet-fed mice. <i>Pharmacological Research</i> , 2021 , 166, 105486	10.2	2
115	Lysozyme is a component of the innate immune system linked to obesity associated-chronic low-grade inflammation and altered glucose tolerance. <i>Clinical Nutrition</i> , 2021 , 40, 1420-1429	5.9	6
114	Activation of Endogenous HS Biosynthesis or Supplementation with Exogenous HS Enhances Adipose Tissue Adipogenesis and Preserves Adipocyte Physiology in Humans. <i>Antioxidants and Redox Signaling</i> , 2021 , 35, 319-340	8.4	8
113	Identification and validation of circulating miRNAs as endogenous controls in obstructive sleep apnea. <i>PLoS ONE</i> , 2019 , 14, e0213622	3.7	8
112	Glycated Hemoglobin, but not Insulin Sensitivity, is Associated with Memory in Subjects with Obesity. <i>Obesity</i> , 2019 , 27, 932-942	8	6
111	Neuregulin 4 Is a Novel Marker of Beige Adipocyte Precursor Cells in Human Adipose Tissue. <i>Frontiers in Physiology</i> , 2019 , 10, 39	4.6	12
110	Preoperative Circulating Succinate Levels as a Biomarker for Diabetes Remission After Bariatric Surgery. <i>Diabetes Care</i> , 2019 , 42, 1956-1965	14.6	27
109	Circulating Irisin and Myostatin as Markers of Muscle Strength and Physical Condition in Elderly Subjects. <i>Frontiers in Physiology</i> , 2019 , 10, 871	4.6	24
108	Iron Status and Metabolically Unhealthy Obesity in Prepubertal Children. <i>Obesity</i> , 2019 , 27, 636-644	8	7
107	Analysis of miRNA signatures in CSF identifies upregulation of miR-21 and miR-146a/b in patients with multiple sclerosis and active lesions. <i>Journal of Neuroinflammation</i> , 2019 , 16, 220	10.1	30
106	Microbiota impacts on chronic inflammation and metabolic syndrome - related cognitive dysfunction. <i>Reviews in Endocrine and Metabolic Disorders</i> , 2019 , 20, 473-480	10.5	20
105	The gut microbiota modulates both browning of white adipose tissue and the activity of brown adipose tissue. <i>Reviews in Endocrine and Metabolic Disorders</i> , 2019 , 20, 387-397	10.5	30
104	Adipose tissue TSH as a new modulator of human adipocyte mitochondrial function. <i>International Journal of Obesity</i> , 2019 , 43, 1611-1619	5.5	7
103	The complement system is dysfunctional in metabolic disease: Evidences in plasma and adipose tissue from obese and insulin resistant subjects. <i>Seminars in Cell and Developmental Biology</i> , 2019 , 85, 164-172	7.5	32
102	Elevated circulating levels of succinate in human obesity are linked to specific gut microbiota. <i>ISME Journal</i> , 2018 , 12, 1642-1657	11.9	132

101	Obesity status influences the relationship among serum osteocalcin, iron stores and insulin sensitivity. <i>Clinical Nutrition</i> , 2018 , 37, 2091-2096	5.9	0
100	Increased Small Intestine Expression of Non-Heme Iron Transporters in Morbidly Obese Patients With Newly Diagnosed Type 2 Diabetes. <i>Molecular Nutrition and Food Research</i> , 2018 , 62, 1700301	5.9	0
99	Adipose TSHB in Humans and Serum TSH in Hypothyroid Rats Inform About Cellular Senescence. <i>Cellular Physiology and Biochemistry</i> , 2018 , 51, 142-153	3.9	5
98	Ferritin, metabolic syndrome and its components: A systematic review and meta-analysis. <i>Atherosclerosis</i> , 2018 , 275, 97-106	3.1	29
97	Genetic deficiency of indoleamine 2,3-dioxygenase promotes gut microbiota-mediated metabolic health. <i>Nature Medicine</i> , 2018 , 24, 1113-1120	50.5	121
96	miRNAs in cerebrospinal fluid identify patients with MS and specifically those with lipid-specific oligoclonal IgM bands. <i>Multiple Sclerosis Journal</i> , 2017 , 23, 1716-1726	5	36
95	Physiology and role of irisin in glucose homeostasis. <i>Nature Reviews Endocrinology</i> , 2017 , 13, 324-337	15.2	259
94	Metformin alters the gut microbiome of individuals with treatment-naive type 2 diabetes, contributing to the therapeutic effects of the drug. <i>Nature Medicine</i> , 2017 , 23, 850-858	50.5	732
93	Adipocyte lipopolysaccharide binding protein (LBP) is linked to a specific lipidomic signature. <i>Obesity</i> , 2017 , 25, 391-400	8	6
92	Adipose tissue and serum CCDC80 in obesity and its association with related metabolic disease. <i>Molecular Medicine</i> , 2017 , 23, 225-234	6.2	7
91	Increased adipose tissue heme levels and exportation are associated with altered systemic glucose metabolism. <i>Scientific Reports</i> , 2017 , 7, 5305	4.9	6
90	Hepatic iron content is independently associated with serum hepcidin levels in subjects with obesity. <i>Clinical Nutrition</i> , 2017 , 36, 1434-1439	5.9	19
89	Nicotinamide N-methyltransferase expression decreases in iron overload, exacerbating toxicity in mouse hepatocytes. <i>Hepatology Communications</i> , 2017 , 1, 803-815	6	2
88	Adipocyte Differentiation 2017 , 69-90		9
87	Genetic identification of thiosulfate sulfurtransferase as an adipocyte-expressed antidiabetic target in mice selected for leanness. <i>Nature Medicine</i> , 2016 , 22, 771-9	50.5	33
86	Obesity Is Associated With Gene Expression and Imaging Markers of Iron Accumulation in Skeletal Muscle. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 1282-9	5.6	18
85	Influence of Dietary Factors on Gut Microbiota: The Role on Insulin Resistance and Diabetes Mellitus 2016 , 147-154		
84	Adipose tissue R2* signal is increased in subjects with obesity: A preliminary MRI study. <i>Obesity</i> , 2016 , 24, 352-8	8	6

83	Contrasting association of circulating sCD14 with insulin sensitivity in non-obese and morbidly obese subjects. <i>Molecular Nutrition and Food Research</i> , 2016 , 60, 103-9	5.9	7
82	Soluble transferrin receptor levels are positively associated with insulin resistance but not with the metabolic syndrome or its individual components. <i>British Journal of Nutrition</i> , 2016 , 116, 1165-1174	3.6	10
81	Lipopolysaccharide binding protein is an adipokine involved in the resilience of the mouse adipocyte to inflammation. <i>Diabetologia</i> , 2015 , 58, 2424-34	10.3	25
80	PRDM16 sustains white fat gene expression profile in human adipocytes in direct relation with insulin action. <i>Molecular and Cellular Endocrinology</i> , 2015 , 405, 84-93	4.4	9
79	Nicotinamide N-methyltransferase regulates hepatic nutrient metabolism through Sirt1 protein stabilization. <i>Nature Medicine</i> , 2015 , 21, 887-94	50.5	129
78	Mechanisms Linking Glucose Homeostasis and Iron Metabolism Toward the Onset and Progression of Type 2 Diabetes. <i>Diabetes Care</i> , 2015 , 38, 2169-76	14.6	107
77	Inflammation triggers specific microRNA profiles in human adipocytes and macrophages and in their supernatants. <i>Clinical Epigenetics</i> , 2015 , 7, 49	7.7	71
76	Olive Oil and the Senescent Bone 2015 , 505-512		
75	Circulating irisin levels are positively associated with metabolic risk factors in sedentary subjects. <i>PLoS ONE</i> , 2015 , 10, e0124100	3.7	53
74	Brain iron overload, insulin resistance, and cognitive performance in obese subjects: a preliminary MRI case-control study. <i>Diabetes Care</i> , 2014 , 37, 3076-83	14.6	40
73	Metabolism: Irisin, the metabolic syndrome and follistatin in humans. <i>Nature Reviews Endocrinology</i> , 2014 , 10, 11-2	15.2	26
72	Olive oil phenolic compounds decrease the postprandial inflammatory response by reducing postprandial plasma lipopolysaccharide levels. <i>Food Chemistry</i> , 2014 , 162, 161-71	8.5	45
71	Fine-tuned iron availability is essential to achieve optimal adipocyte differentiation and mitochondrial biogenesis. <i>Diabetologia</i> , 2014 , 57, 1957-67	10.3	39
70	Human omental and subcutaneous adipose tissue exhibit specific lipidomic signatures. <i>FASEB Journal</i> , 2014 , 28, 1071-81	0.9	38
69	Profiling of circulating microRNAs reveals common microRNAs linked to type 2 diabetes that change with insulin sensitization. <i>Diabetes Care</i> , 2014 , 37, 1375-83	14.6	241
68	Effects of iron overload on chronic metabolic diseases. <i>Lancet Diabetes and Endocrinology</i> , 2014 , 2, 513-26	18.1	131
67	Circulating tryptase as a marker for subclinical atherosclerosis in obese subjects. <i>PLoS ONE</i> , 2014 , 9, e97014	9.14	16
66	Insulin resistance modulates iron-related proteins in adipose tissue. <i>Diabetes Care</i> , 2014 , 37, 1092-100	14.6	43

65	The possible role of antimicrobial proteins in obesity-associated immunologic alterations. <i>Expert Review of Clinical Immunology</i> , 2014 , 10, 855-66	5.1	4
64	Adipose tissue Ecrystallin is a thyroid hormone-binding protein associated with systemic insulin sensitivity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E2259-68	5.6	6
63	Loss of control over eating: a description of the eating disorder/obesity spectrum in women. <i>European Eating Disorders Review</i> , 2014 , 22, 25-31	5.3	32
62	A role for adipocyte-derived lipopolysaccharide-binding protein in inflammation- and obesity-associated adipose tissue dysfunction. <i>Diabetologia</i> , 2013 , 56, 2524-37	10.3	75
61	The gut microbiota profile is associated with insulin action in humans. <i>Acta Diabetologica</i> , 2013 , 50, 753-61	6.1	39
60	Targeting the circulating microRNA signature of obesity. <i>Clinical Chemistry</i> , 2013 , 59, 781-92	5.5	281
59	Study of lactoferrin gene expression in human and mouse adipose tissue, human preadipocytes and mouse 3T3-L1 fibroblasts. Association with adipogenic and inflammatory markers. <i>Journal of Nutritional Biochemistry</i> , 2013 , 24, 1266-75	6.3	24
58	Irisin is expressed and produced by human muscle and adipose tissue in association with obesity and insulin resistance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E769-78	5.6	501
57	Iron and obesity status-associated insulin resistance influence circulating fibroblast-growth factor-23 concentrations. <i>PLoS ONE</i> , 2013 , 8, e58961	3.7	25
56	The MRC1/CD68 ratio is positively associated with adipose tissue lipogenesis and with muscle mitochondrial gene expression in humans. <i>PLoS ONE</i> , 2013 , 8, e70810	3.7	14
55	Lifetime obesity in patients with eating disorders: increasing prevalence, clinical and personality correlates. <i>European Eating Disorders Review</i> , 2012 , 20, 250-4	5.3	127
54	The L-lysophosphatidylinositol/GPR55 system and its potential role in human obesity. <i>Diabetes</i> , 2012 , 61, 281-91	0.9	112
53	Lucha por sobrevivir, diabetes tipo 2 y obesidad. <i>Avances En Diabetología</i> , 2012 , 28, 55-58		1
52	Back to past leeches: repeated phlebotomies and cardiovascular risk. <i>BMC Medicine</i> , 2012 , 10, 53	11.4	3
51	Adipocyte Differentiation 2012 , 17-38		36
50	Circulating zonulin, a marker of intestinal permeability, is increased in association with obesity-associated insulin resistance. <i>PLoS ONE</i> , 2012 , 7, e37160	3.7	165
49	Serum and urinary concentrations of calprotectin as markers of insulin resistance and type 2 diabetes. <i>European Journal of Endocrinology</i> , 2012 , 167, 569-78	6.5	44
48	Total and undercarboxylated osteocalcin predict changes in insulin sensitivity and β cell function in elderly men at high cardiovascular risk. <i>American Journal of Clinical Nutrition</i> , 2012 , 95, 249-55	7	65

47	A Mediterranean diet enriched with olive oil is associated with higher serum total osteocalcin levels in elderly men at high cardiovascular risk. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 3792-8	5.6	63
46	Genetic variation near IRS1 associates with reduced adiposity and an impaired metabolic profile. <i>Nature Genetics</i> , 2011 , 43, 753-60	36.3	237
45	Mirror extreme BMI phenotypes associated with gene dosage at the chromosome 16p11.2 locus. <i>Nature</i> , 2011 , 478, 97-102	50.4	322
44	Increased levels of calprotectin in obesity are related to macrophage content: impact on inflammation and effect of weight loss. <i>Molecular Medicine</i> , 2011 , 17, 1157-67	6.2	77
43	Antimicrobial-sensing proteins in obesity and type 2 diabetes: the buffering efficiency hypothesis. <i>Diabetes Care</i> , 2011 , 34 Suppl 2, S335-41	14.6	17
42	OCT1 Expression in adipocytes could contribute to increased metformin action in obese subjects. <i>Diabetes</i> , 2011 , 60, 168-76	0.9	73
41	CD14 modulates inflammation-driven insulin resistance. <i>Diabetes</i> , 2011 , 60, 2179-86	0.9	78
40	Transferrin receptor-1 gene polymorphisms are associated with type 2 diabetes. <i>European Journal of Clinical Investigation</i> , 2010 , 40, 600-7	4.6	19
39	Surfactant protein d, a marker of lung innate immunity, is positively associated with insulin sensitivity. <i>Diabetes Care</i> , 2010 , 33, 847-53	14.6	33
38	Complement factor H is expressed in adipose tissue in association with insulin resistance. <i>Diabetes</i> , 2010 , 59, 200-9	0.9	74
37	Extracellular fatty acid synthase: a possible surrogate biomarker of insulin resistance. <i>Diabetes</i> , 2010 , 59, 1506-11	0.9	38
36	Circulating pigment epithelium-derived factor levels are associated with insulin resistance and decrease after weight loss. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 4720-8	5.6	75
35	Lipopolysaccharide-binding protein and soluble CD14 in the vitreous fluid of patients with proliferative diabetic retinopathy. <i>Retina</i> , 2010 , 30, 345-52	3.6	10
34	Circulating bactericidal/permeability-increasing protein (BPI) is associated with serum lipids and endothelial function. <i>Thrombosis and Haemostasis</i> , 2010 , 103, 780-7	7	9
33	The decrease of serum levels of human neutrophil alpha-defensins parallels with the surgery-induced amelioration of NASH in obesity. <i>Obesity Surgery</i> , 2010 , 20, 1682-9	3.7	13
32	Fatty acid synthase: association with insulin resistance, type 2 diabetes, and cancer. <i>Clinical Chemistry</i> , 2009 , 55, 425-38	5.5	140
31	Adipocytokines and insulin resistance: the possible role of lipocalin-2, retinol binding protein-4, and adiponectin. <i>Diabetes Care</i> , 2009 , 32 Suppl 2, S362-7	14.6	125
30	Study of circulating prohepcidin in association with insulin sensitivity and changing iron stores. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 982-8	5.6	26

29	Increase in plasma endotoxin concentrations and the expression of Toll-like receptors and suppressor of cytokine signaling-3 in mononuclear cells after a high-fat, high-carbohydrate meal: implications for insulin resistance. <i>Diabetes Care</i> , 2009 , 32, 2281-7	14.6	365
28	Grape-seed procyanidins modulate inflammation on human differentiated adipocytes in vitro. <i>Cytokine</i> , 2009 , 47, 137-42	4	97
27	The relationship of serum osteocalcin concentration to insulin secretion, sensitivity, and disposal with hypocaloric diet and resistance training. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 237-45	5.6	223
26	Fatty acid synthase activity regulates HER2 extracellular domain shedding into the circulation of HER2-positive metastatic breast cancer patients. <i>International Journal of Oncology</i> , 2009 , 35, 1369-76	4.4	17
25	Resistencia a la insulina y aterosclerosis. Impacto del estr� oxidativo en la funci� endotelial. <i>Revista Espanola De Cardiologia Suplementos</i> , 2008 , 8, 45C-52C	0.2	
24	Innate immunity, insulin resistance and type 2 diabetes. <i>Trends in Endocrinology and Metabolism</i> , 2008 , 19, 10-6	8.8	139
23	Bloodletting ameliorates insulin sensitivity and secretion in parallel to reducing liver iron in carriers of HFE gene mutations. <i>Diabetes Care</i> , 2008 , 31, 3-8	14.6	42
22	Circulating surfactant protein A (SP-A), a marker of lung injury, is associated with insulin resistance. <i>Diabetes Care</i> , 2008 , 31, 958-63	14.6	20
21	Circulating retinol-binding protein-4 concentration might reflect insulin resistance-associated iron overload. <i>Diabetes</i> , 2008 , 57, 1918-25	0.9	37
20	Association of circulating lactoferrin concentration and 2 nonsynonymous LTF gene polymorphisms with dyslipidemia in men depends on glucose-tolerance status. <i>Clinical Chemistry</i> , 2008 , 54, 301-9	5.5	52
19	Monocyte chemoattractant protein-1 in obesity and type 2 diabetes. Insulin sensitivity study. <i>Obesity</i> , 2007 , 15, 664-72	8	40
18	Insulin-like growth factor binding protein-related protein 1 (IGFBP-rP1/MAC25) is linked to endothelial-dependent vasodilation in high-ferritin type 2 diabetes. <i>Diabetes Care</i> , 2007 , 30, 1615-7	14.6	6
17	Circulating visfatin is associated with parameters of iron metabolism in subjects with altered glucose tolerance. <i>Diabetes Care</i> , 2007 , 30, 616-21	14.6	45
16	Circulating soluble transferrin receptor according to glucose tolerance status and insulin sensitivity. <i>Diabetes Care</i> , 2007 , 30, 604-8	14.6	37
15	An alternatively spliced soluble TNF-alpha receptor is associated with metabolic disorders: a replication study. <i>Clinical Immunology</i> , 2006 , 121, 236-41	9	9
14	An alternative spliced variant of circulating soluble tumor necrosis factor-alpha receptor-2 is paradoxically associated with insulin action. <i>European Journal of Endocrinology</i> , 2006 , 154, 723-30	6.5	10
13	Natural antibiotics and insulin sensitivity: the role of bactericidal/permeability-increasing protein. <i>Diabetes</i> , 2006 , 55, 216-24	0.9	30
12	Dyslipidemia and inflammation: an evolutionary conserved mechanism. <i>Clinical Nutrition</i> , 2005 , 24, 16-31; 9		285

11	Iron stores, blood donation, and insulin sensitivity and secretion. <i>Clinical Chemistry</i> , 2005 , 51, 1201-5	5.5	114
10	Circulating soluble CD14 monocyte receptor is associated with increased alanine aminotransferase. <i>Clinical Chemistry</i> , 2004 , 50, 1456-8	5.5	9
9	Adiponectin, hepatocellular dysfunction and insulin sensitivity. <i>Clinical Endocrinology</i> , 2004 , 60, 256-63	3.4	83
8	Insulin resistance and chronic cardiovascular inflammatory syndrome. <i>Endocrine Reviews</i> , 2003 , 24, 278-301	3.2	653
7	Novel interactions of adiponectin with the endocrine system and inflammatory parameters. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 2714-8	5.6	136
6	CD14 monocyte receptor, involved in the inflammatory cascade, and insulin sensitivity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 1780-4	5.6	79
5	A polymorphism in the promoter of the tumor necrosis factor-alpha gene (-308) is associated with coronary heart disease in type 2 diabetic patients. <i>Atherosclerosis</i> , 2003 , 167, 257-64	3.1	93
4	Blood letting in high-ferritin type 2 diabetes: effects on insulin sensitivity and beta-cell function. <i>Diabetes</i> , 2002 , 51, 1000-4	0.9	270
3	Blood letting in high-ferritin type 2 diabetes: effects on vascular reactivity. <i>Diabetes Care</i> , 2002 , 25, 2249-55	1.5	63
2	Cross-talk between iron metabolism and diabetes. <i>Diabetes</i> , 2002 , 51, 2348-54	0.9	453
1	Plasma levels of the soluble fraction of tumor necrosis factor receptors 1 and 2 are independent determinants of plasma cholesterol and LDL-cholesterol concentrations in healthy subjects. <i>Atherosclerosis</i> , 1999 , 146, 321-7	3.1	40