

# Marcos Palatnik

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11332222/publications.pdf>

Version: 2024-02-01

20  
papers

1,044  
citations

471509

17  
h-index

752698

20  
g-index

20  
all docs

20  
docs citations

20  
times ranked

868  
citing authors

#	ARTICLE	IF	CITATIONS
1	A Chimera Containing CD4+ and CD8+ T-Cell Epitopes of the Leishmania donovani Nucleoside Hydrolase (NH36) Optimizes Cross-Protection against Leishmania amazonensis Infection. <i>Frontiers in Immunology</i> , 2017, 8, 100.	4.8	35
2	Leishmania donovani Nucleoside Hydrolase (NH36) Domains Induce T-Cell Cytokine Responses in Human Visceral Leishmaniasis. <i>Frontiers in Immunology</i> , 2017, 8, 227.	4.8	27
3	Cross-Protective Immunity to Leishmania amazonensis is Mediated by CD4+ and CD8+ Epitopes of Leishmania donovani Nucleoside Hydrolase Terminal Domains. <i>Frontiers in Immunology</i> , 2014, 5, 189.	4.8	31
4	Leishmania donovani Nucleoside Hydrolase Terminal Domains in Cross-Protective Immunotherapy Against Leishmania amazonensis Murine Infection. <i>Frontiers in Immunology</i> , 2014, 5, 273.	4.8	22
5	The Leishmune <sup>®</sup> 's Nucleoside hydrolase DNA vaccine as an aid in immunotherapy of canine visceral leishmaniasis. <i>Procedia in Vaccinology</i> , 2012, 6, 64-73.	0.4	13
6	Resistance to visceral leishmaniasis is severely compromised in mice deficient of bradykinin B2-receptors. <i>Parasites and Vectors</i> , 2012, 5, 261.	2.5	13
7	Adaptive Immunity against Leishmania Nucleoside Hydrolase Maps Its C-Terminal Domain as the Target of the CD4+ T Cell <sup>driven</sup> Protective Response. <i>PLoS Neglected Tropical Diseases</i> , 2010, 4, e866.	3.0	48
8	Decrease of the incidence of human and canine visceral leishmaniasis after dog vaccination with Leishmune <sup>®</sup> in Brazilian endemic areas. <i>Vaccine</i> , 2009, 27, 3505-3512.	3.8	91
9	Effective immunotherapy against canine visceral leishmaniasis with the FML-vaccine. <i>Vaccine</i> , 2004, 22, 2234-2243.	3.8	121
10	Improving methods for epidemiological control of canine visceral leishmaniasis based on a mathematical model. Impact on the incidence of the canine and human disease. <i>Anais Da Academia Brasileira De Ciencias</i> , 2004, 76, 583-593.	0.8	35
11	IgG1/IgG2 antibody dichotomy in sera of vaccinated or naturally infected dogs with visceral leishmaniasis. <i>Vaccine</i> , 2003, 21, 2589-2597.	3.8	62
12	Immunotherapy against murine experimental visceral leishmaniasis with the FML-vaccine. <i>Vaccine</i> , 2003, 21, 4668-4676.	3.8	44
13	Saponins, IL12 and BCG adjuvant in the FML-vaccine formulation against murine visceral leishmaniasis. <i>Vaccine</i> , 2002, 21, 30-43.	3.8	98
14	Occurrence of Leishmania donovani parasitemia in plasma of infected hamsters. <i>Acta Tropica</i> , 2001, 80, 69-75.	2.0	16
15	Vaccination of Balb/c mice against experimental visceral leishmaniasis with the GP36 glycoprotein antigen of Leishmania donovani. <i>Vaccine</i> , 2001, 19, 3104-3115.	3.8	43
16	A phase III trial of efficacy of the FML-vaccine against canine kala-azar in an endemic area of Brazil (Sicilia) Tj ETQq000 rgt / Overlock 102	3.8	102
17	Vaccination of Swiss Albino mice against experimental visceral leishmaniasis with the FML antigen of Leishmania donovani. <i>Vaccine</i> , 1999, 17, 2554-2561.	3.8	43
18	Haemolytic activities of plant saponins and adjuvants. Effect of Periandra mediterranea saponin on the humoral response to the FML antigen of Leishmania donovani. <i>Vaccine</i> , 1997, 15, 1024-1029.	3.8	84

#	ARTICLE	IF	CITATIONS
19	Prevalence of Anti-Leishmania donovani Antibody Among Brazilian Blood Donors and Multiply Transfused Hemodialysis Patients. <i>American Journal of Tropical Medicine and Hygiene</i> , 1997, 57, 168-171.	1.4	67
20	<i>Leishmania donovani</i> : titration of antibodies to the fucose-mannose ligand as an aid in diagnosis and prognosis of visceral leishmaniasis. <i>Transactions of the Royal Society of Tropical Medicine and Hygiene</i> , 1995, 89, 390-393.	1.8	49