

Eiji Kikuchi

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1133788/eiji-kikuchi-publications-by-citations.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

281
papers

7,550
citations

42
h-index

74
g-index

298
ext. papers

8,880
ext. citations

3.7
avg, IF

5.33
L-index

#	Paper	IF	Citations
281	Outcomes of radical nephroureterectomy: a series from the Upper Tract Urothelial Carcinoma Collaboration. <i>Cancer</i> , 2009 , 115, 1224-33	6.4	739
280	Lymphovascular invasion predicts clinical outcomes in patients with node-negative upper tract urothelial carcinoma. <i>Journal of Clinical Oncology</i> , 2009 , 27, 612-8	2.2	209
279	Adjuvant chemotherapy for high risk upper tract urothelial carcinoma: results from the Upper Tract Urothelial Carcinoma Collaboration. <i>Journal of Urology</i> , 2009 , 182, 900-6	2.5	172
278	Impact of lymph node dissection on cancer specific survival in patients with upper tract urothelial carcinoma treated with radical nephroureterectomy. <i>Journal of Urology</i> , 2009 , 181, 2482-9	2.5	154
277	Predicting clinical outcomes after radical nephroureterectomy for upper tract urothelial carcinoma. <i>European Urology</i> , 2012 , 61, 818-25	10.2	153
276	Preoperative multivariable prognostic model for prediction of nonorgan confined urothelial carcinoma of the upper urinary tract. <i>Journal of Urology</i> , 2010 , 184, 453-8	2.5	151
275	The impact of tumor multifocality on outcomes in patients treated with radical nephroureterectomy. <i>European Urology</i> , 2012 , 61, 245-53	10.2	135
274	Impact of tumor location on prognosis for patients with upper tract urothelial carcinoma managed by radical nephroureterectomy. <i>European Urology</i> , 2010 , 57, 1072-9	10.2	133
273	Epidemiology, diagnosis, preoperative evaluation and prognostic assessment of upper-tract urothelial carcinoma (UTUC). <i>World Journal of Urology</i> , 2017 , 35, 379-387	4	132
272	Tumour architecture is an independent predictor of outcomes after nephroureterectomy: a multi-institutional analysis of 1363 patients. <i>BJU International</i> , 2009 , 103, 307-11	5.6	131
271	Comparison of oncologic outcomes for open and laparoscopic nephroureterectomy: a multi-institutional analysis of 1249 cases. <i>European Urology</i> , 2009 , 56, 1-9	10.2	130
270	The extent of lymphadenectomy seems to be associated with better survival in patients with nonmetastatic upper-tract urothelial carcinoma: how many lymph nodes should be removed?. <i>European Urology</i> , 2009 , 56, 512-8	10.2	126
269	Tumour necrosis is an indicator of aggressive biology in patients with urothelial carcinoma of the upper urinary tract. <i>European Urology</i> , 2010 , 57, 575-81	10.2	125
268	Is tumor volume an independent prognostic factor in clinically localized prostate cancer?. <i>Journal of Urology</i> , 2004 , 172, 508-11	2.5	123
267	Conditional survival after radical cystectomy for bladder cancer: evidence for a patient changing risk profile over time. <i>European Urology</i> , 2014 , 66, 361-70	10.2	94
266	Angiotensin II type 1 receptor antagonist as an angiogenic inhibitor in prostate cancer. <i>Prostate</i> , 2007 , 67, 41-9	4.2	90
265	Interleukin 6 is associated with cachexia in patients with prostate cancer. <i>Urology</i> , 2007 , 69, 113-7	1.6	88

264	Preoperative hydronephrosis grade independently predicts worse pathological outcomes in patients undergoing nephroureterectomy for upper tract urothelial carcinoma. <i>Journal of Urology</i> , 2011 , 185, 1621-6	2.5	84
263	Advanced patient age is associated with inferior cancer-specific survival after radical nephroureterectomy. <i>BJU International</i> , 2010 , 105, 1672-7	5.6	84
262	Lymphovascular invasion independently predicts increased disease specific survival in patients with transitional cell carcinoma of the upper urinary tract. <i>Journal of Urology</i> , 2005 , 174, 2120-3; discussion 2124	2.5	79
261	Impact of histological variants on clinical outcomes of patients with upper urinary tract urothelial carcinoma. <i>Journal of Urology</i> , 2012 , 188, 398-404	2.5	78
260	Gender-specific differences in clinicopathologic outcomes following radical cystectomy: an international multi-institutional study of more than 8000 patients. <i>European Urology</i> , 2014 , 66, 913-9	10.2	77
259	Suppression of hormone-refractory prostate cancer by a novel nuclear factor kappaB inhibitor in nude mice. <i>Cancer Research</i> , 2003 , 63, 107-10	10.1	77
258	Activation of the aryl hydrocarbon receptor pathway enhances cancer cell invasion by upregulating the MMP expression and is associated with poor prognosis in upper urinary tract urothelial cancer. <i>Carcinogenesis</i> , 2010 , 31, 287-95	4.6	76
257	Angiotensin II type 1 receptor antagonist candesartan as an angiogenic inhibitor in a xenograft model of bladder cancer. <i>Clinical Cancer Research</i> , 2006 , 12, 2888-93	12.9	75
256	Comparison of CT urography and excretory urography in the detection and localization of urothelial carcinoma of the upper urinary tract. <i>American Journal of Roentgenology</i> , 2011 , 196, 1102-9	5.4	74
255	Risk stratification of patients with nodal involvement in upper tract urothelial carcinoma: value of lymph-node density. <i>BJU International</i> , 2009 , 103, 302-6	5.6	73
254	Nicotine Induces Tumor Growth and Chemoresistance through Activation of the PI3K/Akt/mTOR Pathway in Bladder Cancer. <i>Molecular Cancer Therapeutics</i> , 2015 , 14, 2112-20	6.1	66
253	Concomitant carcinoma in situ is a feature of aggressive disease in patients with organ confined urothelial carcinoma following radical nephroureterectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2012 , 30, 252-8	2.8	65
252	Evidence-based sex-related outcomes after radical nephroureterectomy for upper tract urothelial carcinoma: results of large multicenter study. <i>Urology</i> , 2009 , 73, 142-6	1.6	62
251	Upper urinary tract urothelial carcinoma with loco-regional nodal metastases: insights from the Upper Tract Urothelial Carcinoma Collaboration. <i>BJU International</i> , 2011 , 108, 1286-91	5.6	61
250	The predictive value of C-reactive protein for prognosis in patients with upper tract urothelial carcinoma treated with radical nephroureterectomy: a multi-institutional study. <i>European Urology</i> , 2014 , 65, 227-34	10.2	56
249	Assessment of long-term quality of life using the FACT-BL questionnaire in patients with an ileal conduit, continent reservoir, or orthotopic neobladder. <i>Japanese Journal of Clinical Oncology</i> , 2006 , 36, 712-6	2.8	53
248	A multi-institutional validation of the prognostic value of the neutrophil-to-lymphocyte ratio for upper tract urothelial carcinoma treated with radical nephroureterectomy. <i>Annals of Surgical Oncology</i> , 2014 , 21, 4041-8	3.1	51
247	Expression of snail in upper urinary tract urothelial carcinoma: prognostic significance and implications for tumor invasion. <i>Clinical Cancer Research</i> , 2010 , 16, 5814-23	12.9	51

246	Vitamin E succinate induced apoptosis and enhanced chemosensitivity to paclitaxel in human bladder cancer cells in vitro and in vivo. <i>Cancer Science</i> , 2010 , 101, 216-23	6.9	50
245	Assessment of the minimum number of lymph nodes needed to detect lymph node invasion at radical nephroureterectomy in patients with upper tract urothelial cancer. <i>Urology</i> , 2009 , 74, 1070-4	1.6	48
244	Gemcitabine and paclitaxel chemotherapy for advanced urothelial carcinoma in patients who have received prior cisplatin-based chemotherapy. <i>International Journal of Clinical Oncology</i> , 2008 , 13, 510-4	4.2	47
243	Technical difficulties of transumbilical laparoendoscopic single-site adrenalectomy: comparison with conventional laparoscopic adrenalectomy. <i>World Journal of Urology</i> , 2013 , 31, 199-203	4	46
242	Dual PI3K/mTOR inhibitor NVP-BEZ235 sensitizes docetaxel in castration resistant prostate cancer. <i>Journal of Urology</i> , 2014 , 191, 227-34	2.5	46
241	Prognostic value of plasma fibrinogen levels in patients with localized upper tract urothelial carcinoma. <i>BJU International</i> , 2013 , 111, 857-64	5.6	42
240	Prognostic value of extranodal extension and other lymph node parameters in patients with upper tract urothelial carcinoma. <i>Journal of Urology</i> , 2012 , 187, 845-51	2.5	42
239	Risk of cancer-specific mortality following recurrence after radical nephroureterectomy. <i>Annals of Surgical Oncology</i> , 2012 , 19, 4337-44	3.1	42
238	Prostate specific antigen adjusted for transition zone volume: the most powerful method for detecting prostate carcinoma. <i>Cancer</i> , 2000 , 89, 842-9	6.4	42
237	Evidenced-based clinical practice guideline for upper tract urothelial carcinoma (summary--Japanese Urological Association, 2014 edition). <i>International Journal of Urology</i> , 2015 , 22, 3-13	2.3	41
236	Long-term androgen ablation and docetaxel up-regulate phosphorylated Akt in castration resistant prostate cancer. <i>Journal of Urology</i> , 2011 , 185, 2376-81	2.5	40
235	Role of computed tomography urography in the clinical evaluation of upper tract urothelial carcinoma. <i>International Journal of Urology</i> , 2016 , 23, 284-98	2.3	40
234	Angiotensin II type 1 receptor expression and microvessel density in human bladder cancer. <i>Urology</i> , 2011 , 77, 1009.e19-25	1.6	37
233	Visceral obesity is a strong predictor of perioperative outcome in patients undergoing laparoscopic radical nephrectomy. <i>BJU International</i> , 2012 , 110, E980-4	5.6	36
232	Ets-1 and hypoxia inducible factor-1alpha inhibition by angiotensin II type-1 receptor blockade in hormone-refractory prostate cancer. <i>Prostate</i> , 2010 , 70, 162-9	4.2	36
231	Prevention of cancer cachexia by a novel nuclear factor {kappa}B inhibitor in prostate cancer. <i>Clinical Cancer Research</i> , 2005 , 11, 5590-4	12.9	36
230	The predictive value of positive urine cytology for outcomes following radical nephroureterectomy in patients with primary upper tract urothelial carcinoma: a multi-institutional study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 48.e19-26	2.8	35
229	Clinical outcome of tumor recurrence for Ta, T1 non-muscle invasive bladder cancer from the data on registered bladder cancer patients in Japan: 1999-2001 report from the Japanese Urological Association. <i>International Journal of Urology</i> , 2009 , 16, 279-86	2.3	35

228	Preoperative T categorization and prediction of histopathologic grading of urothelial carcinoma in renal pelvis using diffusion-weighted MRI. <i>American Journal of Roentgenology</i> , 2011 , 197, 1130-6	5.4	34
227	Subclassification of pT3 urothelial carcinoma of the renal pelvicalyceal system is associated with recurrence-free and cancer-specific survival: proposal for a revision of the current TNM classification. <i>European Urology</i> , 2012 , 62, 224-31	10.2	33
226	Metastatic behavior of upper tract urothelial carcinoma after radical nephroureterectomy: association with primary tumor location. <i>Annals of Surgical Oncology</i> , 2014 , 21, 1038-45	3.1	32
225	The prognostic significance of vasohibin-1 expression in patients with upper urinary tract urothelial carcinoma. <i>Clinical Cancer Research</i> , 2012 , 18, 4145-53	12.9	32
224	Impact of smoking status on bladder tumor recurrence after radical nephroureterectomy for upper tract urothelial carcinoma. <i>Journal of Urology</i> , 2013 , 189, 2062-8	2.5	31
223	Regulation of monocyte chemoattractant protein-1 through angiotensin II type 1 receptor in prostate cancer. <i>American Journal of Pathology</i> , 2012 , 180, 1008-1016	5.8	31
222	Racial differences in the outcome of patients with urothelial carcinoma of the upper urinary tract: an international study. <i>BJU International</i> , 2011 , 108, E304-9	5.6	31
221	The prognostic significance of vasohibin-1 expression in patients with prostate cancer. <i>British Journal of Cancer</i> , 2013 , 108, 2123-9	8.7	30
220	Inhibition of orthotopic human bladder tumor growth by lentiviral gene transfer of endostatin. <i>Clinical Cancer Research</i> , 2004 , 10, 1835-42	12.9	30
219	Prognostic significance of Bacillus Calmette-Guérin failure classification in non-muscle-invasive bladder cancer. <i>BJU International</i> , 2012 , 110, E216-21	5.6	29
218	Down-regulation of NF kappa B activation is an effective therapeutic modality in acquired platinum-resistant bladder cancer. <i>BMC Cancer</i> , 2015 , 15, 324	4.8	28
217	Delivery of replication-competent retrovirus expressing Escherichia coli purine nucleoside phosphorylase increases the metabolism of the prodrug, fludarabine phosphate and suppresses the growth of bladder tumor xenografts. <i>Cancer Gene Therapy</i> , 2007 , 14, 279-86	5.4	28
216	Activation of aryl hydrocarbon receptor promotes invasion of clear cell renal cell carcinoma and is associated with poor prognosis and cigarette smoke. <i>International Journal of Cancer</i> , 2015 , 137, 299-310	7.5	27
215	Extraperitoneal approach induces postoperative inguinal hernia compared with transperitoneal approach after laparoscopic radical prostatectomy. <i>Japanese Journal of Clinical Oncology</i> , 2010 , 40, 349-52	2.8	27
214	The predictors of local recurrence after radical cystectomy in patients with invasive bladder cancer. <i>Japanese Journal of Clinical Oncology</i> , 2008 , 38, 360-4	2.8	27
213	Sulfasalazine could modulate the CD44v9-xCT system and enhance cisplatin-induced cytotoxic effects in metastatic bladder cancer. <i>Cancer Science</i> , 2019 , 110, 1431-1441	6.9	26
212	Tumor budding, a novel prognostic indicator for predicting stage progression in T1 bladder cancers. <i>Cancer Science</i> , 2016 , 107, 1338-44	6.9	26
211	Patient characteristics and outcomes in metastatic upper tract urothelial carcinoma after radical nephroureterectomy: the experience of Japanese multi-institutions. <i>BJU International</i> , 2013 , 112, E28-34	5.6	26

210	Acquired platinum resistance enhances tumour angiogenesis through angiotensin II type 1 receptor in bladder cancer. <i>British Journal of Cancer</i> , 2011 , 105, 1331-7	8.7	26
209	Enhancement of radiosensitivity by a unique novel NF- κ B inhibitor, DHMEQ, in prostate cancer. <i>British Journal of Cancer</i> , 2012 , 107, 652-7	8.7	26
208	Effect of angiotensin II type 1 receptor antagonist on tumor growth and angiogenesis in a xenograft model of human bladder cancer. <i>Human Cell</i> , 2007 , 20, 1-9	4.5	26
207	High Absolute Monocyte Count Predicts Poor Clinical Outcome in Patients with Castration-Resistant Prostate Cancer Treated with Docetaxel Chemotherapy. <i>Annals of Surgical Oncology</i> , 2016 , 23, 4115-4122	3.1	26
206	Prognostic value of renin-angiotensin system blockade in non-muscle-invasive bladder cancer. <i>Annals of Surgical Oncology</i> , 2012 , 19, 3987-93	3.1	25
205	Discontinuance of bacille Calmette-Guérin instillation therapy for nonmuscle-invasive bladder cancer has negative effect on tumor recurrence. <i>Urology</i> , 2009 , 73, 1318-22	1.6	25
204	Neoadjuvant gemcitabine plus cisplatin for muscle-invasive bladder cancer. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 908-14	2.8	25
203	Preoperative prognostic nomogram (probability table) for renal cell carcinoma based on TNM classification. <i>Journal of Urology</i> , 2009 , 181, 480-5; discussion 485	2.5	25
202	Herpes simplex virus based gene therapy enhances the efficacy of mitomycin C for the treatment of human bladder transitional cell carcinoma. <i>Journal of Urology</i> , 2005 , 174, 741-6	2.5	25
201	Laparoscopic Versus Open Nephroureterectomy in Muscle-Invasive Upper Tract Urothelial Carcinoma: Subanalysis of the Multi-Institutional National Database of the Japanese Urological Association. <i>Journal of Endourology</i> , 2016 , 30, 520-5	2.7	24
200	Essential content of evidence-based clinical practice guidelines for bladder cancer: The Japanese Urological Association 2015 update. <i>International Journal of Urology</i> , 2016 , 23, 640-5	2.3	24
199	Postoperative Nomogram for Relapse-Free Survival in Patients with High Grade Upper Tract Urothelial Carcinoma. <i>Journal of Urology</i> , 2017 , 197, 580-589	2.5	23
198	The clinical significance of urine cytology after a radical cystectomy for urothelial cancer. <i>International Journal of Urology</i> , 2010 , 17, 527-32	2.3	23
197	Prognostic role of KiSS-1 and possibility of therapeutic modality of metastatin, the final peptide of the KiSS-1 gene, in urothelial carcinoma. <i>Molecular Cancer Therapeutics</i> , 2012 , 11, 853-63	6.1	23
196	Highly efficient gene delivery for bladder cancers by intravesically administered replication-competent retroviral vectors. <i>Clinical Cancer Research</i> , 2007 , 13, 4511-8	12.9	23
195	Visceral obesity is associated with better recurrence-free survival after curative surgery for Japanese patients with localized clear cell renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2015 , 45, 210-6	2.8	22
194	Impact of an adjuvant chemotherapeutic regimen on the clinical outcome in high risk patients with upper tract urothelial carcinoma: a Japanese multi-institution experience. <i>Journal of Urology</i> , 2015 , 193, 1122-8	2.5	22
193	Predictive factors for severe and febrile neutropenia during docetaxel chemotherapy for castration-resistant prostate cancer. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 605-12	4.2	22

192	Cis-dichlorodiammineplatinum upregulates angiotensin II type 1 receptors through reactive oxygen species generation and enhances VEGF production in bladder cancer. <i>Molecular Cancer Therapeutics</i> , 2010 , 9, 2982-92	6.1	22
191	Tumour length of the largest focus predicts prostate-specific antigen-based recurrence after radical prostatectomy in clinically localized prostate cancer. <i>BJU International</i> , 2009 , 104, 1215-8	5.6	22
190	Prognostic impact of renin-angiotensin system blockade in localised upper-tract urothelial carcinoma. <i>British Journal of Cancer</i> , 2012 , 106, 290-6	8.7	22
189	Detection and quantitative analysis of early stage orthotopic murine bladder tumor using in vivo magnetic resonance imaging. <i>Journal of Urology</i> , 2003 , 170, 1375-8	2.5	22
188	Variant isoforms of CD44 involves acquisition of chemoresistance to cisplatin and has potential as a novel indicator for identifying a cisplatin-resistant population in urothelial cancer. <i>BMC Cancer</i> , 2018 , 18, 113	4.8	21
187	Contrast-enhanced ultrasonography of the prostate with Sonazoid. <i>Japanese Journal of Clinical Oncology</i> , 2010 , 40, 1099-104	2.8	21
186	The Conditional Survival with Time of Intravesical Recurrence of Upper Tract Urothelial Carcinoma. <i>Journal of Urology</i> , 2017 , 198, 1278-1285	2.5	20
185	Laparoendoscopic single-site adrenalectomy versus conventional laparoscopic adrenalectomy: a comparison of surgical outcomes and an analysis of a single surgeon's learning curve. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2014 , 28, 2911-9	5.2	20
184	Long term follow-up in patients with initially diagnosed low grade Ta non-muscle invasive bladder tumors: tumor recurrence and worsening progression. <i>BMC Urology</i> , 2014 , 14, 5	2.2	20
183	Transumbilical approach for laparo-endoscopic single-site adrenalectomy: initial experience and short-term outcome. <i>International Journal of Urology</i> , 2012 , 19, 331-5	2.3	20
182	Prognostic significance of Bcl-xL expression and efficacy of Bcl-xL targeting therapy in urothelial carcinoma. <i>British Journal of Cancer</i> , 2013 , 108, 2312-20	8.7	20
181	Late recurrence and progression in non-muscle-invasive bladder cancers after 5-year tumor-free periods. <i>Urology</i> , 2010 , 75, 1385-90	1.6	20
180	Angiotensin II type 1 receptor antagonist enhances cis-dichlorodiammineplatinum-induced cytotoxicity in mouse xenograft model of bladder cancer. <i>Urology</i> , 2009 , 73, 655-60	1.6	20
179	Angiotensin II Type 1 Receptor Antagonist as an Angiogenic Inhibitor in Urogenital Cancer. <i>Reviews on Recent Clinical Trials</i> , 2009 , 4, 75-8	1.2	20
178	Initial experience of transumbilical laparoendoscopic single-site surgery of partial adrenalectomy in patient with aldosterone-producing adenoma. <i>BMC Urology</i> , 2010 , 10, 19	2.2	20
177	Simple stratification of survival using bone scan and serum C-reactive protein in prostate cancer patients with metastases. <i>Urologia Internationalis</i> , 2008 , 80, 129-33	1.9	20
176	Acquired platinum resistance involves epithelial to mesenchymal transition through ubiquitin ligase FBXO32 dysregulation. <i>JCI Insight</i> , 2016 , 1, e83654	9.9	20
175	Bladder cancer develops 6 years earlier in current smokers: analysis of bladder cancer registry data collected by the cancer registration committee of the Japanese Urological Association. <i>International Journal of Urology</i> , 2009 , 16, 64-9	2.3	19

174	Clinical Practice Guidelines for Bladder Cancer 2019 update by the Japanese Urological Association: Summary of the revision. <i>International Journal of Urology</i> , 2020 , 27, 702-709	2.3	18
173	The pretreatment neutrophil-to-lymphocyte ratio is a novel biomarker for predicting clinical responses to pembrolizumab in platinum-resistant metastatic urothelial carcinoma patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 602.e1-602.e10	2.8	18
172	Relationship Between Increased Expression of the Axl/Gas6 Signal Cascade and Prognosis of Patients with Upper Tract Urothelial Carcinoma. <i>Annals of Surgical Oncology</i> , 2016 , 23, 663-70	3.1	18
171	The prognostic significance of OCT4 expression in patients with prostate cancer. <i>Human Pathology</i> , 2016 , 51, 1-8	3.7	18
170	A prospective longitudinal survey of erectile dysfunction in patients with localized prostate cancer treated with permanent prostate brachytherapy. <i>Journal of Urology</i> , 2013 , 189, 1014-8	2.5	18
169	Intravesical interleukin-15 gene therapy in an orthotopic bladder cancer model. <i>Human Gene Therapy</i> , 2011 , 22, 1423-32	4.8	18
168	Preoperative hydronephrosis: independent predictor for changes in renal function following nephroureterectomy. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 202-7	2.8	18
167	Prognostic Impact of Renin-Angiotensin System Blockade on Renal Cell Carcinoma After Surgery. <i>Annals of Surgical Oncology</i> , 2015 , 22, 3751-9	3.1	17
166	Refraining from Smoking for 15 Years or More Reduced the Risk of Tumor Recurrence in Non-muscle Invasive Bladder Cancer Patients. <i>Annals of Surgical Oncology</i> , 2016 , 23, 1752-9	3.1	17
165	How does visceral obesity affect surgical performance in laparoscopic radical nephrectomy?. <i>Japanese Journal of Clinical Oncology</i> , 2015 , 45, 373-7	2.8	17
164	Predictors of survival in patients with disease recurrence after radical nephroureterectomy. <i>BJU International</i> , 2014 , 113, 911-7	5.6	17
163	Prognostic value of preoperative multiparametric magnetic resonance imaging (MRI) for predicting biochemical recurrence after radical prostatectomy. <i>BJU International</i> , 2014 , 113, 741-7	5.6	17
162	Risk factors for perioperative complications of laparoscopic adrenalectomy including single-site surgery. <i>Journal of Endourology</i> , 2012 , 26, 1463-7	2.7	17
161	Extrapancreatic solid pseudopapillary tumor: case report and review of the literature. <i>International Journal of Clinical Oncology</i> , 2012 , 17, 165-8	4.2	17
160	Surgical and chemotherapeutic options for urachal carcinoma: report of ten cases and literature review. <i>Urologia Internationalis</i> , 2012 , 88, 209-14	1.9	17
159	Plasma fibrinogen level: an independent prognostic factor for disease-free survival and cancer-specific survival in patients with localised renal cell carcinoma. <i>BJU International</i> , 2016 , 118, 598-603	5.6	17
158	Surgical outcome of laparoscopic surgery, including laparoendoscopic single-site surgery, for retroperitoneal paraganglioma compared with adrenal pheochromocytoma. <i>Journal of Endourology</i> , 2014 , 28, 686-92	2.7	16
157	Renin-angiotensin system blockade: Its contribution and controversy. <i>International Journal of Urology</i> , 2015 , 22, 721-30	2.3	16

156	Visceral fat is correlated with prolonged operative time in laparoendoscopic single-site adrenalectomy and laparoscopic adrenalectomy. <i>Urology</i> , 2013 , 82, 1312-8	1.6	16
155	Urinary pH is highly associated with tumor recurrence during intravesical mitomycin C therapy for nonmuscle invasive bladder tumor. <i>Journal of Urology</i> , 2011 , 185, 802-6	2.5	16
154	Role of pelvic lymph node dissection in lymph node-negative patients with invasive bladder cancer. <i>Japanese Journal of Clinical Oncology</i> , 2010 , 40, 247-51	2.8	16
153	Gemcitabine and cisplatin chemotherapy induced reversible posterior leukoencephalopathy syndrome in a bladder cancer patient. <i>International Journal of Clinical Oncology</i> , 2010 , 15, 508-11	4.2	16
152	Papillary renal cell carcinoma: clinicopathological characteristics in 40 patients. <i>Clinical and Experimental Nephrology</i> , 2008 , 12, 195-9	2.5	16
151	Single-cell RNA-seq analysis reveals the platinum resistance gene COX7B and the surrogate marker CD63. <i>Cancer Medicine</i> , 2018 , 7, 6193-6204	4.8	16
150	Clinical Role of Programmed Cell Death-1 Expression in Patients with Non-muscle-invasive Bladder Cancer Recurring After Initial Bacillus Calmette-Guérin Therapy. <i>Annals of Surgical Oncology</i> , 2018 , 25, 2484-2491	3.1	15
149	Independent predictors for bladder outcomes after treatment of intravesical recurrence following radical nephroureterectomy in patients with primary upper tract urothelial carcinoma. <i>Annals of Surgical Oncology</i> , 2014 , 21, 3151-8	3.1	15
148	C-reactive protein: a biomarker of survival in patients with localized upper tract urothelial carcinoma treated with radical nephroureterectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1725-30	2.8	15
147	Unidirectional barbed suture for vesicourethral anastomosis during laparoscopic radical prostatectomy. <i>Asian Journal of Endoscopic Surgery</i> , 2014 , 7, 241-5	1.4	15
146	Potent cytotoxic effect of a novel nuclear factor-kappaB inhibitor dehydroxymethylepoxyquinomicin on human bladder cancer cells producing various cytokines. <i>Urology</i> , 2010 , 75, 805-12	1.6	15
145	The impact of previous ureteroscopic tumor ablation on oncologic outcomes after radical nephroureterectomy for upper urinary tract urothelial carcinoma. <i>Journal of Endourology</i> , 2011 , 25, 775-9	2.7	15
144	Does pelvic lymph node dissection improve the biochemical relapse-free survival in low-risk prostate cancer patients treated by laparoscopic radical prostatectomy?. <i>Journal of Endourology</i> , 2012 , 26, 1199-202	2.7	15
143	The novel NF-kappaB activation inhibitor dehydroxymethyl-epoxyquinomicin suppresses anti-Thy1.1-induced glomerulonephritis in rats. <i>Nephron Experimental Nephrology</i> , 2008 , 110, e17-24		15
142	Impact of baseline visceral fat accumulation on prognosis in patients with metastatic renal cell carcinoma treated with systemic therapy. <i>Medical Oncology</i> , 2017 , 34, 47	3.7	14
141	Could patient age influence tumor recurrence rate in non-muscle-invasive bladder cancer patients treated with BCG immunotherapy?. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 565-70	2.8	14
140	Prediction of extraprostatic extension by prostate specific antigen velocity, endorectal MRI, and biopsy Gleason score in clinically localized prostate cancer. <i>International Journal of Urology</i> , 2008 , 15, 520-3	2.3	14
139	Variant isoforms of CD44 expression in upper tract urothelial cancer as a predictive marker for recurrence and mortality. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016 , 34, 337.e19-26	2.8	14

138	The Preoperative Neutrophil-to-lymphocyte Ratio is a Novel Biomarker for Predicting Worse Clinical Outcomes in Non-muscle Invasive Bladder Cancer Patients with a Previous History of Smoking. <i>Annals of Surgical Oncology</i> , 2016 , 23, 1039-1047	3.1	14
137	Comparison of continence outcomes of early catheter removal on postoperative day 2 and 4 after laparoscopic radical prostatectomy: a randomized controlled trial. <i>BMC Urology</i> , 2015 , 15, 77	2.2	13
136	Prognostic Value of Baseline Serum C-Reactive Protein Level in Intermediate-Risk Group Patients With Metastatic Renal-Cell Carcinoma Treated by First-Line Vascular Endothelial Growth Factor-Targeted Therapy. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, e927-e933	3.3	13
135	Human castration resistant prostate cancer rather prefer to decreased 5 α -reductase activity. <i>Scientific Reports</i> , 2013 , 3, 1268	4.9	13
134	Docetaxel in combination with prednisolone for hormone refractory prostate cancer. <i>Japanese Journal of Clinical Oncology</i> , 2010 , 40, 79-84	2.8	13
133	Impact of Combined Use of Blood-based Inflammatory Markers on Patients with Upper Tract Urothelial Carcinoma Following Radical Nephroureterectomy: Proposal of a Cumulative Marker Score as a Novel Predictive Tool for Prognosis. <i>European Urology Focus</i> , 2015 , 1, 54-63	5.1	12
132	Purified Protein Derivative Skin test prior to bacillus Calmette-Gu \acute{e} rin Therapy May have Therapeutic Impact in Patients with Nonmuscle Invasive Bladder Cancer. <i>Journal of Urology</i> , 2018 , 199, 1446-1451	2.5	12
131	Risk of subsequent tumour recurrence and stage progression in bacille Calmette-Gu \acute{e} rin relapsing non-muscle-invasive bladder cancer. <i>BJU International</i> , 2012 , 110, E508-13	5.6	12
130	Phosphorylated Akt up-regulates angiotensin II type-1 receptor expression in castration resistant prostate cancer. <i>Prostate</i> , 2011 , 71, 1510-7	4.2	12
129	Prediction of PSA bounce after permanent prostate brachytherapy for localized prostate cancer. <i>International Journal of Clinical Oncology</i> , 2009 , 14, 502-6	4.2	12
128	Carcinoma in situ of the upper urinary tract treated with radical nephroureterectomy--results from a multicenter study. <i>European Urology</i> , 2008 , 54, 961-3	10.2	12
127	Establishment of orthotopic mouse superficial bladder tumor model for studies on intravesical treatments. <i>Human Cell</i> , 2008 , 21, 57-63	4.5	12
126	Prolonged pneumoperitoneum time is an independent risk factor for intravesical recurrence after laparoscopic radical nephroureterectomy in upper tract urothelial carcinoma. <i>Surgical Oncology</i> , 2017 , 26, 73-79	2.5	11
125	Solifenacin or mirabegron could improve persistent overactive bladder symptoms after dutasteride treatment in patients with benign prostatic hyperplasia. <i>Urology</i> , 2015 , 85, 1151-1155	1.6	11
124	Determining When to Stop Prostate Specific Antigen Monitoring after Radical Prostatectomy: the Role of Ultrasensitive Prostate Specific Antigen. <i>Journal of Urology</i> , 2017 , 197, 655-661	2.5	11
123	Do metastatic upper tract urothelial carcinoma and bladder carcinoma have similar clinical responses to systemic chemotherapy? A Japanese multi-institutional experience. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 163-9	2.8	11
122	Prognostic significance of 5-fluorouracil metabolism-relating enzymes and enhanced chemosensitivity to 5-fluorouracil by 5-chloro 2,4-dihydroxy-pyridine in urothelial carcinoma. <i>BMC Cancer</i> , 2012 , 12, 420	4.8	11
121	Clinical practice patterns for upper tract urothelial carcinoma: a nationwide survey in Japan. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 768-74	2.8	11

120	Preoperative sarcopenia status is associated with lymphovascular invasion in upper tract urothelial carcinoma patients treated with radical nephroureterectomy. <i>Canadian Urological Association Journal</i> , 2018 , 12, E132-E136	1.2	10
119	Impact of renal function of patients with advanced urothelial cancer on eligibility for first-line chemotherapy and treatment outcomes. <i>Japanese Journal of Clinical Oncology</i> , 2015 , 45, 867-73	2.8	10
118	Transumbilical laparo-endoscopic single site surgery for adrenal cortical adenoma inducing primary aldosteronism: initial experience. <i>BMC Research Notes</i> , 2011 , 4, 364	2.3	10
117	How do young residents practice laparoscopic surgical skills?. <i>Urology</i> , 2010 , 76, 352-6	1.6	10
116	Simple and practical nomograms for predicting the stone-free rate after shock wave lithotripsy in patients with a solitary upper ureteral stone. <i>World Journal of Urology</i> , 2017 , 35, 1455-1461	4	9
115	Impact of acute kidney injury defined by CTCAE v4.0 during first course of cisplatin-based chemotherapy on treatment outcomes in advanced urothelial cancer patients. <i>Clinical and Experimental Nephrology</i> , 2017 , 21, 732-740	2.5	9
114	Intravesical dual PI3K/mTOR complex 1/2 inhibitor NVP-BEZ235 therapy in an orthotopic bladder cancer model. <i>International Journal of Oncology</i> , 2015 , 47, 377-83	4.4	9
113	Effect of a novel nuclear factor- κ B activation inhibitor on renal ischemia-reperfusion injury. <i>Transplantation</i> , 2013 , 96, 863-70	1.8	9
112	Frequency of tumor recurrence: a strong predictor of stage progression in initially diagnosed nonmuscle invasive bladder cancer. <i>Journal of Urology</i> , 2011 , 185, 450-5	2.5	9
111	Enhanced antitumor effect of combination intravesical mitomycin C and bacillus Calmette-Guerin therapy in an orthotopic bladder cancer model. <i>Oncology Letters</i> , 2011 , 2, 13-19	2.6	9
110	Regulation of human dendritic cells by a novel specific nuclear factor-kappaB inhibitor, dehydroxymethylepoxyquinomicin. <i>Human Immunology</i> , 2010 , 71, 763-70	2.3	9
109	Prognostic significance of high nuclear grade in patients with pathologic T1a renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 831-5	2.8	9
108	Therapeutic effect of intravesical administration of paclitaxel solubilized with poly(2-methacryloyloxyethyl phosphorylcholine-co-n-butyl methacrylate) in an orthotopic bladder cancer model. <i>BMC Cancer</i> , 2015 , 15, 317	4.8	8
107	History of Non-Muscle-Invasive Bladder Cancer May Have a Worse Prognostic Impact in cT2-4aN0M0 Bladder Cancer Patients Treated With Radical Cystectomy. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, e969-e976	3.3	8
106	Patient age was an independent predictor of cancer-specific survival in male patients with upper tract urothelial carcinoma treated by radical nephroureterectomy. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 554-559	2.8	8
105	Establishment of the optimal follow-up schedule after radical prostatectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 341.e9-341.e14	2.8	8
104	Long-Term Oncologic Outcomes of Laparoscopic Versus Open Radical Nephroureterectomy for Patients with T3N0M0 Upper Tract Urothelial Carcinoma: A Multicenter Cohort Study with Adjustment by Propensity Score Matching. <i>Annals of Surgical Oncology</i> , 2019 , 26, 3774-3781	3.1	8
103	Does laparoendoscopic single-site adrenalectomy increase surgical risk in patients with pheochromocytoma?. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2013 , 27, 593-8	5.2	8

102	Impact of smoking on the age at diagnosis of upper tract urothelial carcinoma: Subanalysis of the Japanese Urological Association multi-institutional national database. <i>International Journal of Urology</i> , 2015 , 22, 1023-7	2.3	8
101	Predicting the probability of significant prostate cancer in Japanese men with serum prostate-specific antigen less than 10 ng/mL: development of a novel pre-biopsy nomogram. <i>International Journal of Urology</i> , 2010 , 17, 274-80	2.3	8
100	Benefit of laparoscopic radical nephrectomy in patients with a high BMI. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 400-4	2.8	8
99	The benefit of laparoscopic partial nephrectomy in high body mass index patients. <i>Japanese Journal of Clinical Oncology</i> , 2012 , 42, 619-24	2.8	8
98	Carcinoma in situ developing in an ileal neobladder. <i>Urology</i> , 2007 , 69, 576.e9-11	1.6	8
97	Can urologists introduce the concept of "oligometastasis" for metastatic bladder cancer after total cystectomy?. <i>Oncotarget</i> , 2017 , 8, 111819-111835	3.3	8
96	Metastatic urothelial carcinoma to pericardia manifested by dyspnea from cardiac tamponade during systemic chemotherapy: Case report and literature review. <i>Canadian Urological Association Journal</i> , 2012 , 6, E184-8	1.2	8
95	Bacillus Calmette-Guérin-unresponsive non-muscle-invasive bladder cancer: Its definition and future therapeutic strategies. <i>International Journal of Urology</i> , 2020 , 27, 108-116	2.3	8
94	Purified protein derivative skin test reactions are associated with clinical outcomes of patients with nonmuscle invasive bladder cancer treated with induction bacillus Calmette-Guérin therapy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 77.e15-77.e21	2.8	8
93	No significant impact of patient age and prior treatment profile with docetaxel on the efficacy of cabazitaxel in patient with castration-resistant prostate cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2018 , 82, 1061-1066	3.5	8
92	Impact of lymph node dissection on clinical outcomes during nephroureterectomy in patients with clinically node-negative upper urinary tract urothelial cancer: subanalysis of a multi-institutional nationwide case series of the Japanese Urological Association. <i>Japanese Journal of Clinical Oncology</i> , 2017 , 47, 652-659	2.8	7
91	PKM2 under hypoxic environment causes resistance to mTOR inhibitor in human castration resistant prostate cancer. <i>Oncotarget</i> , 2018 , 9, 27698-27707	3.3	7
90	TNFAIP2 expression induces epithelial-to-mesenchymal transition and confers platinum resistance in urothelial cancer cells. <i>Laboratory Investigation</i> , 2019 , 99, 1702-1713	5.9	7
89	Radiological findings of primary localized amyloidosis of the ureter. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 35, 431-5	5.6	7
88	What is the predictor of prolonged operative time during laparoscopic radical prostatectomy?. <i>International Journal of Urology</i> , 2013 , 20, 330-6	2.3	7
87	How can lymphocele development be prevented after laparoscopic radical prostatectomy?. <i>Journal of Endourology</i> , 2013 , 27, 447-51	2.7	7
86	Periprostatic fat area is an independent factor that prolonged operative time in laparoscopic radical prostatectomy. <i>Urology</i> , 2013 , 82, 1304-9	1.6	7
85	Prognostic factors of recurrent disease in upper urinary tract urothelial cancer after radical nephroureterectomy: Subanalysis of the multi-institutional national database of the Japanese Urological Association. <i>International Journal of Urology</i> , 2015 , 22, 1013-20	2.3	7

84	Cancer death from non-muscle invasive bladder cancer: report of the Japanese Urological Association of data from the 1999-2001 registry in Japan. <i>International Journal of Urology</i> , 2010 , 17, 905-912	2.3	7
83	Three-dimensional ultrasonography in evaluation of benign prostatic hyperplasia. <i>International Journal of Urology</i> , 2004 , 11, 1087-91	2.3	7
82	Obstructive sleep apnea syndrome should be considered as a cause of nocturia in younger patients without other voiding symptoms. <i>Canadian Urological Association Journal</i> , 2016 , 10, E241-E245	1.2	7
81	Impact of multimodal treatment on prognosis for patients with metastatic upper urinary tract urothelial cancer: Subanalysis of the multi-institutional nationwide case series study of the Japanese Urological Association. <i>International Journal of Urology</i> , 2016 , 23, 224-30	2.3	7
80	The learning curve of laparoendoscopic single-site adrenalectomy: an analysis of over 100 cases. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2017 , 31, 170-177	5.2	6
79	A Case of Gemcitabine and Cisplatin Chemotherapy in a Patient With Metastatic Urothelial Carcinoma Receiving Hemodialysis. <i>Clinical Genitourinary Cancer</i> , 2016 , 14, e413-6	3.3	6
78	Frequency and prognostic significance of incidental prostate cancer at radical cystectomy: Results from an international retrospective study. <i>European Journal of Surgical Oncology</i> , 2017 , 43, 2193-2199	3.6	6
77	Vasohibin-1 as a novel microenvironmental biomarker for patient risk reclassification in low-risk prostate cancer. <i>Oncotarget</i> , 2018 , 9, 10203-10210	3.3	6
76	Evaluation of prostate-specific antigen density in the diagnosis of prostate cancer combined with magnetic resonance imaging before biopsy in men aged 70 years and older with elevated PSA. <i>Molecular and Clinical Oncology</i> , 2018 , 9, 656-660	1.6	6
75	Significance of a frozen section analysis of the ureteral margin in bladder cancer patients treated with radical cystectomy and neoadjuvant chemotherapy. <i>Medical Oncology</i> , 2017 , 34, 187	3.7	5
74	Lymphovascular invasion status at transurethral resection of bladder tumors may predict subsequent poor response of T1 tumors to bacillus Calmette-Guérin. <i>BMC Urology</i> , 2016 , 16, 5	2.2	5
73	A prospective longitudinal survey of erectile function status in symptomatic benign prostatic hyperplasia patients treated with dutasteride. <i>Aging Male</i> , 2016 , 19, 111-6	2.1	5
72	Performance of diffusion-weighted MRI post-CT urography for the diagnosis of upper tract urothelial carcinoma: Comparison with selective urine cytology sampling. <i>Clinical Imaging</i> , 2018 , 52, 208-215	2.7	5
71	Limited in vitro efficacy of CYP17A1 inhibition on human castration resistant prostate cancer. <i>Steroids</i> , 2014 , 92, 39-44	2.8	5
70	Low-dose docetaxel enhances the sensitivity of S-1 in a xenograft model of human castration resistant prostate cancer. <i>International Journal of Cancer</i> , 2012 , 130, 431-42	7.5	5
69	Therapeutic enhancement of S-1 with CPT-11 through down-regulation of thymidylate synthase in bladder cancer. <i>Cancer Medicine</i> , 2013 , 2, 488-95	4.8	5
68	Renal damage inhibited in mice lacking angiotensinogen gene subjected to unilateral ureteral obstruction. <i>Urology</i> , 2009 , 74, 938-43	1.6	5
67	The clinical impact of the classification of carcinoma in situ on tumor recurrence and their clinical course in patients with bladder tumor. <i>Japanese Journal of Clinical Oncology</i> , 2011 , 41, 424-9	2.8	5

66	Clinical impact of bladder biopsies with TUR-BT according to cytology results in patients with bladder cancer: a case control study. <i>BMC Urology</i> , 2010 , 10, 12	2.2	5
65	Prostate specific antigen adjusted for transition zone epithelial volume: the powerful predictor for the detection of prostate cancer on repeat biopsy. <i>Journal of Urology</i> , 2005 , 173, 1541-5	2.5	5
64	Usefulness of alpha1-antichymotrypsin-PSA complex for predicting bone metastases of prostate cancer. <i>Urology</i> , 2006 , 68, 371-5	1.6	5
63	Enhancement of Diethylstilbestrol Induced Cytotoxicity by Bcl-2 Antisense Oligodeoxynucleotides and A Glutathione Depletor for Prostate Cancer. <i>Journal of Urology</i> , 2003 , 169, 730-734	2.5	5
62	Percutaneous endoscopic marsupialization of a pyelocaliceal diverticulum with milk of calcium stones. <i>Urologia Internationalis</i> , 1998 , 60, 62-5	1.9	5
61	Role of the MUC1-C oncoprotein in the acquisition of cisplatin resistance by urothelial carcinoma. <i>Cancer Science</i> , 2020 , 111, 3639-3652	6.9	5
60	Are the Formulas Used to Estimate Renal Function Adequate for Patients Treated With Cisplatin-Based Chemotherapy After Nephroureterectomy for Upper Tract Urothelial Carcinoma?. <i>Clinical Genitourinary Cancer</i> , 2016 , 14, e501-e507	3.3	5
59	Urinary pH Levels are Strongly Associated with Bladder Recurrence After Nephroureterectomy in Upper Tract Urothelial Carcinoma Patients with a Smoking History. <i>Annals of Surgical Oncology</i> , 2016 , 23, 1029-1038	3.1	4
58	Optimal timing of hormonal therapy for prostate-specific antigen recurrence after radical prostatectomy. <i>Medical Oncology</i> , 2014 , 31, 45	3.7	4
57	Prognostic Model for Predicting Survival in Patients with Disease Recurrence Following Radical Cystectomy. <i>European Urology Focus</i> , 2015 , 1, 75-81	5.1	4
56	A case of spinal and bulbar muscular atrophy with high-stage and high-Gleason Score prostate cancer responded to maximal androgen blockade therapy. <i>Journal of Andrology</i> , 2012 , 33, 563-5		4
55	Are predictive models for cancer volume clinically useful in localized prostate cancer?. <i>International Journal of Urology</i> , 2009 , 16, 936-40	2.3	4
54	Castration-resistant prostate cancer patients who had poor response on first androgen deprivation therapy would obtain certain clinical benefit from early docetaxel administration. <i>International Journal of Clinical Oncology</i> , 2019 , 24, 546-553	4.2	4
53	Evaluating the Oncological Outcomes of Pure Laparoscopic Radical Nephroureterectomy Performed for Upper-Tract Urothelial Carcinoma Patients: A Multicenter Cohort Study Adjusted by Propensity Score Matching. <i>Annals of Surgical Oncology</i> , 2021 , 28, 465-473	3.1	4
52	Does neoadjuvant chemotherapy have therapeutic benefit for node-positive upper tract urothelial carcinoma? Results of a multi-center cohort study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021 , 40, 105.e19-105.e19	2.8	4
51	The clinicopathological characteristics of muscle-invasive bladder recurrence in upper tract urothelial carcinoma. <i>Cancer Science</i> , 2021 , 112, 1084-1094	6.9	4
50	Characterizing intermediate-risk non-muscle-invasive bladder cancer: Implications for the definition of intermediate risk and treatment strategy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017 , 35, 208-214	2.8	3
49	Spontaneous Regression of Epstein-Barr Virus-Positive Primary Diffuse Large Cell B-Cell Lymphoma of the Urinary Bladder After the Cessation of Enzalutamide. <i>Clinical Genitourinary Cancer</i> , 2016 , 14, e215-28 ³		3

48	Novel algorithm for management of acute epididymitis. <i>International Journal of Urology</i> , 2017 , 24, 82-87	2.3	3
47	Visceral to total obesity ratio and severe hydronephrosis are independently associated with prolonged pneumoperitoneum operative time in patients undergoing laparoscopic radical nephroureterectomy for upper tract urothelial carcinoma. <i>SpringerPlus</i> , 2015 , 4, 290		3
46	Efficacy of treatment with a GnRH antagonist in prostate cancer patients previously treated with a GnRH agonist. <i>Cancer Chemotherapy and Pharmacology</i> , 2015 , 76, 301-6	3.5	3
45	Transumbilical laparoendoscopic single-site partial nephrectomy using a microwave tissue coagulator. <i>International Cancer Conference Journal</i> , 2012 , 1, 176-179	0.9	3
44	Clinical implications of tumor size and local extent of primary prostatic lesions in prostate cancer patients with metastases: value of endorectal magnetic resonance imaging in patients with metastases. <i>Urology</i> , 2007 , 70, 86-90	1.6	3
43	Potent increased risk of the initiation of DNA replication in human prostate cancer with the use of 5β-reductase inhibitors. <i>American Journal of Clinical and Experimental Urology</i> , 2014 , 2, 136-44	1.6	3
42	Role of Previous Malignancy History in Clinical Outcomes in Patients with Initially Diagnosed Non-Muscle Invasive Bladder Cancer. <i>Annals of Surgical Oncology</i> , 2021 , 28, 5349-5359	3.1	3
41	A new bioinformatics approach identifies overexpression of GRB2 as a poor prognostic biomarker for prostate cancer. <i>Scientific Reports</i> , 2021 , 11, 5696	4.9	3
40	Should We Try Antiandrogen Withdrawal in Castration-Resistant Prostate Cancer Patients? Insights From a Retrospective Study. <i>Clinical Genitourinary Cancer</i> , 2016 , 14, e569-e573	3.3	3
39	A Novel Risk-based Approach Simulating Oncological Surveillance After Radical Nephroureterectomy in Patients with Upper Tract Urothelial Carcinoma. <i>European Urology Oncology</i> , 2020 , 3, 756-763	6.7	3
38	Ureteral pseudodiverticulosis accompanied by urothelial carcinoma diagnosed by CT urography: a case report and review of the literature. <i>BJR case Reports</i> , 2018 , 4, 20170111	0.7	3
37	Prognostic significance of erythrocyte protein band 4.1-like5 expression in upper urinary tract urothelial carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017 , 35, 543.e17-543.e24	2.8	2
36	Appropriate timing for a biochemical evaluation after adrenalectomy for unilateral aldosterone-producing adenoma. <i>Clinical Endocrinology</i> , 2020 , 92, 503-508	3.4	2
35	Does switching the bacillus Calmette-Guérin strain affect clinical outcome in patients with recurrent non-muscle-invasive bladder cancer after initial bacillus Calmette-Guérin therapy?. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 306.e1-306.e8	2.8	2
34	Syndrome of inappropriate antidiuretic hormone secretion in a patient with castration-resistant prostate cancer treated with enzalutamide. <i>Acta Oncologica</i> , 2016 , 55, 399-400	3.2	2
33	Factors influencing the operating time for single-port laparoscopic radical nephrectomy: focus on the anatomy and distribution of the renal artery and vein. <i>Japanese Journal of Clinical Oncology</i> , 2017 , 47, 976-980	2.8	2
32	Androgen deprivation therapy improves pneumoperitoneum time during laparoscopic radical prostatectomy in Japanese patients with enlarged prostates. <i>Asian Journal of Endoscopic Surgery</i> , 2014 , 7, 145-51	1.4	2
31	Three cases of granulocyte colony-stimulating factor-producing urinary upper tract carcinomas. <i>Scandinavian Journal of Urology and Nephrology</i> , 2012 , 46, 136-41		2

30	Prognostic stratification in patients who received hormonal therapy for prostate-specific antigen recurrence after radical prostatectomy. <i>Japanese Journal of Clinical Oncology</i> , 2010 , 40, 177-80	2.8	2
29	Brain metastasis of a papillary renal cell carcinoma, identified as type 2. <i>International Journal of Clinical Oncology</i> , 2009 , 14, 270-2	4.2	2
28	Non-muscle invasive bladder cancer with concomitant vaginal urothelial carcinoma: a case report and review of the literature. <i>International Journal of Clinical Oncology</i> , 2010 , 15, 626-30	4.2	2
27	A comparison of perioperative outcomes of laparoscopic versus open nephroureterectomy for upper tract urothelial carcinoma: a propensity score matching analysis. <i>Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology</i> , 2021 ,	4.4	2
26	Relationship between radiation doses and erectile function deterioration in patients with localized prostate cancer treated with permanent prostate brachytherapy. <i>International Journal of Urology</i> , 2020 , 27, 1087-1093	2.3	2
25	Diversity in treatment modalities of Stage II/III urothelial cancer in Japan: sub-analysis of the multi-institutional national database of the Japanese Urological Association. <i>Japanese Journal of Clinical Oncology</i> , 2016 , 46, 468-74	2.8	2
24	Natural course of asymptomatic abnormal prostate findings incidentally detected by CT after intravesical BCG therapy. <i>International Journal of Clinical Oncology</i> , 2017 , 22, 554-562	4.2	1
23	A comparison of nephrotoxicity between patients with a solitary-functioning kidney and those with bilateral-functioning kidneys in cisplatin-based chemotherapy for advanced urothelial carcinoma: a Japanese retrospective multi-institutional study. <i>BMC Cancer</i> , 2018 , 18, 290	4.8	1
22	Influence of physician years on urological journal publication productivity among Japanese urologists. <i>SpringerPlus</i> , 2016 , 5, 2040		1
21	Clinical Background of Patients with Sperm in Their Urinary Sediment. <i>PLoS ONE</i> , 2015 , 10, e0136844	3.7	1
20	A comparison of perioperative outcomes of laparoscopic versus open nephroureterectomy for upper tract urothelial carcinoma: a propensity score matching analysis. <i>Minerva Urology and Nephrology</i> , 2022 , 74,	2.3	1
19	Urinary pH is an independent predictor of upper tract recurrence in non-muscle-invasive bladder cancer patients with a smoking history. <i>Scientific Reports</i> , 2021 , 11, 20675	4.9	1
18	ASO Author Reflections: Previous History of Non-urothelial Malignancy May Provide Predictive Information of Worse Clinical Outcome for Initially Diagnosed Non-Muscle Invasive Bladder Cancer. <i>Annals of Surgical Oncology</i> , 2021 , 28, 5360-5361	3.1	1
17	Guideline adherence for radical cystectomy significantly affects survival outcomes in non-muscle-invasive bladder cancer patients. <i>Japanese Journal of Clinical Oncology</i> , 2021 , 51, 1303-1312	2.8	1
16	Erectile function status is highly associated with prostate-specific antigen bounce in localized prostate cancer patients treated with permanent prostate brachytherapy. <i>International Journal of Urology</i> , 2016 , 23, 247-52	2.3	1
15	Can random bladder biopsies be eliminated after bacillus Calmette-Guérin therapy against carcinoma in situ?. <i>International Urology and Nephrology</i> , 2021 , 53, 465-469	2.3	1
14	Significance of prophylactic urethrectomy at the time of radical cystectomy for bladder cancer. <i>Japanese Journal of Clinical Oncology</i> , 2021 , 51, 287-295	2.8	1
13	Individualized prostate-specific antigen threshold values to avoid overdiagnosis of prostate cancer and reduce unnecessary biopsy in elderly men. <i>Japanese Journal of Clinical Oncology</i> , 2014 , 44, 852-9	2.8	0

12	Renal cell carcinoma arising from a horseshoe kidney in a chronic hemodialysis patient. <i>Clinical and Experimental Nephrology</i> , 2012 , 16, 647-51	2.5	o
11	Oncological outcomes of dose reductions in cisplatin due to renal dysfunction for patients with metastatic urothelial carcinoma.. <i>BJUJ Compass</i> , 2021 , 2, 322-330	0.9	o
10	Effects of transurethral resection under general anesthesia on tumor recurrence in non-muscle invasive bladder cancer. <i>International Journal of Clinical Oncology</i> , 2021 , 26, 2094-2103	4.2	o
9	Umbilical closure using 2-octyl cyanoacrylate in transumbilical laparoscopic adrenalectomy: A randomized controlled trial. <i>International Journal of Urology</i> , 2020 , 27, 670-675	2.3	
8	Current status of systemic chemotherapy for octogenarians with advanced urothelial cancer in Japan: a Japanese multi-institutional study (CURE study). <i>International Journal of Clinical Oncology</i> , 2016 , 21, 1142-1149	4.2	
7	Editorial comment to template-based lymphadenectomy in urothelial carcinoma of the renal pelvis: a prospective study. <i>International Journal of Urology</i> , 2014 , 21, 460	2.3	
6	Expression level of dihydropyrimidine dehydrogenase is associated with clinical outcome in patients with T1G3 bladder cancer treated with Bacillus Calmette-Guerin. <i>BMC Research Notes</i> , 2014 , 7, 646	2.3	
5	RE: Prognostic significance of Bacillus Calmette-Guérin failure classification in non-muscle-invasive bladder cancer. <i>BJU International</i> , 2012 , 110, E164	5.6	
4	Renal cell carcinoma accompanied by left-sided IVC. <i>Internal Medicine</i> , 2011 , 50, 1127	1.1	
3	CP8-9 NF-B function inhibitor, DHMEQ as a treatment option for advanced urological cancer. <i>Japanese Journal of Urology</i> , 2004 , 95, 163		
2	Site-specific differences in survival among upper and lower tract urothelial carcinoma patients treated with radical surgery. <i>Japanese Journal of Clinical Oncology</i> , 2021 , 51, 984-991	2.8	
1	Pre-operative and post-operative estimated glomerular filtration rate in Japanese patients with urological malignancies. <i>Clinical and Experimental Nephrology</i> , 2018 , 22, 1205-1212	2.5	