

Marion M Simmons

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1134259/publications.pdf>

Version: 2024-02-01

13
papers

464
citations

1163117

8
h-index

1125743

13
g-index

13
all docs

13
docs citations

13
times ranked

491
citing authors

#	ARTICLE	IF	CITATIONS
1	Evaluation of an alternative method for production of biodiesel from processed fats derived from Category 1, 2 and 3 animal by-products (submitted by College Proteins). <i>EFSA Journal</i> , 2020, 18, e06089.	1.8	3
2	Inactivation of H-type and L-type bovine spongiform encephalopathy following recommended autoclave decontamination procedures. <i>Transboundary and Emerging Diseases</i> , 2020, 67, 1872.	3.0	1
3	The Scrapie Prevalence in a Goat Herd Is Underestimated by Using a Rapid Diagnostic Test. <i>Frontiers in Bioengineering and Biotechnology</i> , 2020, 8, 164.	4.1	8
4	Incomplete inactivation of atypical scrapie following recommended autoclave decontamination procedures. <i>Transboundary and Emerging Diseases</i> , 2019, 66, 1993-2001.	3.0	7
5	DISCONTTOOLS: Identifying gaps in controlling bovine spongiform encephalopathy. <i>Transboundary and Emerging Diseases</i> , 2018, 65, 9-21.	3.0	6
6	Abnormalities in Brainstem Auditory Evoked Potentials in Sheep with Transmissible Spongiform Encephalopathies and Lack of a Clear Pathological Relationship. <i>Frontiers in Veterinary Science</i> , 2016, 3, 60.	2.2	2
7	Evidence of scrapie transmission to sheep via goat milk. <i>BMC Veterinary Research</i> , 2016, 12, 208.	1.9	21
8	First case of chronic wasting disease in Europe in a Norwegian free-ranging reindeer. <i>Veterinary Research</i> , 2016, 47, 88.	3.0	244
9	Whole Blood Gene Expression Profiling in Preclinical and Clinical Cattle Infected with Atypical Bovine Spongiform Encephalopathy. <i>PLoS ONE</i> , 2016, 11, e0153425.	2.5	10
10	The pathological and molecular but not clinical phenotypes are maintained after second passage of experimental atypical bovine spongiform encephalopathy in cattle. <i>BMC Veterinary Research</i> , 2014, 10, 243.	1.9	15
11	Chronic wasting disease and atypical forms of bovine spongiform encephalopathy and scrapie are not transmissible to mice expressing wild-type levels of human prion protein. <i>Journal of General Virology</i> , 2012, 93, 1624-1629.	2.9	78
12	Experimental H-type and L-type bovine spongiform encephalopathy in cattle: observation of two clinical syndromes and diagnostic challenges. <i>BMC Veterinary Research</i> , 2012, 8, 22.	1.9	56
13	The interpretation of disease phenotypes to identify TSE strains in mice: characterisation of BSE using PrP ^{Sc} distribution patterns in the brain. <i>Veterinary Research</i> , 2012, 43, 86.	3.0	13