

Antti Koskela

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1135090/publications.pdf>

Version: 2024-02-01

26
papers

928
citations

567144

15
h-index

610775

24
g-index

27
all docs

27
docs citations

27
times ranked

1597
citing authors

#	ARTICLE	IF	CITATIONS
1	Osteoblast-derived WNT16 represses osteoclastogenesis and prevents cortical bone fragility fractures. <i>Nature Medicine</i> , 2014, 20, 1279-1288.	15.2	303
2	Estrogen receptor- β in osteocytes is important for trabecular bone formation in male mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 2294-2299.	3.3	118
3	Perfluoroalkyl substances, bone density, and cardio-metabolic risk factors in obese 8-12 year old children: A pilot study. <i>Environmental Research</i> , 2018, 160, 314-321.	3.7	77
4	The bone-sparing effects of estrogen and WNT16 are independent of each other. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 14972-14977.	3.3	50
5	Expression of the Hutchinson-Gilford Progeria Mutation during Osteoblast Development Results in Loss of Osteocytes, Irregular Mineralization, and Poor Biomechanical Properties. <i>Journal of Biological Chemistry</i> , 2012, 287, 33512-33522.	1.6	39
6	Estrogen receptor- β expression in neuronal cells affects bone mass. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 983-988.	3.3	37
7	Porcupine inhibitors impair trabecular and cortical bone mass and strength in mice. <i>Journal of Endocrinology</i> , 2018, 238, 13-23.	1.2	37
8	Inducible Wnt16 inactivation: WNT16 regulates cortical bone thickness in adult mice. <i>Journal of Endocrinology</i> , 2018, 237, 113-122.	1.2	32
9	Osteoblast-derived NOTUM reduces cortical bone mass in mice and the <i>NOTUM</i> locus is associated with bone mineral density in humans. <i>FASEB Journal</i> , 2019, 33, 11163-11179.	0.2	24
10	Synergistic effects of tributyltin and 2,3,7,8-tetrachlorodibenzo-p-dioxin on differentiating osteoblasts and osteoclasts. <i>Toxicology and Applied Pharmacology</i> , 2012, 263, 210-217.	1.3	23
11	The role of activation functions 1 and 2 of estrogen receptor- β for the effects of estradiol and selective estrogen receptor modulators in male mice. <i>Journal of Bone and Mineral Research</i> , 2013, 28, 1117-1126.	3.1	23
12	Transgene silencing of the Hutchinson-Gilford progeria syndrome mutation results in a reversible bone phenotype, whereas resveratrol treatment does not show overall beneficial effects. <i>FASEB Journal</i> , 2015, 29, 3193-3205.	0.2	21
13	Enzalutamide Reduces the Bone Mass in the Axial But Not the Appendicular Skeleton in Male Mice. <i>Endocrinology</i> , 2016, 157, 969-977.	1.4	20
14	SERMs have substance-specific effects on bone, and these effects are mediated via ER- β -AF-1 in female mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 310, E912-E918.	1.8	20
15	RSPO3 is important for trabecular bone and fracture risk in mice and humans. <i>Nature Communications</i> , 2021, 12, 4923.	5.8	19
16	The androgen receptor is required for maintenance of bone mass in adult male mice. <i>Molecular and Cellular Endocrinology</i> , 2019, 479, 159-169.	1.6	19
17	Clinically relevant doses of vitamin A decrease cortical bone mass in mice. <i>Journal of Endocrinology</i> , 2018, 239, 389-402.	1.2	17
18	Liver-derived IGF-I regulates cortical bone mass but is dispensable for the osteogenic response to mechanical loading in female mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 311, E138-E144.	1.8	12

#	ARTICLE	IF	CITATIONS
19	Endocrine, metabolic and apical effects of in utero and lactational exposure to non-dioxin-like 2,2,3,4,4,5,5-heptachlorobiphenyl (PCB 180): A postnatal follow-up study in rats. <i>Reproductive Toxicology</i> , 2021, 102, 109-127.	1.3	8
20	Androgen receptor SUMOylation regulates bone mass in male mice. <i>Molecular and Cellular Endocrinology</i> , 2019, 479, 117-122.	1.6	7
21	Osteocyte- and late osteoblast-derived NOTUM reduces cortical bone mass in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2021, 320, E967-E975.	1.8	6
22	The Bone Sparing Effects of 2-Methoxyestradiol Are Mediated via Estrogen Receptor- α in Male Mice. <i>Endocrinology</i> , 2016, 157, 4200-4205.	1.4	5
23	Acute fat loss does not affect bone mass. <i>Scientific Reports</i> , 2021, 11, 14177.	1.6	5
24	Perfluoroalkyl Substances and Abdominal Aortic Calcification. <i>Journal of Occupational and Environmental Medicine</i> , 2022, 64, 287-294.	0.9	3
25	Maternal beef and postweaning herring diets increase bone mineral density and strength in mouse offspring. <i>Experimental Biology and Medicine</i> , 2013, 238, 1362-1369.	1.1	2
26	Estradiol and RSPO3 regulate vertebral trabecular bone mass independent of each other. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2022, , .	1.8	1