
Robin Whelpton

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11353643/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The binding of physostigmine to human serum albumin. Journal of Pharmacy and Pharmacology, 2011, 42, 804-805.	2.4	9
2	lonization constants, octanol partition coefficients and Cholinesterase inhibitor constants for chlorpromazine and its metabolites. Journal of Pharmacy and Pharmacology, 2011, 41, 856-858.	2.4	6
3	Chromatographic separation of thioridazine sulphoxide and N-oxide diastereoisomers: identification as metabolites in the rat. Journal of Pharmacy and Pharmacology, 2011, 38, 506-509.	2.4	15
4	Kinetics of Metabolism and Excretion. , 2010, , 121-136.		0
5	Substance P modulation of striatal dopamine outflow is determined by M1 and M2 muscarinic receptors in male Wistar rats. Neuroscience Letters, 2000, 293, 179-182.	2.1	11
6	Double-blind, randomized study of the effect of cisapride on gastric emptying in critically ill patients. Critical Care Medicine, 1997, 25, 447-451.	0.9	53
7	Evidence for modulatory effects of substance P fragments (1–4) and (8–11) on endogenous dopamine outflow in rat striatal slices. Neuroscience Letters, 1996, 205, 33-36.	2.1	16
8	Distribution and Excretion of Radiolabeled Temoporfin in a Murine Tumor Model. Photochemistry and Photobiology, 1996, 63, 885-891.	2.5	35
9	DISTRIBUTION OF TEMOPORFIN, A NEW PHOTOSENSITIZER FOR THE PHOTODYNAMIC THERAPY OF CANCER, IN A MURINE TUMOR MODEL. Photochemistry and Photobiology, 1995, 61, 397-401.	2.5	65
10	Substance P-(1–7) and substance P-(5–11) locally modulate dopamine release in rat striatum. European Journal of Pharmacology, 1995, 282, 229-233.	3.5	25
11	Biphasic inhibition of stimulated endogenous dopamine release by 7â€OHâ€DPAT in slices of rat nucleus accumbens. British Journal of Pharmacology, 1995, 115, 421-426.	5.4	24
12	Determination of paracetamol (acetaminophen) in blood and plasma using high performance liquid chromatography with dual electrode coulometric quantification in the redox mode. Biomedical Chromatography, 1993, 7, 90-93.	1.7	20
13	Liquid chromatographic determination of hyoscine (scopolamine) in urine using solid phase extraction. Biomedical Chromatography, 1992, 6, 198-204.	1.7	7
14	Determination of plasma nefopam by liquid chromatography and electrochemical detection. Biomedical Applications, 1990, 526, 159-168.	1.7	10
15	Determination of prilocaine in human plasma samples using high-performance liquid chromatography with dual-electrode electrochemical detection. Biomedical Applications, 1990, 526, 215-222.	1.7	12
16	Iterative least squares fitting of pH-partition data. Trends in Pharmacological Sciences, 1989, 10, 182-183.	8.7	12
17	High-performance liquid chromatographic resolution of the enantiomers of thioridazine, its metabolites and related compounds. Biomedical Applications, 1988, 426, 223-228.	1.7	12

18 HPLC-EC Determination of Physostigmine in Biological Samples. , 1986, , 181-187.

0

ROBIN WHELPTON

#	Article	IF	CITATIONS
19	Sensitive liquid chromatographic method for physostigmine in biological fluids using dual-electrode electrochemical detection. Biomedical Applications, 1985, 341, 361-371.	1.7	29
20	PHYSOSTIGMINE CONCENTRATIONS AFTER ORAL DOSES. Lancet, The, 1985, 325, 695-696.	13.7	18
21	Isotope-Labelled Materials as Internal Standards. , 1984, , 39-44.		0
22	Analysis of plasma physostigmine concentrations by liquid chromatography. Biomedical Applications, 1983, 272, 216-220.	1.7	43
23	Analysis of plasma trifluoperazine by gas chromatography and selected ion monitoring. Biomedical Applications, 1982, 228, 321-326.	1.7	13
24	Basic Laboratory Operations. , 0, , 353-398.		0
25	Sample Preparation. , 0, , 49-94.		Ο