

Francisco Langeani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1137703/publications.pdf>

Version: 2024-02-01

64
papers

1,278
citations

516561

16
h-index

377752

34
g-index

66
all docs

66
docs citations

66
times ranked

1149
citing authors

#	ARTICLE	IF	CITATIONS
1	Gross brain morphology of Hypoptopomatinae and Neoplecostominae (Siluriformes: Loricariidae): Comparative anatomy and phylogenetic implications. <i>Journal of Zoological Systematics and Evolutionary Research</i> , 2021, 59, 111-147.	0.6	2
2	Overlooked biodiversity of mitochondrial lineages in <i>Hemiodus</i> (Ostariophysi, Characiformes). <i>Zoologica Scripta</i> , 2021, 50, 337-351.	0.7	5
3	Two new species of <i>Hypostomus</i> suckermouth armoured catfishes (Teleostei: Loricariidae) from central Brazil. <i>Journal of Fish Biology</i> , 2021, 99, 905-920.	0.7	3
4	Molecular species delimitation of the genera <i>Anodus</i> , <i>Argonectes</i> , <i>Bivibranchia</i> and <i>Micromischodus</i> (Ostariophysi: Characiformes). <i>Neotropical Ichthyology</i> , 2021, 19, .	0.5	2
5	Stream fish from recently deforested basins in the Meridional Amazon, Mato Grosso, Brazil. <i>Biota Neotropica</i> , 2020, 20, .	0.2	3
6	DNA barcode and minibarcode identification of freshwater fishes from Cerrado headwater streams in Central Brazil. <i>Journal of Fish Biology</i> , 2019, 95, 1046-1060.	0.7	10
7	<i>Hemiodus bimaculatus</i> , a new species of Hemiodontidae from the Rio Tapaj�s drainage, Brazil (Ostariophysi: Characiformes). <i>Journal of Fish Biology</i> , 2019, 94, 798-803.	0.7	4
8	Poor taxonomic sampling undermines nomenclatural stability: A reply to Roxo et al. (2019). <i>Zootaxa</i> , 2019, 4701, zootaxa.4701.5.10.	0.2	4
9	Checklist and key for the identification of fish fauna of the Uberaba River, Upper Paran� River system, Brazil. <i>ZooKeys</i> , 2019, 875, 129-155.	0.5	6
10	A new species of <i>Astyanax</i> (Characiformes: Characidae) from Costa Rica and Panama, with a key to the lower Central American species of the genus. <i>Journal of Fish Biology</i> , 2018, 92, 1866-1887.	0.7	4
11	Redescription of <i>Trichomycterus striatus</i> (Meek & Hildebrand, 1913) (Siluriformes: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	0.2	7
12	<i>Microlepidogaster negomata</i> , a new hypoptopomatine catfish (Loricariidae: Hypoptopomatinae) from Rio S�o Francisco basin, southeastern Brazil. <i>Zootaxa</i> , 2017, 4276, 233-244.	0.2	1
13	Gross brain morphology of the armoured catfish <i>Rineloricaria heteroptera</i> (Loricariidae: Loricariinae): A descriptive and quantitative approach. <i>Journal of Morphology</i> , 2017, 278, 1689-1705.	0.6	8
14	Annotated catalog of the type specimens deposited at the fish collection of the Universidade Estadual Paulista, C�mpus de S�o Jos� do Rio Preto, S�o Paulo, Brazil (DZSJRP). <i>Zootaxa</i> , 2017, 4319, 435.	0.2	0
15	Redescription of <i>Hisonotus notatus</i> Eigenmann & Eigenmann, 1889 (Loricariidae: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 drainages of southeastern Brazil. <i>Neotropical Ichthyology</i> , 2016, 14, .	0.5	2
16	Reconciling more than 150 years of taxonomic confusion: The true identity of <i>Moenkhausia lepidura</i> , with a key to the species of the <i>M. lepidura</i> group (Characiformes: Characidae). <i>Zootaxa</i> , 2016, 4107, 338-52.	0.2	11
17	A new species of <i>Ituglanis</i> (Siluriformes: Trichomycteridae) from the Tocantins and Parana� river basins, central Brazil, with remarks on the systematics of the genus. <i>Zootaxa</i> , 2016, 4171, 439-458.	0.2	10
18	Photography-based taxonomy is inadequate, unnecessary, and potentially harmful for biological sciences. <i>Zootaxa</i> , 2016, 4196, zootaxa.4196.3.9.	0.2	63

#	ARTICLE	IF	CITATIONS
19	A new species of the catfish <i>Neoplecostomus</i> (Loricariidae: Neoplecostominae) from a coastal drainage in southeastern Brazil. <i>Neotropical Ichthyology</i> , 2016, 14, .	0.5	4
20	A new <i>Hasemania</i> Ellis from the upper rio Paran basin, with the redescription of <i>Hasemania crenuroides</i> Zarske & Gry (Characiformes: Characidae). <i>Neotropical Ichthyology</i> , 2015, 13, 479-486.	0.5	9
21	The stream fish fauna from three regions of the Upper Paran River basin. <i>Biota Neotropica</i> , 2015, 15, .	1.0	13
22	First record of <i>Crenicichla semifasciata</i> (Heckel, 1840) (Teleostei: Cichlidae) in the upper rio Paran basin, Brazil. <i>Check List</i> , 2015, 11, 1610.	0.1	3
23	Miniaturization in <i>Otothyris</i> Myers, 1927 (Loricariidae: Hypoptopomatinae). <i>Neotropical Ichthyology</i> , 2014, 12, 53-60.	0.5	12
24	Systematics of <i>Pseudotothyris</i> (Loricariidae: Hypoptopomatinae). <i>Zoological Journal of the Linnean Society</i> , 2014, 170, 822-874.	1.0	16
25	Redescription of <i>Hyphessobrycon flammeus</i> Myers, 1924 (Ostariophysi: Characidae), a threatened species from Brazil. <i>Neotropical Ichthyology</i> , 2014, 12, 247-256.	0.5	8
26	A new spiny species of <i>Hypostomus</i> Lacpde (Loricariidae: Hypostominae) from thermal waters, upper rio Paran basin, central Brazil. <i>Neotropical Ichthyology</i> , 2014, 12, 729-736.	0.5	7
27	<i>Microlepidogaster discus</i> , a new species of Hypoptopomatinae (Siluriformes: Loricariidae) from the rio Jequitinhonha basin, southeastern Brazil. <i>Zootaxa</i> , 2014, 3873, 560-70.	0.2	6
28	Systematics of <i>Pseudotothyris</i> (Loricariidae: Hypoptopomatinae). <i>Zoological Journal of the Linnean Society</i> , 2014, 170, 822-874.	1.0	12
29	A new species of <i>Neoplecostomus</i> Eigenmann & Eigenmann, 1888 (Siluriformes: Loricariidae:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	0.5	5
30	<i>Hemiodus iratapuru</i> , a new species of Hemiodontidae from the Rio Jari, Amazon Basin, Brazil (Teleostei, Characiformes). <i>Journal of Fish Biology</i> , 2013, 82, 1259-1268.	0.7	4
31	<i>Hyphessobrycon uaiso</i> : new characid fish from the rio Grande, upper rio Paran basin, Minas Gerais State (Ostariophysi: Characidae), with a brief comment about some types of <i>Hyphessobrycon</i> . <i>Neotropical Ichthyology</i> , 2013, 11, 525-536.	0.5	24
32	Spatial and seasonal patterns in fish assemblage in Corrego Rico, upper Parana River basin. <i>Neotropical Ichthyology</i> , 2013, 11, 143-152.	0.5	8
33	<i>Harttia absaberi</i> , a new species of loricariid catfish (Siluriformes: Loricariidae: Loricariinae) from the upper rio Paran basin, Brazil. <i>Neotropical Ichthyology</i> , 2013, 11, 779-786.	0.5	6
34	<i>Microlepidogaster arachs</i> , a new species of hypoptopomatine catfish (Siluriformes: Loricariidae) from the upper rio Paran basin, Brazil. <i>Zootaxa</i> , 2013, 3608, 379-88.	0.2	11
35	A New Species of <i>Hypostomus</i> (Siluriformes: Loricariidae) from the Upper Rio Paraguay Basin, Brazil. <i>Copeia</i> , 2012, 2012, 494-500.	1.4	4
36	A ictiofauna de cabeceiras do rio Tiet, So Paulo, Brasil. <i>Biota Neotropica</i> , 2011, 11, 217-236.	1.0	10

#	ARTICLE	IF	CITATIONS
37	Rhinolekos, a new genus with three new species of Hypoptopomatinae (Siluriformes: Loricariidae) from upper rio Paran�. Neotropical Ichthyology, 2011, 9, 65-78.	0.5	13
38	Microlepidogaster dimorpha, a new species of Hypoptopomatinae (Siluriformes: Loricariidae) from the upper rio Paran� system. Neotropical Ichthyology, 2011, 9, 79-86.	0.5	19
39	<i>Moenkhausia celibela</i> : a new species from the Amazon basin, Brazil (Characiformes: Characidae). Journal of Fish Biology, 2010, 77, 879-889.	0.7	8
40	A new species of Moenkhausia from the rio Amazonas and rio Orinoco basins (Characiformes: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 622	0.2	9
41	Coptobrycon bilineatus (Ellis, 1911) (Characiformes: Characidae): redescription and comments on its phylogenetic relationships. Neotropical Ichthyology, 2010, 8, 727-736.	0.5	5
42	Color pattern variation in Trichomycterus iheringi (Eigenmann, 1917) (Siluriformes: Trichomycteridae) from rio Itatinga and rio Claro, S�o Paulo, Brazil. Neotropical Ichthyology, 2010, 8, 49-56.	0.5	14
43	Redescription of Astyanax goyacensis Eigenmann, 1908 (Ostariophysi: Characiformes: Characidae). Neotropical Ichthyology, 2009, 7, 371-376.	0.5	19
44	Burrowing behavior of Dermatonotus muelleri (Anura, Microhylidae) with reference to the origin of the burrowing behavior of Anura. Journal of Ethology, 2009, 27, 195-201.	0.4	41
45	A fish-based biotic integrity index for assessment of lowland streams in southeastern Brazil. Hydrobiologia, 2009, 623, 173-189.	1.0	91
46	Characidium xanthopterus (Ostariophysi: Characiformes: Crenuchidae): a new species from the Central Brazilian Plateau. Neotropical Ichthyology, 2008, 6, 169-174.	0.5	16
47	A new Hemigrammus Gill from upper rio Parana system, Southeastern Brazil (Characiformes: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 7	0.2	7
48	Neoplecostomus (Teleostei: Loricariidae) from the upper Rio Parana basin, Brazil, with description of three new species. Zootaxa, 2008, 1757, 31.	0.2	27
49	Hypheobrycon rutiliflavus n. sp., a new characid fish from the upper rio Paraguai, State of Mato Grosso, Brazil (Characiformes: Characidae). Zootaxa, 2008, 1674, 39.	0.2	7
50	Diversidade da ictiofauna do Alto Rio Paran�: composi�o atual e perspectivas futuras. Biota Neotropica, 2007, 7, 181-197.	1.0	213
51	Ichthyofauna of the rio Itatinga in the Parque das Neblinas, Bertioga, S�o Paulo State: composition and biogeography. Biota Neotropica, 2007, 7, 81-86.	1.0	31
52	Fish, Hasemanina crenuchoides Zarske and �ry, 1999 (Ostariophysi: Characiformes: Characidae): rediscovery and distribution extension in the upper rio Paran� system, Minas Gerais, Brazil. Check List, 2007, 3, 119.	0.1	6
53	Redescr�o e osteologia de Bryconamericus exodon Eigenmann, 1907 (Ostariophysi, Characiformes,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 8	1.0	8
54	Stream fish, water and habitat quality in a pasture dominated basin, southeastern Brazil. Brazilian Journal of Biology, 2006, 66, 681-696.	0.4	106

#	ARTICLE	IF	CITATIONS
55	Effects of Physical Habitat Degradation on the Stream Fish Assemblage Structure in a Pasture Region. <i>Environmental Management</i> , 2006, 38, 974-982.	1.2	90
56	<i>Bryconamericus turiuba</i> , a New Species from the Upper Rio Paraná System (Ostariophysi). <i>Journal of Fish Biology</i> , 2006, 69, 1011-1022.	1.4	16
57	Riffle and pool fish communities in a large stream of southeastern Brazil. <i>Neotropical Ichthyology</i> , 2005, 3, 305-311.	0.5	41
58	<i>Hemiodus jatuarana</i> , a new species of Hemiodontidae from the rio Trombetas, Amazon Basin, Brazil (Teleostei, Characiformes). <i>Zootaxa</i> , 2004, 546, 1-6.	0.2	6
59	Estrutura e composição da ictiofauna de riachos da bacia do Rio Grande no estado de São Paulo, sudeste do Brasil. <i>Biota Neotropica</i> , 2004, 4, 01-39.	1.0	54
60	Geographic polymorphism of P element in populations of <i>Drosophila sturtevantii</i> . <i>Genetics and Molecular Biology</i> , 2003, 26, 175-179.	0.6	1
61	New Species of <i>Harttia</i> (Loricariidae, Loricariinae) from the Rio São Francisco Basin. <i>Copeia</i> , 2001, 2001, 136-142.	1.4	4
62	Peixes de riacho do parque estadual Morro do Diabo, Bacia do Alto Rio Paraná, SP. <i>Biota Neotropica</i> , 2001, 1, 1-15.	1.0	93
63	New Species of <i>Hemiodus</i> (Ostariophysi, Characiformes, Hemiodontidae) from the Rio Tocantins, Brazil, with Comments on Color Patterns and Tooth Shapes within the Species and Genus. <i>Copeia</i> , 1999, 1999, 718.	1.4	7
64	<i>Pimelodus paranaensis</i> , sp. n., um novo Pimelodidae (Pisces, Siluriformes) do Alto Paraná, Brasil. <i>Revista Brasileira De Zoologia</i> , 1988, 5, 409-417.	0.5	19