

Susmita Ghosh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11385596/publications.pdf>

Version: 2024-02-01

12
papers

437
citations

840776

11
h-index

1199594

12
g-index

12
all docs

12
docs citations

12
times ranked

815
citing authors

#	ARTICLE	IF	CITATIONS
1	AIM1 is an actin-binding protein that suppresses cell migration and micrometastatic dissemination. <i>Nature Communications</i> , 2017, 8, 142.	12.8	36
2	Frequent inactivation of SLIT2 and ROBO1 signaling in head and neck lesions: clinical and prognostic implications. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology</i> , 2015, 119, 202-212.	0.4	12
3	Prostate adenocarcinomas aberrantly expressing p63 are molecularly distinct from usual-type prostatic adenocarcinomas. <i>Modern Pathology</i> , 2015, 28, 446-456.	5.5	49
4	mTOR Signaling Feedback Modulates Mammary Epithelial Differentiation and Restrains Invasion Downstream of <i>PTEN</i> Loss. <i>Cancer Research</i> , 2013, 73, 5218-5231.	0.9	13
5	Overexpression of EGFR in Head and Neck Squamous Cell Carcinoma Is Associated with Inactivation of SH3GL2 and CDC25A Genes. <i>PLoS ONE</i> , 2013, 8, e63440.	2.5	68
6	Frequent alterations of <i>LOH11CR2A</i> , <i>PIG8</i> and <i>CHEK1</i> genes at chromosomal 11q24.1-24.2 region in breast carcinoma: Clinical and prognostic implications. <i>Molecular Oncology</i> , 2011, 5, 454-464.	4.6	23
7	PI3K/mTOR signaling regulates prostatic branching morphogenesis. <i>Developmental Biology</i> , 2011, 360, 329-342.	2.0	31
8	Frequent alterations of the candidate genes <i>hMLH1</i> , <i>ITGA9</i> and <i>RBSP3</i> in early dysplastic lesions of head and neck: Clinical and prognostic significance. <i>Cancer Science</i> , 2010, 101, 1511-1520.	3.9	50
9	LIMD1 is more frequently altered than RB1 in head and neck squamous cell carcinoma: clinical and prognostic implications. <i>Molecular Cancer</i> , 2010, 9, 58.	19.2	22
10	Alterations of ROBO1/DUTT1 and ROBO2 loci in early dysplastic lesions of head and neck: clinical and prognostic implications. <i>Human Genetics</i> , 2009, 125, 189-198.	3.8	28
11	<i>SH3GL2</i> and <i>CDKN2A/2B</i> loci are independently altered in early dysplastic lesions of head and neck: correlation with HPV infection and tobacco habit. <i>Journal of Pathology</i> , 2009, 217, 408-419.	4.5	48
12	Alterations of 3p21.31 tumor suppressor genes in head and neck squamous cell carcinoma: Correlation with progression and prognosis. <i>International Journal of Cancer</i> , 2008, 123, 2594-2604.	5.1	57