

Francisco Blanco-Vaca

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1139612/publications.pdf>

Version: 2024-02-01

158
papers

5,630
citations

81743

39
h-index

102304

66
g-index

166
all docs

166
docs citations

166
times ranked

6995
citing authors

#	ARTICLE	IF	CITATIONS
1	The A β ¹⁻⁴² /A β ¹⁻⁴⁰ ratio in CSF is more strongly associated to tau markers and clinical progression than A β ¹⁻⁴² alone. <i>Alzheimer's Research and Therapy</i> , 2022, 14, 20.	3.0	18
2	TMAO and Gut Microbial-Derived Metabolites TML and β BB Are Not Associated with Thrombotic Risk in Patients with Venous Thromboembolism. <i>Journal of Clinical Medicine</i> , 2022, 11, 1425.	1.0	2
3	Importance of cerebrospinal fluid storage conditions for the Alzheimer's disease diagnostics on an automated platform. <i>Clinical Chemistry and Laboratory Medicine</i> , 2022, 60, 1058-1063.	1.4	4
4	LDL, HDL and endocrine-related cancer: From pathogenic mechanisms to therapies. <i>Seminars in Cancer Biology</i> , 2021, 73, 134-157.	4.3	30
5	(r)HDL in theranostics: how do we apply HDL's biology for precision medicine in atherosclerosis management?. <i>Biomaterials Science</i> , 2021, 9, 3185-3208.	2.6	5
6	Comprehensive Genetic Testing of CYP21A2: A Retrospective Analysis in Patients with Suspected Congenital Adrenal Hyperplasia. <i>Journal of Clinical Medicine</i> , 2021, 10, 1183.	1.0	2
7	Reverse Cholesterol Transport Dysfunction Is a Feature of Familial Hypercholesterolemia. <i>Current Atherosclerosis Reports</i> , 2021, 23, 29.	2.0	8
8	Nicotinamide Protects Against Diet-Induced Body Weight Gain, Increases Energy Expenditure, and Induces White Adipose Tissue Beiging. <i>Molecular Nutrition and Food Research</i> , 2021, 65, e2100111.	1.5	9
9	The Capacity of APOB-Depleted Plasma in Inducing ATP-Binding Cassette A1/G1-Mediated Macrophage Cholesterol Efflux But Not Gut Microbial-Derived Metabolites Is Independently Associated with Mortality in Patients with ST-Segment Elevation Myocardial Infarction. <i>Biomedicines</i> , 2021, 9, 1336.	1.4	3
10	NAD ⁺ -Increasing Strategies to Improve Cardiometabolic Health?. <i>Frontiers in Endocrinology</i> , 2021, 12, 815565.	1.5	4
11	Therapeutic Potential of Emerging NAD ⁺ -Increasing Strategies for Cardiovascular Diseases. <i>Antioxidants</i> , 2021, 10, 1939.	2.2	11
12	Polygenic Markers in Patients Diagnosed of Autosomal Dominant Hypercholesterolemia in Catalonia: Distribution of Weighted LDL-c-Raising SNP Scores and Refinement of Variant Selection. <i>Biomedicines</i> , 2020, 8, 353.	1.4	6
13	Nicotinamide Prevents Apolipoprotein B-Containing Lipoprotein Oxidation, Inflammation and Atherosclerosis in Apolipoprotein E-Deficient Mice. <i>Antioxidants</i> , 2020, 9, 1162.	2.2	11
14	Phenol-Enriched Virgin Olive Oil Promotes Macrophage-Specific Reverse Cholesterol Transport In Vivo. <i>Biomedicines</i> , 2020, 8, 266.	1.4	9
15	Modulation of the Gut Microbiota by Olive Oil Phenolic Compounds: Implications for Lipid Metabolism, Immune System, and Obesity. <i>Nutrients</i> , 2020, 12, 2200.	1.7	48
16	Evaluation of biochemical and hematological parameters in adults with Down syndrome. <i>Scientific Reports</i> , 2020, 10, 13755.	1.6	4
17	LDL Receptor Regulates the Reverse Transport of Macrophage-Derived Unesterified Cholesterol via Concerted Action of the HDL-LDL Axis. <i>Circulation Research</i> , 2020, 127, 778-792.	2.0	45
18	Low-density lipoprotein receptor-related protein 1 deficiency in cardiomyocytes reduces susceptibility to insulin resistance and obesity. <i>Metabolism: Clinical and Experimental</i> , 2020, 106, 154191.	1.5	7

#	ARTICLE	IF	CITATIONS
19	Macrophage Cholesterol Efflux Downregulation Is Not Associated with Abdominal Aortic Aneurysm (AAA) Progression. <i>Biomolecules</i> , 2020, 10, 662.	1.8	2
20	HDL and LDL: Potential New Players in Breast Cancer Development. <i>Journal of Clinical Medicine</i> , 2019, 8, 853.	1.0	93
21	APOA1 oxidation is associated to dysfunctional high-density lipoproteins in human abdominal aortic aneurysm. <i>EBioMedicine</i> , 2019, 43, 43-53.	2.7	40
22	Human ApoA-I Overexpression Enhances Macrophage-Specific Reverse Cholesterol Transport but Fails to Prevent Inherited Diabetes in Mice. <i>International Journal of Molecular Sciences</i> , 2019, 20, 655.	1.8	6
23	Molecular analysis of APOB, SAR1B, ANGPTL3, and MTP in patients with primary hypocholesterolemia in a clinical laboratory setting: Evidence supporting polygenicity in mutation-negative patients. <i>Atherosclerosis</i> , 2019, 283, 52-60.	0.4	15
24	Phytosterols in Cancer: From Molecular Mechanisms to Preventive and Therapeutic Potentials. <i>Current Medicinal Chemistry</i> , 2019, 26, 6735-6749.	1.2	37
25	Vitamin B3 impairs reverse cholesterol transport in Apolipoprotein E-deficient mice. <i>Clínica E Investigaci3n En Arteriosclerosis (English Edition)</i> , 2019, 31, 251-260.	0.1	0
26	Lipid Profile Rather Than the LCAT Mutation Explains Renal Disease in Familial LCAT Deficiency. <i>Journal of Clinical Medicine</i> , 2019, 8, 1860.	1.0	10
27	Novel Insights into the Role of HDL-Associated Sphingosine-1-Phosphate in Cardiometabolic Diseases. <i>International Journal of Molecular Sciences</i> , 2019, 20, 6273.	1.8	18
28	Vitamin B3 impairs reverse cholesterol transport in Apolipoprotein E-deficient mice. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2019, 31, 251-260.	0.4	2
29	Altered HDL Remodeling and Functionality in Familial Hypercholesterolemia. <i>Journal of the American College of Cardiology</i> , 2018, 71, 466-468.	1.2	13
30	Autosomal dominant hypercholesterolemia in Catalonia: Correspondence between clinical-biochemical and genetic diagnostics in 967 patients studied in a multicenter clinical setting. <i>Journal of Clinical Lipidology</i> , 2018, 12, 1452-1462.	0.6	14
31	Administration of CORM-2 inhibits diabetic neuropathy but does not reduce dyslipidemia in diabetic mice. <i>PLoS ONE</i> , 2018, 13, e0204841.	1.1	12
32	A rare STAP1 mutation incompletely associated with familial hypercholesterolemia. <i>Clinica Chimica Acta</i> , 2018, 487, 270-274.	0.5	19
33	Impaired HDL (High-Density Lipoprotein)-Mediated Macrophage Cholesterol Efflux in Patients With Abdominal Aortic Aneurysm—Brief Report. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 2750-2754.	1.1	13
34	Trimethylamine N-Oxide: A Link among Diet, Gut Microbiota, Gene Regulation of Liver and Intestine Cholesterol Homeostasis and HDL Function. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3228.	1.8	138
35	LXR-dependent regulation of macrophage-specific reverse cholesterol transport is impaired in a model of genetic diabetes. <i>Translational Research</i> , 2017, 186, 19-35.e5.	2.2	5
36	Antiatherogenic potential of ezetimibe in sitosterolemia: Beyond plant sterols lowering. <i>Atherosclerosis</i> , 2017, 260, 94-96.	0.4	4

#	ARTICLE	IF	CITATIONS
37	Phytosterol-mediated inhibition of intestinal cholesterol absorption in mice is independent of liver X receptor. <i>Molecular Nutrition and Food Research</i> , 2017, 61, 1700055.	1.5	13
38	A novel homozygous mutation causing lecithinâ€“cholesterol acyltransferase deficiency in a proband of Romanian origin with a record of extreme gestational hyperlipidemia. <i>Journal of Clinical Lipidology</i> , 2017, 11, 1475-1479.e3.	0.6	6
39	Effect of PPAR- δ/γ agonist GW0742 treatment in the acute phase response and bloodâ€“brain barrier permeability following brain injury. <i>Translational Research</i> , 2017, 182, 27-48.	2.2	17
40	Human hepatic lipase overexpression in mice induces hepatic steatosis and obesity through promoting hepatic lipogenesis and white adipose tissue lipolysis and fatty acid uptake. <i>PLoS ONE</i> , 2017, 12, e0189834.	1.1	21
41	Lipoprotein hydrophobic core lipids are partially extruded to surface in smaller HDL: â€œHerniatedâ€“HDL, a common feature in diabetes. <i>Scientific Reports</i> , 2016, 6, 19249.	1.6	25
42	Modulation of autoimmune arthritis severity in mice by apolipoprotein E (ApoE) and cholesterol. <i>Clinical and Experimental Immunology</i> , 2016, 186, 292-303.	1.1	5
43	Clinically used selective estrogen receptor modulators affect different steps of macrophage-specific reverse cholesterol transport. <i>Scientific Reports</i> , 2016, 6, 32105.	1.6	14
44	ApoA-I mimetic administration, but not increased apoA-I-containing HDL, inhibits tumour growth in a mouse model of inherited breast cancer. <i>Scientific Reports</i> , 2016, 6, 36387.	1.6	34
45	Homozygous Familial Hypercholesterolemia in Spain. <i>Circulation: Cardiovascular Genetics</i> , 2016, 9, 504-510.	5.1	61
46	Chylomicrons: Advances in biology, pathology, laboratory testing, and therapeutics. <i>Clinica Chimica Acta</i> , 2016, 455, 134-148.	0.5	59
47	Chronic intermittent psychological stress promotes macrophage reverse cholesterol transport by impairing bile acid absorption in mice. <i>Physiological Reports</i> , 2015, 3, e12402.	0.7	21
48	Consumption of polyunsaturated fat improves the saturated fatty acidâ€“mediated impairment of HDL antioxidant potential. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 1987-1996.	1.5	16
49	PPAR- δ/γ activation promotes phospholipid transfer protein expression. <i>Biochemical Pharmacology</i> , 2015, 94, 101-108.	2.0	23
50	Enhanced vascular permeability facilitates entry of plasma HDL and promotes macrophage-reverse cholesterol transport from skin in mice. <i>Journal of Lipid Research</i> , 2015, 56, 241-253.	2.0	14
51	HDL and Lifestyle Interventions. <i>Handbook of Experimental Pharmacology</i> , 2015, 224, 569-592.	0.9	19
52	Quantification of In Vitro Macrophage Cholesterol Efflux and In Vivo Macrophage-Specific Reverse Cholesterol Transport. <i>Methods in Molecular Biology</i> , 2015, 1339, 211-233.	0.4	29
53	Remarkable quantitative and qualitative differences in HDL after niacin or fenofibrate therapy in type 2 diabetic patients. <i>Atherosclerosis</i> , 2015, 238, 213-219.	0.4	23
54	Genetic analysis does not confirm non-classical congenital adrenal hyperplasia in more than a third of the women followed with this diagnosis. <i>Hormones</i> , 2014, 13, 585-7.	0.9	1

#	ARTICLE	IF	CITATIONS
55	High-density lipoprotein cholesterol targeting for novel drug discovery: where have we gone wrong?. <i>Expert Opinion on Drug Discovery</i> , 2014, 9, 119-124.	2.5	8
56	Sitosterolemia: Diagnosis, Investigation, and Management. <i>Current Atherosclerosis Reports</i> , 2014, 16, 424.	2.0	92
57	Molecular analysis of chylomicronemia in a clinical laboratory setting: Diagnosis of 13 cases of lipoprotein lipase deficiency. <i>Clinica Chimica Acta</i> , 2014, 429, 61-68.	0.5	34
58	Bariatric surgery in morbidly obese patients improves the atherogenic qualitative properties of the plasma lipoproteins. <i>Atherosclerosis</i> , 2014, 234, 200-205.	0.4	29
59	The role of the gut in reverse cholesterol transport – Focus on the enterocyte. <i>Progress in Lipid Research</i> , 2013, 52, 317-328.	5.3	33
60	Resveratrol administration or SIRT1 overexpression does not increase LXR signaling and macrophage-to-feces reverse cholesterol transport in vivo. <i>Translational Research</i> , 2013, 161, 110-117.	2.2	8
61	Impact of the LDL subfraction phenotype on Lp-PLA2 distribution, LDL modification and HDL composition in type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2013, 12, 112.	2.7	47
62	Hepatic lipase- and endothelial lipase-deficiency in mice promotes macrophage-to-feces RCT and HDL antioxidant properties. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2013, 1831, 691-697.	1.2	24
63	Latent autoimmune diabetes in adults is perched between type 1 and type 2: evidence from adults in one region of Spain. <i>Diabetes/Metabolism Research and Reviews</i> , 2013, 29, 446-451.	1.7	49
64	Phytosterols inhibit the tumor growth and lipoprotein oxidizability induced by a high-fat diet in mice with inherited breast cancer. <i>Journal of Nutritional Biochemistry</i> , 2013, 24, 39-48.	1.9	41
65	Structural and functional analysis of APOA5 mutations identified in patients with severe hypertriglyceridemia. <i>Journal of Lipid Research</i> , 2013, 54, 649-661.	2.0	34
66	Methionine-induced hyperhomocysteinemia impairs the antioxidant ability of high-density lipoproteins without reducing in vivo macrophage-specific reverse cholesterol transport. <i>Molecular Nutrition and Food Research</i> , 2013, 57, 1814-1824.	1.5	18
67	Human scavenger protein AIM increases foam cell formation and CD36-mediated oxLDL uptake. <i>Journal of Leukocyte Biology</i> , 2013, 95, 509-520.	1.5	36
68	Acute Psychological Stress Accelerates Reverse Cholesterol Transport via Corticosterone-Dependent Inhibition of Intestinal Cholesterol Absorption. <i>Circulation Research</i> , 2012, 111, 1459-1469.	2.0	28
69	Identification of a novel mutation in the ANGPTL3 gene in two families diagnosed of familial hypobetalipoproteinemia without APOB mutation. <i>Clinica Chimica Acta</i> , 2012, 413, 552-555.	0.5	63
70	Effect of Improving Glycemic Control in Patients With Type 2 Diabetes Mellitus on Low-Density Lipoprotein Size, Electronegative Low-Density Lipoprotein and Lipoprotein-Associated Phospholipase A2 Distribution. <i>American Journal of Cardiology</i> , 2012, 110, 67-71.	0.7	37
71	Effect of atorvastatin on lipoprotein (a) and interleukin-10: A randomized placebo-controlled trial. <i>Diabetes and Metabolism</i> , 2011, 37, 124-130.	1.4	25
72	Differential effects of gemfibrozil and fenofibrate on reverse cholesterol transport from macrophages to feces in vivo. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2011, 1811, 104-110.	1.2	25

#	ARTICLE	IF	CITATIONS
73	ATP-binding cassette G5/G8 deficiency causes hypertriglyceridemia by affecting multiple metabolic pathways. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2011, 1811, 1186-1193.	1.2	20
74	The Cholesterol Content of Western Diets Plays a Major Role in the Paradoxical Increase in High-Density Lipoprotein Cholesterol and Upregulates the Macrophage Reverse Cholesterol Transport Pathway. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 2493-2499.	1.1	64
75	Mast Cell Activation In Vivo Impairs the Macrophage Reverse Cholesterol Transport Pathway in the Mouse. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 520-527.	1.1	20
76	Seeking Novel Targets for Improving In Vivo Macrophage-Specific Reverse Cholesterol Transport: Translating Basic Science into New Therapies for the Prevention and Treatment of Atherosclerosis. <i>Current Vascular Pharmacology</i> , 2011, 9, 220-237.	0.8	13
77	Disodium ascorbyl phytostanol phosphate (FM-VP4), a modified phytostanol, is a highly active hypocholesterolaemic agent that affects the enterohepatic circulation of both cholesterol and bile acids in mice. <i>British Journal of Nutrition</i> , 2010, 103, 153-160.	1.2	15
78	Human Apolipoprotein A-II Determines Plasma Triglycerides by Regulating Lipoprotein Lipase Activity and High-Density Lipoprotein Proteome. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2010, 30, 232-238.	1.1	69
79	Cyclooxygenase 2 Inhibition Exacerbates Palmitate-Induced Inflammation and Insulin Resistance in Skeletal Muscle Cells. <i>Endocrinology</i> , 2010, 151, 537-548.	1.4	52
80	La apolipoproteína A-II altera la composición apolipoproteica de HDL y su capacidad para activar la lipoproteína lipasa. <i>Clínica E Investigación En Arteriosclerosis</i> , 2010, 22, 192-197.	0.4	1
81	Increased plasma levels of plant sterols and atherosclerosis: A controversial issue. <i>Current Atherosclerosis Reports</i> , 2009, 11, 391-398.	2.0	18
82	Apolipoprotein Modulation of Streptococcal Serum Opacity Factor Activity against Human Plasma High-Density Lipoproteins. <i>Biochemistry</i> , 2009, 48, 8070-8076.	1.2	18
83	Serum soluble transferrin receptor concentrations are increased in central obesity. Results from a screening programme for hereditary hemochromatosis in men with hyperferritinemia. <i>Clinica Chimica Acta</i> , 2009, 400, 111-116.	0.5	30
84	New insights into the molecular actions of plant sterols and stanols in cholesterol metabolism. <i>Atherosclerosis</i> , 2009, 203, 18-31.	0.4	241
85	In vivo macrophage-specific RCT and antioxidant and antiinflammatory HDL activity measurements: New tools for predicting HDL atheroprotection. <i>Atherosclerosis</i> , 2009, 206, 321-327.	0.4	56
86	Efecto de la expresión de la PTEC, el gemfibrozilo y la rosiglitazona en el transporte inverso de colesterol desde macrófagos a heces in vivo. <i>Clínica E Investigación En Arteriosclerosis</i> , 2009, 21, 232-239.	0.4	0
87	Identification of ZNF366 and PTPRD as novel determinants of plasma homocysteine in a family-based genome-wide association study. <i>Blood</i> , 2009, 114, 1417-1422.	0.6	30
88	Standardization of a method to evaluate the antioxidant capacity of high-density lipoproteins. <i>International Journal of Biomedical Science</i> , 2009, 5, 402-10.	0.5	8
89	The Effects of Liposuction Removal of Subcutaneous Abdominal Fat on Lipid Metabolism are Independent of Insulin Sensitivity in Normal-Overweight Individuals. <i>Obesity Surgery</i> , 2008, 18, 408-414.	1.1	56
90	Genome-wide linkage analysis for identifying quantitative trait loci involved in the regulation of lipoprotein a (Lpa) levels. <i>European Journal of Human Genetics</i> , 2008, 16, 1372-1379.	1.4	24

#	ARTICLE	IF	CITATIONS
91	Role of vitamin D in the pathogenesis of type 2 diabetes mellitus. <i>Diabetes, Obesity and Metabolism</i> , 2008, 10, 185-197.	2.2	410
92	Phytosterols do not change susceptibility to obesity, insulin resistance, and diabetes induced by a high-fat diet in mice. <i>Metabolism: Clinical and Experimental</i> , 2008, 57, 1497-1501.	1.5	14
93	CETP activity variation in mice does not affect two major HDL antiatherogenic properties: Macrophage-specific reverse cholesterol transport and LDL antioxidant protection. <i>Atherosclerosis</i> , 2008, 196, 505-513.	0.4	17
94	Liver X receptor-mediated activation of reverse cholesterol transport from macrophages to feces in vivo requires ABCG5/G8. <i>Journal of Lipid Research</i> , 2008, 49, 1904-1911.	2.0	74
95	Homocysteine and Cognitive Impairment. <i>Dementia and Geriatric Cognitive Disorders</i> , 2008, 26, 506-512.	0.7	41
96	Unraveling the functions of macrophage transporters by measuring macrophage-specific reverse cholesterol transport in vivo. <i>Future Lipidology</i> , 2007, 2, 609-613.	0.5	0
97	Genetically based hypertension generated through interaction of mild hypoalphalipoproteinemia and mild hyperhomocysteinemia. <i>Journal of Hypertension</i> , 2007, 25, 1597-1607.	0.3	11
98	Folic acid supplementation delays atherosclerotic lesion development in apoE-deficient mice. <i>Life Sciences</i> , 2007, 80, 638-643.	2.0	26
99	Dietary phytosterols modulate T-helper immune response but do not induce apparent anti-inflammatory effects in a mouse model of acute, aseptic inflammation. <i>Life Sciences</i> , 2007, 80, 1951-1956.	2.0	42
100	Are LXR-regulated genes a major molecular target of plant sterols/stanols?. <i>Atherosclerosis</i> , 2007, 195, 210-211.	0.4	47
101	Differential intestinal mucosal protein expression in hypercholesterolemic mice fed a phytosterol-enriched diet. <i>Proteomics</i> , 2007, 7, 2659-2666.	1.3	9
102	Deficiency in monocyte chemoattractant protein-1 modifies lipid and glucose metabolism. <i>Experimental and Molecular Pathology</i> , 2007, 83, 361-366.	0.9	26
103	Liver Triglyceride Content in HIV-1-Infected Patients on Combination Antiretroviral Therapy Studied with ¹ H-MR Spectroscopy. <i>Antiviral Therapy</i> , 2007, 12, 195-204.	0.6	19
104	Manipulation of inflammation modulates hyperlipidemia in apolipoprotein E-deficient mice: A possible role for interleukin-6. <i>Cytokine</i> , 2006, 34, 224-232.	1.4	16
105	Antiatherogenic role of high-density lipoproteins: insights from genetically engineered-mice. <i>Frontiers in Bioscience - Landmark</i> , 2006, 11, 1328.	3.0	18
106	Phytosterol-mediated inhibition of intestinal cholesterol absorption is independent of ATP-binding cassette transporter A1. <i>British Journal of Nutrition</i> , 2006, 95, 618-622.	1.2	23
107	Apolipoprotein A5 S19W May Play a Role in Dysbetalipoproteinemia in Patients with the Apo E2/E2 Genotype. <i>Clinical Chemistry</i> , 2006, 52, 1974-1975.	1.5	14
108	Atorvastatin does not decrease or delay diabetes onset in two different mouse models of type 1 diabetes. <i>Diabetologia</i> , 2005, 48, 1671-1673.	2.9	11

#	ARTICLE	IF	CITATIONS
109	Overexpression of Human Apolipoprotein A-II in Transgenic Mice Does Not Impair Macrophage-Specific Reverse Cholesterol Transport In Vivo. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005, 25, e128-32.	1.1	61
110	Changes in intestinal and liver global gene expression in response to a phytosterol-enriched diet. <i>Atherosclerosis</i> , 2005, 181, 75-85.	0.4	84
111	A Genomewide Exploration Suggests a New Candidate Gene at Chromosome 11q23 as the Major Determinant of Plasma Homocysteine Levels: Results from the GAIT Project. <i>American Journal of Human Genetics</i> , 2005, 76, 925-933.	2.6	90
112	Turpentine-induced inflammation reduces the hepatic expression of the multiple drug resistance gene, the plasma cholesterol concentration and the development of atherosclerosis in apolipoprotein E deficient mice. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1733, 192-198.	1.2	22
113	Paradoxical exacerbation of combined hyperlipidemia in human apolipoprotein A-II transgenic mice treated with fenofibrate. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1737, 130-137.	1.2	20
114	Direct evidence in vivo of impaired macrophage-specific reverse cholesterol transport in ATP-binding cassette transporter A1-deficient mice. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1738, 6-9.	1.2	34
115	Patients with MEN-1 are more insulin-resistant than their non-affected relatives. <i>European Journal of Internal Medicine</i> , 2005, 16, 507-509.	1.0	7
116	Human Apolipoprotein A-II Enrichment Displaces Paraoxonase From HDL and Impairs Its Antioxidant Properties. <i>Circulation Research</i> , 2004, 95, 789-797.	2.0	118
117	Evaluation of Two Nonisotopic Immunoassays for Determination of Glutamic Acid Decarboxylase and Tyrosine Phosphatase Autoantibodies in Serum. <i>Clinical Chemistry</i> , 2004, 50, 1378-1382.	1.5	14
118	Moderate beer consumption does not change early or mature atherosclerosis in mice. <i>Nutrition Journal</i> , 2004, 3, 1.	1.5	123
119	Emerging cardiovascular risk factors in subclinical hypothyroidism: Lack of change after restoration of euthyroidism. <i>Metabolism: Clinical and Experimental</i> , 2004, 53, 1512-1515.	1.5	47
120	Phenytoin treatment reduces atherosclerosis in mice through mechanisms independent of plasma HDL-cholesterol concentration. <i>Atherosclerosis</i> , 2004, 174, 275-285.	0.4	9
121	Apolipoprotein A-II, genetic variation on chromosome 1q21-q24, and disease susceptibility. <i>Current Opinion in Lipidology</i> , 2004, 15, 247-253.	1.2	45
122	A Quantitative Trait Locus for Cholesterol/Low Density Lipoprotein within the Promoter of the Factor IX Gene. <i>Blood</i> , 2004, 104, 4000-4000.	0.6	0
123	Patient presenting multiple consecutive venous and arterial thrombotic events despite intensive conventional treatment: response after normalization of plasma homocysteine and N-acetylcysteine therapy. <i>Journal of Internal Medicine</i> , 2003, 254, 397-400.	2.7	3
124	Platelet-Activating Factor Acetylhydrolase Is Mainly Associated With Electronegative Low-Density Lipoprotein Subfraction. <i>Circulation</i> , 2003, 108, 92-96.	1.6	101
125	Mechanisms of HDL deficiency in mice overexpressing human apoA-II. <i>Journal of Lipid Research</i> , 2002, 43, 1734-1742.	2.0	25
126	Changes in low-density lipoprotein electronegativity and oxidizability after aerobic exercise are related to the increase in associated non-esterified fatty acids. <i>Atherosclerosis</i> , 2002, 160, 223-232.	0.4	77

#	ARTICLE	IF	CITATIONS
127	Thromboplastin-thrombomodulin-mediated Time and Serum Folate Levels Are Genetically Correlated with the Risk of Thromboembolic Disease: Results from the GAIT Project. <i>Thrombosis and Haemostasis</i> , 2002, 87, 68-73.	1.8	12
128	Overexpression of human apolipoprotein A-II in transgenic mice does not increase their susceptibility to insulin resistance and obesity. <i>Diabetologia</i> , 2002, 45, 600-601.	2.9	14
129	Density distribution of electronegative LDL in normolipemic and hyperlipemic subjects. <i>Journal of Lipid Research</i> , 2002, 43, 699-705.	2.0	81
130	ApoA-IMALLORCA impairs LCAT activation and induces dominant familial hypoalphalipoproteinemia. <i>Journal of Lipid Research</i> , 2002, 43, 115-123.	2.0	24
131	ApoA-I(MALLORCA) impairs LCAT activation and induces dominant familial hypoalphalipoproteinemia. <i>Journal of Lipid Research</i> , 2002, 43, 115-23.	2.0	18
132	Density distribution of electronegative LDL in normolipemic and hyperlipemic subjects. <i>Journal of Lipid Research</i> , 2002, 43, 699-705.	2.0	66
133	Apo(B)-dependent dyslipidemic phenotypes in type 1 diabetic patients. <i>European Journal of Internal Medicine</i> , 2001, 12, 496-502.	1.0	1
134	Which Cholesterol Are We Measuring with the Roche Direct, Homogeneous LDL-C Plus Assay?. <i>Clinical Chemistry</i> , 2001, 47, 124-126.	1.5	25
135	Role of apoA-II in lipid metabolism and atherosclerosis: advances in the study of an enigmatic protein. <i>Journal of Lipid Research</i> , 2001, 42, 1727-1739.	2.0	118
136	ApoA-II expression in CETP transgenic mice increases VLDL production and impairs VLDL clearance. <i>Journal of Lipid Research</i> , 2001, 42, 241-248.	2.0	42
137	Homocyst(e)ine and the C677T mutation of methylenetetrahydrofolate reductase in survivors of premature myocardial infarction. <i>Clinical Biochemistry</i> , 2000, 33, 509-512.	0.8	6
138	Determinants of plasma homocyst(e)ine in patients with nephrotic syndrome. <i>Journal of Molecular Medicine</i> , 2000, 78, 147-154.	1.7	9
139	Inaccuracy of Calculated LDL-Cholesterol in Type 2 Diabetes: Consequences for Patient Risk Classification and Therapeutic Decisions. <i>Clinical Chemistry</i> , 2000, 46, 1830-1832.	1.5	23
140	Comparison of the Abbott IMx [®] and a High-Performance Liquid Chromatography Method for Measuring Total Plasma Homocysteine. <i>Clinical Chemistry and Laboratory Medicine</i> , 2000, 38, 327-9.	1.4	25
141	Genetic Susceptibility to Thrombosis and Its Relationship to Physiological Risk Factors: The GAIT Study. <i>American Journal of Human Genetics</i> , 2000, 67, 1452-1459.	2.6	306
142	Increased production of very-low-density lipoproteins in transgenic mice overexpressing human apolipoprotein A-II and fed with a high-fat diet. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2000, 1488, 233-244.	1.2	31
143	Expression of human apolipoprotein A-II in apolipoprotein E-deficient mice induces features of familial combined hyperlipidemia. <i>Journal of Lipid Research</i> , 2000, 41, 1328-1338.	2.0	59
144	Pitfalls of Direct HDL-Cholesterol Measurements in Mouse Models of Hyperlipidemia and Atherosclerosis. <i>Clinical Chemistry</i> , 1999, 45, 1567-1569.	1.5	19

#	ARTICLE	IF	CITATIONS
145	Effect of simvastatin treatment on the electronegative low-density lipoprotein present in patients with heterozygous familial hypercholesterolemia. <i>American Journal of Cardiology</i> , 1999, 84, 655-659.	0.7	76
146	Free cholesterol deposition in the cornea of human apolipoprotein A-II transgenic mice with functional lecithin: Cholesterol acyltransferase deficiency. <i>Metabolism: Clinical and Experimental</i> , 1999, 48, 415-421.	1.5	23
147	Molecular Pathology of Multiple Endocrine Neoplasia Type I. <i>Diagnostic Molecular Pathology</i> , 1999, 8, 195-204.	2.1	10
148	A novel germline mutation in exon 5 of the multiple endocrine neoplasia type 1 gene. <i>Journal of Molecular Medicine</i> , 1998, 76, 837-839.	1.7	11
149	Plasma homocysteine is related to albumin excretion rate in patients with diabetes mellitus: a new link between diabetic nephropathy and cardiovascular disease?. <i>Diabetologia</i> , 1998, 41, 684-693.	2.9	186
150	Molecular Diagnosis of Lecithin: Cholesterol Acyltransferase Deficiency in a Presymptomatic Proband. <i>Clinical Chemistry and Laboratory Medicine</i> , 1998, 36, 443-8.	1.4	5
151	Human apolipoprotein A-II is a pro-atherogenic molecule when it is expressed in transgenic mice at a level similar to that in humans: evidence of a potentially relevant species-specific interaction with diet. <i>Journal of Lipid Research</i> , 1998, 39, 457-462.	2.0	61
152	Molecular Basis of Fish-Eye Disease in a Patient From Spain. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1997, 17, 1382-1391.	1.1	18
153	Functional Lecithin:Cholesterol Acyltransferase Deficiency and High Density Lipoprotein Deficiency in Transgenic Mice Overexpressing Human Apolipoprotein A-II. <i>Journal of Biological Chemistry</i> , 1996, 271, 6720-6728.	1.6	68
154	Effects of site-directed mutagenesis on the serine residues of human lecithin: Cholesterol acyltransferase. <i>Lipids</i> , 1994, 29, 803-809.	0.7	8
155	Structure of Human Apolipoprotein D: Locations of the Intermolecular and Intramolecular Disulfide Links. <i>Biochemistry</i> , 1994, 33, 12451-12455.	1.2	43
156	Disulfide linked dimers of apolipoprotein D in urine. <i>Electrophoresis</i> , 1993, 14, 1086-1087.	1.3	9
157	Roles of cysteines in human lecithin:cholesterol acyltransferase. <i>Biochemistry</i> , 1993, 32, 3089-3094.	1.2	26
158	Effects of site-directed mutagenesis on the N-glycosylation sites of human lecithin:cholesterol acyltransferase. <i>Biochemistry</i> , 1993, 32, 8732-8736.	1.2	35