

Shanshan Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11432135/publications.pdf>

Version: 2024-02-01

18
papers

562
citations

687363

13
h-index

794594

19
g-index

19
all docs

19
docs citations

19
times ranked

966
citing authors

#	ARTICLE	IF	CITATIONS
1	TAZ is indispensable for c-MYC-induced hepatocarcinogenesis. <i>Journal of Hepatology</i> , 2022, 76, 123-134.	3.7	28
2	The Hippo pathway effector TAZ induces intrahepatic cholangiocarcinoma in mice and is ubiquitously activated in the human disease. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022, 41, .	8.6	10
3	Distinct and Overlapping Roles of Hippo Effectors YAP and TAZ During Human and Mouse Hepatocarcinogenesis. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021, 11, 1095-1117.	4.5	21
4	Autophagy is involved in the neuroprotective effect of nicotiflorin. <i>Journal of Ethnopharmacology</i> , 2021, 278, 114279.	4.1	8
5	Combined CDK4/6 and Pan-mTOR Inhibition Is Synergistic Against Intrahepatic Cholangiocarcinoma. <i>Clinical Cancer Research</i> , 2019, 25, 403-413.	7.0	56
6	Hippo Cascade Controls Lineage Commitment of Liver Tumors in Mice and Humans. <i>American Journal of Pathology</i> , 2018, 188, 995-1006.	3.8	29
7	Association between QTc interval prolongation and outcomes of diabetic foot ulcers: Data from a 4-year follow-up study in China. <i>Diabetes Research and Clinical Practice</i> , 2018, 138, 26-34.	2.8	7
8	Efficacy of MEK inhibition in a K-Ras-driven cholangiocarcinoma preclinical model. <i>Cell Death and Disease</i> , 2018, 9, 31.	6.3	23
9	Notch2 controls hepatocyte-derived cholangiocarcinoma formation in mice. <i>Oncogene</i> , 2018, 37, 3229-3242.	5.9	79
10	Molecular profiling of intrahepatic cholangiocarcinoma: the search for new therapeutic targets. <i>Expert Review of Gastroenterology and Hepatology</i> , 2017, 11, 349-356.	3.0	16
11	Association between estimated glomerular filtration rate and outcomes in patients with diabetic foot ulcers: a 3-year follow-up study. <i>European Journal of Endocrinology</i> , 2017, 177, 41-50.	3.7	25
12	Synthesis and evaluation of asymmetric curcuminoid analogs as potential anticancer agents that downregulate NF- κ B activation and enhance the sensitivity of gastric cancer cell lines to irinotecan chemotherapy. <i>European Journal of Medicinal Chemistry</i> , 2017, 139, 917-925.	5.5	31
13	Pan-mTOR inhibitor MLN0128 is effective against intrahepatic cholangiocarcinoma in mice. <i>Journal of Hepatology</i> , 2017, 67, 1194-1203.	3.7	77
14	Activated mutant forms of $\text{PIK}3\text{CA}$ cooperate with RasV12 or c-Met to induce liver tumour formation in mice via AKT2/mTORC1 cascade. <i>Liver International</i> , 2016, 36, 1176-1186.	3.9	26
15	A novel double carbonyl analog of curcumin induces the apoptosis of human lung cancer H460 cells via the activation of the endoplasmic reticulum stress signaling pathway. <i>Oncology Reports</i> , 2016, 36, 1640-1648.	2.6	7
16	Contribution and Mobilization of Mesenchymal Stem Cells in a mouse model of carbon tetrachloride-induced liver fibrosis. <i>Scientific Reports</i> , 2015, 5, 17762.	3.3	38
17	Corticosterone Mediates the Inhibitory Effect of Restraint Stress on the Migration of Mesenchymal Stem Cell to Carbon Tetrachloride-Induced Fibrotic Liver by Downregulating CXCR4/7 Expression. <i>Stem Cells and Development</i> , 2015, 24, 587-596.	2.1	13
18	Maintenance Therapy With Continuous or Switch Strategy in Advanced Non-small Cell Lung Cancer. <i>Chest</i> , 2011, 140, 117-126.	0.8	67