

Richard E Gilbert

List of Publications by Citations

Source: <https://exaly.com/author-pdf/11455510/richard-e-gilbert-publications-by-citations.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

105
papers

7,636
citations

51
h-index

87
g-index

107
ext. papers

8,570
ext. citations

7
avg, IF

5.76
L-index

#	Paper	IF	Citations
105	The tubulointerstitium in progressive diabetic kidney disease: more than an aftermath of glomerular injury?. <i>Kidney International</i> , 1999 , 56, 1627-37	9.9	483
104	Hypertension Canada's 2018 Guidelines for Diagnosis, Risk Assessment, Prevention, and Treatment of Hypertension in Adults and Children. <i>Canadian Journal of Cardiology</i> , 2018 , 34, 506-525	3.8	348
103	Hypertension Canada's 2016 Canadian Hypertension Education Program Guidelines for Blood Pressure Measurement, Diagnosis, Assessment of Risk, Prevention, and Treatment of Hypertension. <i>Canadian Journal of Cardiology</i> , 2016 , 32, 569-88	3.8	314
102	The 2015 Canadian Hypertension Education Program recommendations for blood pressure measurement, diagnosis, assessment of risk, prevention, and treatment of hypertension. <i>Canadian Journal of Cardiology</i> , 2015 , 31, 549-68	3.8	222
101	Hypertension Canada's 2017 Guidelines for Diagnosis, Risk Assessment, Prevention, and Treatment of Hypertension in Adults. <i>Canadian Journal of Cardiology</i> , 2017 , 33, 557-576	3.8	205
100	Effect of Empagliflozin on Left Ventricular Mass in Patients With Type 2 Diabetes Mellitus and Coronary Artery Disease: The EMPA-HEART CardioLink-6 Randomized Clinical Trial. <i>Circulation</i> , 2019 , 140, 1693-1702	16.7	205
99	The (Pro)renin receptor: site-specific and functional linkage to the vacuolar H ⁺ -ATPase in the kidney. <i>Hypertension</i> , 2009 , 54, 261-9	8.5	205
98	Retinal neovascularization is prevented by blockade of the renin-angiotensin system. <i>Hypertension</i> , 2000 , 36, 1099-104	8.5	201
97	The 2014 Canadian Hypertension Education Program recommendations for blood pressure measurement, diagnosis, assessment of risk, prevention, and treatment of hypertension. <i>Canadian Journal of Cardiology</i> , 2014 , 30, 485-501	3.8	198
96	Heart failure in diabetes: effects of anti-hyperglycaemic drug therapy. <i>Lancet, The</i> , 2015 , 385, 2107-17	40	188
95	Direct actions of urotensin II on the heart: implications for cardiac fibrosis and hypertrophy. <i>Circulation Research</i> , 2003 , 93, 246-53	15.7	184
94	Are beta-blockers as efficacious in patients with diabetes mellitus as in patients without diabetes mellitus who have chronic heart failure? A meta-analysis of large-scale clinical trials. <i>American Heart Journal</i> , 2003 , 146, 848-53	4.9	152
93	Protein kinase C beta inhibition attenuates the progression of experimental diabetic nephropathy in the presence of continued hypertension. <i>Diabetes</i> , 2003 , 52, 512-8	0.9	151
92	Transforming growth factor beta 1 and renal injury following subtotal nephrectomy in the rat: role of the renin-angiotensin system. <i>Kidney International</i> , 1997 , 51, 1553-67	9.9	149
91	The 2013 Canadian Hypertension Education Program recommendations for blood pressure measurement, diagnosis, assessment of risk, prevention, and treatment of hypertension. <i>Canadian Journal of Cardiology</i> , 2013 , 29, 528-42	3.8	147
90	Effect of angiotensin II type 1 receptor blockade on experimental hepatic fibrogenesis. <i>Journal of Hepatology</i> , 2001 , 35, 376-85	13.4	139
89	Proximal Tubulopathy: Prime Mover and Key Therapeutic Target in Diabetic Kidney Disease. <i>Diabetes</i> , 2017 , 66, 791-800	0.9	133

88	The 2012 Canadian hypertension education program recommendations for the management of hypertension: blood pressure measurement, diagnosis, assessment of risk, and therapy. <i>Canadian Journal of Cardiology</i> , 2012 , 28, 270-87	3.8	133
87	The 2010 Canadian Hypertension Education Program recommendations for the management of hypertension: part 2 - therapy. <i>Canadian Journal of Cardiology</i> , 2010 , 26, 249-58	3.8	131
86	Pathological expression of renin and angiotensin II in the renal tubule after subtotal nephrectomy. Implications for the pathogenesis of tubulointerstitial fibrosis. <i>American Journal of Pathology</i> , 1999 , 155, 429-40	5.8	121
85	Demographics and concomitant disorders in heart failure. <i>Lancet, The</i> , 2003 , 362, 147-58	4.0	118
84	Role of VEGF in maintaining renal structure and function under normotensive and hypertensive conditions. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 14448-53	11.5	116
83	Effect of Empagliflozin on Erythropoietin Levels, Iron Stores, and Red Blood Cell Morphology in Patients With Type 2 Diabetes Mellitus and Coronary Artery Disease. <i>Circulation</i> , 2020 , 141, 704-707	16.7	115
82	Long-term administration of the histone deacetylase inhibitor vorinostat attenuates renal injury in experimental diabetes through an endothelial nitric oxide synthase-dependent mechanism. <i>American Journal of Pathology</i> , 2011 , 178, 2205-14	5.8	114
81	Inhibition of platelet-derived growth factor promotes pericyte loss and angiogenesis in ischemic retinopathy. <i>American Journal of Pathology</i> , 2004 , 164, 1263-73	5.8	99
80	Angiotensin type 2 receptor is expressed in the adult rat kidney and promotes cellular proliferation and apoptosis. <i>Kidney International</i> , 2000 , 58, 2437-51	9.9	98
79	The 2009 Canadian Hypertension Education Program recommendations for the management of hypertension: Part 2--therapy. <i>Canadian Journal of Cardiology</i> , 2009 , 25, 287-98	3.8	97
78	The renin-angiotensin system influences ocular endothelial cell proliferation in diabetes: transgenic and interventional studies. <i>American Journal of Pathology</i> , 2003 , 162, 151-60	5.8	97
77	The 2011 Canadian Hypertension Education Program recommendations for the management of hypertension: blood pressure measurement, diagnosis, assessment of risk, and therapy. <i>Canadian Journal of Cardiology</i> , 2011 , 27, 415-433.e1-2	3.8	93
76	Urinary connective tissue growth factor excretion in patients with type 1 diabetes and nephropathy. <i>Diabetes Care</i> , 2003 , 26, 2632-6	14.6	91
75	Expression, localization, and function of the thioredoxin system in diabetic nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2009 , 20, 730-41	12.7	89
74	Histone deacetylase inhibition attenuates diabetes-associated kidney growth: potential role for epigenetic modification of the epidermal growth factor receptor. <i>Kidney International</i> , 2011 , 79, 1312-21	9.9	89
73	Aminoguanidine ameliorates overexpression of pro-sclerotic growth factors and collagen deposition in experimental diabetic nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2001 , 12, 2098-2107	12.7	88
72	COX-2 inhibition and retinal angiogenesis in a mouse model of retinopathy of prematurity. <i>Investigative Ophthalmology and Visual Science</i> , 2003 , 44, 974-9		87
71	Tranilast attenuates cardiac matrix deposition in experimental diabetes: role of transforming growth factor-beta. <i>Cardiovascular Research</i> , 2005 , 65, 694-701	9.9	84

70	Sodium-glucose linked transporter-2 inhibitors: potential for renoprotection beyond blood glucose lowering?. <i>Kidney International</i> , 2014 , 86, 693-700	9.9	81
69	The CXCR4/CXCR7/SDF-1 pathway contributes to the pathogenesis of Shiga toxin-associated hemolytic uremic syndrome in humans and mice. <i>Journal of Clinical Investigation</i> , 2012 , 122, 759-76	15.9	79
68	Blockade of the renin-angiotensin and endothelin systems on progressive renal injury. <i>Hypertension</i> , 2000 , 36, 561-8	8.5	77
67	Effects of endothelin or angiotensin II receptor blockade on diabetes in the transgenic (mRen-2)27 rat. <i>Kidney International</i> , 2000 , 57, 1882-94	9.9	74
66	Long-term glycemic control and the rate of progression of early diabetic kidney disease. <i>Kidney International</i> , 1993 , 44, 855-9	9.9	74
65	Protein kinase Cbeta inhibition attenuates osteopontin expression, macrophage recruitment, and tubulointerstitial injury in advanced experimental diabetic nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2005 , 16, 1654-60	12.7	69
64	Attenuation of tubular apoptosis by blockade of the renin-angiotensin system in diabetic Ren-2 rats. <i>Kidney International</i> , 2002 , 61, 31-9	9.9	68
63	High glucose-induced thioredoxin-interacting protein in renal proximal tubule cells is independent of transforming growth factor-beta1. <i>American Journal of Pathology</i> , 2007 , 171, 744-54	5.8	67
62	Role of hyperlipidemia in progressive renal disease: focus on diabetic nephropathy. <i>Kidney International</i> , 1999 , 71, S31-6	9.9	67
61	Renal expression of transforming growth factor-beta inducible gene-h3 (beta ig-h3) in normal and diabetic rats. <i>Kidney International</i> , 1998 , 54, 1052-62	9.9	66
60	Endothelin receptor antagonism ameliorates mast cell infiltration, vascular hypertrophy, and epidermal growth factor expression in experimental diabetes. <i>Circulation Research</i> , 2000 , 86, 158-65	15.7	65
59	Osteopontin expression in progressive renal injury in remnant kidney: role of angiotensin II. <i>Kidney International</i> , 2000 , 58, 1469-80	9.9	64
58	Vasopeptidase inhibition attenuates the progression of renal injury in subtotal nephrectomized rats. <i>Kidney International</i> , 2001 , 60, 715-21	9.9	60
57	Mast cell infiltration and chemokine expression in progressive renal disease. <i>Kidney International</i> , 2003 , 64, 906-13	9.9	56
56	Diabetes-induced vascular hypertrophy is accompanied by activation of Na(+)-H(+) exchange and prevented by Na(+)-H(+) exchange inhibition. <i>Circulation Research</i> , 2000 , 87, 1133-40	15.7	55
55	Culture-modified bone marrow cells attenuate cardiac and renal injury in a chronic kidney disease rat model via a novel antifibrotic mechanism. <i>PLoS ONE</i> , 2010 , 5, e9543	3.7	51
54	Tranilast attenuates diastolic dysfunction and structural injury in experimental diabetic cardiomyopathy. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007 , 293, H2860-9	5.2	50
53	Inhibition of the epidermal growth factor receptor preserves podocytes and attenuates albuminuria in experimental diabetic nephropathy. <i>Nephrology</i> , 2011 , 16, 573-81	2.2	49

52	Diabetic vascular complications. <i>Clinical and Experimental Pharmacology and Physiology</i> , 1997 , 24, 770-5	3	48
51	SB-267268, a nonpeptidic antagonist of alpha(v)beta3 and alpha(v)beta5 integrins, reduces angiogenesis and VEGF expression in a mouse model of retinopathy of prematurity. <i>Investigative Ophthalmology and Visual Science</i> , 2006 , 47, 1600-5		48
50	Transforming growth factor-beta in human diabetic nephropathy: effects of ACE inhibition. <i>Diabetes Care</i> , 2006 , 29, 2670-5	14.6	47
49	Heart failure and nephropathy: catastrophic and interrelated complications of diabetes. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2006 , 1, 193-208	6.9	44
48	Sirtuin 1 activation attenuates cardiac fibrosis in a rodent pressure overload model by modifying Smad2/3 transactivation. <i>Cardiovascular Research</i> , 2018 , 114, 1629-1641	9.9	42
47	Macrophage infiltration and cellular proliferation in the non-ischemic kidney and heart following prolonged unilateral renal ischemia. <i>Nephron Physiology</i> , 2007 , 106, p54-62		41
46	Sodium-Glucose Linked Cotransporter-2 Inhibition Does Not Attenuate Disease Progression in the Rat Remnant Kidney Model of Chronic Kidney Disease. <i>PLoS ONE</i> , 2016 , 11, e0144640	3.7	37
45	A purpose-synthesised anti-fibrotic agent attenuates experimental kidney diseases in the rat. <i>PLoS ONE</i> , 2012 , 7, e47160	3.7	35
44	Hypertension in people with type 2 diabetes: Update on pharmacologic management. <i>Canadian Family Physician</i> , 2011 , 57, 997-1002, e347-53	0.9	35
43	SDF-1/CXCR4 signaling preserves microvascular integrity and renal function in chronic kidney disease. <i>PLoS ONE</i> , 2014 , 9, e92227	3.7	34
42	The cardiac (pro)renin receptor is primarily expressed in myocyte transverse tubules and is increased in experimental diabetic cardiomyopathy. <i>Journal of Hypertension</i> , 2011 , 29, 1175-84	1.9	34
41	Pathophysiology of diabetic nephropathy. <i>Metabolism: Clinical and Experimental</i> , 1998 , 47, 3-6	12.7	34
40	Vascular endothelial growth factor expression and glomerular endothelial cell loss in the remnant kidney model. <i>Nephrology Dialysis Transplantation</i> , 2003 , 18, 1286-92	4.3	32
39	Sirtuin 1 Activation Reduces Transforming Growth Factor- β -Induced Fibrogenesis and Affords Organ Protection in a Model of Progressive, Experimental Kidney and Associated Cardiac Disease. <i>American Journal of Pathology</i> , 2017 , 187, 80-90	5.8	30
38	FT011, a new anti-fibrotic drug, attenuates fibrosis and chronic heart failure in experimental diabetic cardiomyopathy. <i>European Journal of Heart Failure</i> , 2012 , 14, 549-62	12.3	30
37	Early-outgrowth bone marrow cells attenuate renal injury and dysfunction via an antioxidant effect in a mouse model of type 2 diabetes. <i>Diabetes</i> , 2012 , 61, 2114-25	0.9	29
36	SPARC gene expression is reduced in early diabetes-related kidney growth. <i>Kidney International</i> , 1995 , 48, 1216-25	9.9	29
35	Fluorescent microangiography is a novel and widely applicable technique for delineating the renal microvasculature. <i>PLoS ONE</i> , 2011 , 6, e24695	3.7	24

34	Load-independent effects of empagliflozin contribute to improved cardiac function in experimental heart failure with reduced ejection fraction. <i>Cardiovascular Diabetology</i> , 2020 , 19, 13	8.7	23
33	The interaction between the renin-angiotensin system and vascular endothelial growth factor in the pathogenesis of retinal neovascularization in diabetes. <i>Journal of Vascular Research</i> , 2001 , 38, 527-35	1.9	23
32	Dual inhibition of sodium-glucose linked cotransporters 1 and 2 exacerbates cardiac dysfunction following experimental myocardial infarction. <i>Cardiovascular Diabetology</i> , 2018 , 17, 99	8.7	21
31	Fas-induced apoptosis is a feature of progressive diabetic nephropathy in transgenic (mRen-2)27 rats: attenuation with renin-angiotensin blockade. <i>Nephrology</i> , 2004 , 9, 7-13	2.2	20
30	Effect of Basal Insulin Glargine on First and Recurrent Episodes of Heart Failure Hospitalization: The ORIGIN Trial (Outcome Reduction With Initial Glargine Intervention). <i>Circulation</i> , 2018 , 137, 88-90	16.7	19
29	The endothelium in diabetic nephropathy. <i>Current Atherosclerosis Reports</i> , 2014 , 16, 410	6	19
28	Protein kinase C-beta inhibition attenuates the progression of nephropathy in non-diabetic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 1782-90	4.3	18
27	Empagliflozin Reduces Myocardial Extracellular Volume in Patients With Type 2 Diabetes and Coronary Artery Disease. <i>JACC: Cardiovascular Imaging</i> , 2021 , 14, 1164-1173	8.4	18
26	Treatment of Diabetes in People With Heart Failure. <i>Canadian Journal of Diabetes</i> , 2018 , 42 Suppl 1, S196-S200	17	
25	Urinary transforming growth factor-beta in patients with diabetic nephropathy: implications for the pathogenesis of tubulointerstitial pathology. <i>Nephrology Dialysis Transplantation</i> , 2001 , 16, 2442-3	4.3	15
24	Attenuation of diabetes-associated mesenteric vascular hypertrophy with perindopril: morphological and molecular biological studies. <i>Metabolism: Clinical and Experimental</i> , 1998 , 47, 24-7	12.7	14
23	The impact of empagliflozin on kidney injury molecule-1: a subanalysis of the Effects of Empagliflozin on Cardiac Structure, Function, and Circulating Biomarkers in Patients with Type 2 Diabetes CardioLink-6 trial. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35, 895-897	4.3	12
22	Role of the eNOS-NO system in regulating the antiproteinuric effects of VEGF receptor 2 inhibition in diabetes. <i>BioMed Research International</i> , 2013 , 2013, 201475	3	11
21	Effects of Empagliflozin on Left Ventricular Remodeling in Patients with Type 2 Diabetes and Coronary Artery Disease: Echocardiographic Substudy of the EMPA-HEART CardioLink-6 Randomized Clinical Trial. <i>Journal of the American Society of Echocardiography</i> , 2020 , 33, 644-646	5.8	10
20	Impaired cardiac anti-oxidant activity in diabetes: human and correlative experimental studies. <i>Acta Diabetologica</i> , 2014 , 51, 771-82	3.9	10
19	A new anti-fibrotic drug attenuates cardiac remodeling and systolic dysfunction following experimental myocardial infarction. <i>International Journal of Cardiology</i> , 2013 , 168, 1174-85	3.2	10
18	Bone marrow cell therapies for endothelial repair and their relevance to kidney disease. <i>Seminars in Nephrology</i> , 2012 , 32, 215-23	4.8	10
17	Urotensin-II as a novel therapeutic target in the clinical management of cardiorenal disease. <i>Current Opinion in Investigational Drugs</i> , 2004 , 5, 276-82		9

16	Is there a role for endothelin antagonists in diabetic renal disease?. <i>Diabetes, Obesity and Metabolism</i> , 2000 , 2, 15-24	6.7	8
15	Reversing CXCL10 Deficiency Ameliorates Kidney Disease in Diabetic Mice. <i>American Journal of Pathology</i> , 2018 , 188, 2763-2773	5.8	8
14	Diabetes and hypertension: prognostic and therapeutic considerations. <i>Blood Pressure</i> , 1995 , 4, 329-38	1.7	7
13	Treatment of Hypertension. <i>Canadian Journal of Diabetes</i> , 2018 , 42 Suppl 1, S186-S189	2.1	6
12	Impaired SIRT1 activity leads to diminution in glomerular endowment without accelerating age-associated GFR decline. <i>Physiological Reports</i> , 2019 , 7, e14044	2.6	3
11	Heart failure: fatal, forgotten, and frequent in type 1 diabetes too. <i>Lancet Diabetes and Endocrinology</i> , 2015 , 3, 832-4	18.1	3
10	Cell therapy for diabetic nephropathy: is the future, now?. <i>Seminars in Nephrology</i> , 2012 , 32, 486-93	4.8	3
9	Hyperglycemia and renal mass ablation synergistically augment albuminuria in the diabetic subtotal nephrectomized rat: implications for modeling diabetic nephropathy. <i>Nephron Extra</i> , 2012 , 2, 115-24		3
8	Progenitor cell secretory products exert additive renoprotective effects when combined with ace inhibitors in experimental CKD. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2016 , 17,	3	2
7	Vasoactive Molecules and the Kidney 2012 , 384-420		2
6	Impact of sodium glucose linked cotransporter-2 inhibition on renal microvascular oxygen tension in a rodent model of diabetes mellitus. <i>Physiological Reports</i> , 2021 , 9, e14890	2.6	2
5	Application of Modular Therapy for Renoprotection in Experimental Chronic Kidney Disease. <i>Tissue Engineering - Part A</i> , 2015 , 21, 1963-72	3.9	1
4	Impact of empagliflozin on right ventricular parameters and function among patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2021 , 20, 200	8.7	0
3	Late intervention in the remnant kidney model attenuates proteinuria but not glomerular filtration rate decline. <i>Nephrology</i> , 2021 , 26, 270-279	2.2	0
2	Angiotensin-converting enzyme inhibition attenuates renal platelet-derived growth factor gene expression and cell proliferation in subtotal nephrectomy. <i>Nephrology</i> , 2001 , 6, 290-297	2.2	
1	Hypertension revisited. <i>Canadian Family Physician</i> , 2012 , 58, 634-6	0.9	