

# Philippe Gual

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1145987/philippe-gual-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85  
papers

8,536  
citations

39  
h-index

92  
g-index

99  
ext. papers

9,592  
ext. citations

5.8  
avg, IF

5.21  
L-index

#	Paper	IF	Citations
85	SYK-3BP2 Pathway Activity in Parenchymal and Myeloid Cells Is a Key Pathogenic Factor in Metabolic Steatohepatitis. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , <b>2021</b> , 13, 173-191	7.9	1
84	Plasmatic osteopontin and vascular access dysfunction in hemodialysis patients: a cross-sectional, case-control study (The OSMOSIS Study). <i>Journal of Nephrology</i> , <b>2021</b> , 1	4.8	1
83	Metabolic Fatty Liver Disease in Children: A Growing Public Health Problem.. <i>Biomedicines</i> , <b>2021</b> , 9,	4.8	3
82	Chronic Inflammation in Non-Alcoholic Steatohepatitis: Molecular Mechanisms and Therapeutic Strategies. <i>Frontiers in Endocrinology</i> , <b>2020</b> , 11, 597648	5.7	20
81	No association between binge eating disorder and severity of non-alcoholic fatty liver disease in severely obese patients. <i>JGH Open</i> , <b>2020</b> , 4, 525-531	1.8	4
80	REDD1 deficiency protects against nonalcoholic hepatic steatosis induced by high-fat diet. <i>FASEB Journal</i> , <b>2020</b> , 34, 5046-5060	0.9	10
79	Hepatic FNDC5 is a potential local protective factor against Non-Alcoholic Fatty Liver. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , <b>2020</b> , 1866, 165705	6.9	19
78	Endoplasmic reticulum stress mediates resistance to BCL-2 inhibitor in uveal melanoma cells. <i>Cell Death Discovery</i> , <b>2020</b> , 6, 22	6.9	4
77	MCD diet-induced steatohepatitis generates a diurnal rhythm of associated biomarkers and worsens liver injury in Klf10 deficient mice. <i>Scientific Reports</i> , <b>2020</b> , 10, 12139	4.9	6
76	Natural Killer Cells and Type 1 Innate Lymphoid Cells Are New Actors in Non-alcoholic Fatty Liver Disease. <i>Frontiers in Immunology</i> , <b>2019</b> , 10, 1192	8.4	27
75	The Differential Expression of Cide Family Members is Associated with Nafld Progression from Steatosis to Steatohepatitis. <i>Scientific Reports</i> , <b>2019</b> , 9, 7501	4.9	10
74	The mid-infrared spectroscopy: A novel non-invasive diagnostic tool for NASH diagnosis in severe obesity. <i>JHEP Reports</i> , <b>2019</b> , 1, 361-368	10.3	5
73	Bax inhibitor-1 protects from nonalcoholic steatohepatitis by limiting inositol-requiring enzyme 1 alpha signaling in mice. <i>Hepatology</i> , <b>2018</b> , 68, 515-532	11.2	45
72	Visceral Adipose Tissue Drives Cardiac Aging Through Modulation of Fibroblast Senescence by Osteopontin Production. <i>Circulation</i> , <b>2018</b> , 138, 809-822	16.7	63
71	Screening for therapeutic trials and treatment indication in clinical practice: MACK-3, a new blood test for the diagnosis of fibrotic NASH. <i>Alimentary Pharmacology and Therapeutics</i> , <b>2018</b> , 47, 1387-1396	6.1	31
70	Non-invasive Evaluation of NAFLD with Indocyanine Green Clearance Test: a Preliminary Study in Morbidly Obese Patients Undergoing Bariatric Surgery. <i>Obesity Surgery</i> , <b>2018</b> , 28, 735-742	3.7	9
69	Role of Lipid Droplet Proteins in the Development of NAFLD and Hepatic Insulin Resistance <b>2018</b> ,		3

68	Early postnatal soluble FGFR3 therapy prevents the atypical development of obesity in achondroplasia. <i>PLoS ONE</i> , <b>2018</b> , 13, e0195876	3.7	15
67	Metabolism dysregulation induces a specific lipid signature of nonalcoholic steatohepatitis in patients. <i>Scientific Reports</i> , <b>2017</b> , 7, 46658	4.9	99
66	Autophagy in chronic liver diseases: the two faces of Janus. <i>American Journal of Physiology - Cell Physiology</i> , <b>2017</b> , 312, C263-C273	5.4	41
65	CD44 is a key player in non-alcoholic steatohepatitis. <i>Journal of Hepatology</i> , <b>2017</b> , 67, 328-338	13.4	60
64	Baseline Anthropometric and Metabolic Parameters Correlate with Weight Loss in Women 1-Year After Laparoscopic Roux-En-Y Gastric Bypass. <i>Obesity Surgery</i> , <b>2017</b> , 27, 2940-2949	3.7	9
63	Determinants associated with the correction of glomerular hyper-filtration one year after bariatric surgery. <i>Surgery for Obesity and Related Diseases</i> , <b>2017</b> , 13, 1760-1766	3	10
62	Immunoglobulin G4-associated autoimmune hepatitis may be found in Western countries. <i>Digestive and Liver Disease</i> , <b>2016</b> , 48, 302-8	3.3	13
61	Roux-En Y Gastric Bypass Results in Long-Term Remission of Hepatocyte Apoptosis and Hepatic Histological Features of Non-alcoholic Steatohepatitis. <i>Frontiers in Physiology</i> , <b>2016</b> , 7, 344	4.6	20
60	Compounds Triggering ER Stress Exert Anti-Melanoma Effects and Overcome BRAF Inhibitor Resistance. <i>Cancer Cell</i> , <b>2016</b> , 29, 805-819	24.3	110
59	Severe Vitamin D Deficiency Is Not Associated with Liver Damage in Morbidly Obese Patients. <i>Obesity Surgery</i> , <b>2016</b> , 26, 2138-2143	3.7	18
58	Hematopoietic IKBKE limits the chronicity of inflammasome priming and metaflammation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2015</b> , 112, 506-11	11.5	22
57	ER stress induces NLRP3 inflammasome activation and hepatocyte death. <i>Cell Death and Disease</i> , <b>2015</b> , 6, e1879	9.8	204
56	Hypoxia inhibits Cavin-1 and Cavin-2 expression and down-regulates caveolae in adipocytes. <i>Endocrinology</i> , <b>2015</b> , 156, 789-801	4.8	21
55	Defects in mitophagy promote redox-driven metabolic syndrome in the absence of TP53INP1. <i>EMBO Molecular Medicine</i> , <b>2015</b> , 7, 802-18	12	30
54	Severe Vitamin D Deficiency May be an Additional Cofactor for the Occurrence of Alcoholic Steatohepatitis. <i>Alcoholism: Clinical and Experimental Research</i> , <b>2015</b> , 39, 1027-33	3.7	20
53	The promyelocytic leukemia protein is upregulated in conditions of obesity and liver steatosis. <i>International Journal of Biological Sciences</i> , <b>2015</b> , 11, 629-32	11.2	9
52	Effects of sleeve gastrectomy in high fat diet-induced obese mice: respective role of reduced caloric intake, white adipose tissue inflammation and changes in adipose tissue and ectopic fat depots. <i>Surgical Endoscopy and Other Interventional Techniques</i> , <b>2014</b> , 28, 592-602	5.2	20
51	Body composition, anthropometrics, energy expenditure, systemic inflammation, in premenopausal women 1 year after laparoscopic Roux-en-Y gastric bypass. <i>Surgical Endoscopy and Other Interventional Techniques</i> , <b>2014</b> , 28, 500-7	5.2	31

50	Obesity et maladie du foie gras non alcoolique. <i>Obesite</i> , <b>2014</b> , 9, 271-279	0.1	
49	Low Levels of 25-Hydroxy Vitamin D are Independently Associated with the Risk of Bacterial Infection in Cirrhotic Patients. <i>Clinical and Translational Gastroenterology</i> , <b>2014</b> , 5, e56	4.2	33
48	Osteopontin deficiency aggravates hepatic injury induced by ischemia-reperfusion in mice. <i>Cell Death and Disease</i> , <b>2014</b> , 5, e1208	9.8	32
47	Autophagy and non-alcoholic fatty liver disease. <i>BioMed Research International</i> , <b>2014</b> , 2014, 120179	3	81
46	Prophylaxis of bacterial infections in cirrhosis: is an optimal 25-OH vitamin D level required?. <i>Journal of Hepatology</i> , <b>2014</b> , 61, 965-6	13.4	4
45	M2 Kupffer cells promote M1 Kupffer cell apoptosis: a protective mechanism against alcoholic and nonalcoholic fatty liver disease. <i>Hepatology</i> , <b>2014</b> , 59, 130-42	11.2	336
44	Non-alcoholic steatohepatitis in morbidly obese patients. <i>Clinics and Research in Hepatology and Gastroenterology</i> , <b>2013</b> , 37, 17-29	2.4	23
43	Postnatal soluble FGFR3 therapy rescues achondroplasia symptoms and restores bone growth in mice. <i>Science Translational Medicine</i> , <b>2013</b> , 5, 203ra124	17.5	61
42	Solute carrier family 2 member 1 is involved in the development of nonalcoholic fatty liver disease. <i>Hepatology</i> , <b>2013</b> , 57, 505-14	11.2	18
41	Autophagy, signaling and obesity. <i>Pharmacological Research</i> , <b>2012</b> , 66, 513-25	10.2	58
40	Reply to: C-reactive protein levels in non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , <b>2012</b> , 56, 508-510	13.4	1
39	Regular coffee but not espresso drinking is protective against fibrosis in a cohort mainly composed of morbidly obese European women with NAFLD undergoing bariatric surgery. <i>Journal of Hepatology</i> , <b>2012</b> , 57, 1090-6	13.4	106
38	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , <b>2012</b> , 8, 445-544	14.2	2783
37	The osteopontin level in liver, adipose tissue and serum is correlated with fibrosis in patients with alcoholic liver disease. <i>PLoS ONE</i> , <b>2012</b> , 7, e35612	3.7	61
36	Obesity-dependent metabolic signatures associated with nonalcoholic fatty liver disease progression. <i>Journal of Proteome Research</i> , <b>2012</b> , 11, 2521-32	5.6	137
35	Identification of adipose tissue dendritic cells correlated with obesity-associated insulin-resistance and inducing Th17 responses in mice and patients. <i>Diabetes</i> , <b>2012</b> , 61, 2238-47	0.9	250
34	C-reactive protein levels in relation to various features of non-alcoholic fatty liver disease among obese patients. <i>Journal of Hepatology</i> , <b>2011</b> , 55, 660-665	13.4	68
33	Serum markers of hepatocyte death and apoptosis are non invasive biomarkers of severe fibrosis in patients with alcoholic liver disease. <i>PLoS ONE</i> , <b>2011</b> , 6, e17599	3.7	49

32	Hemojuvelin: a new link between obesity and iron homeostasis. <i>Obesity</i> , <b>2011</b> , 19, 1545-51	8	29
31	A new composite model including metabolic syndrome, alanine aminotransferase and cytokeratin-18 for the diagnosis of non-alcoholic steatohepatitis in morbidly obese patients. <i>Alimentary Pharmacology and Therapeutics</i> , <b>2010</b> , 32, 1315-22	6.1	101
30	Tpl2 kinase is upregulated in adipose tissue in obesity and may mediate interleukin-1beta and tumor necrosis factor- $\alpha$ effects on extracellular signal-regulated kinase activation and lipolysis. <i>Diabetes</i> , <b>2010</b> , 59, 61-70	0.9	49
29	Liquid chromatography-mass spectrometry-based parallel metabolic profiling of human and mouse model serum reveals putative biomarkers associated with the progression of nonalcoholic fatty liver disease. <i>Journal of Proteome Research</i> , <b>2010</b> , 9, 4501-12	5.6	119
28	Hepatic expression patterns of inflammatory and immune response genes associated with obesity and NASH in morbidly obese patients. <i>PLoS ONE</i> , <b>2010</b> , 5, e13577	3.7	159
27	Rab4b is a small GTPase involved in the control of the glucose transporter GLUT4 localization in adipocyte. <i>PLoS ONE</i> , <b>2009</b> , 4, e5257	3.7	29
26	Cannabinoid CB2 receptor potentiates obesity-associated inflammation, insulin resistance and hepatic steatosis. <i>PLoS ONE</i> , <b>2009</b> , 4, e5844	3.7	158
25	Elevated expression of osteopontin may be related to adipose tissue macrophage accumulation and liver steatosis in morbid obesity. <i>Diabetes</i> , <b>2009</b> , 58, 125-33	0.9	108
24	Impact of laparoscopic Roux-en-Y gastric bypass on metabolic syndrome, inflammation, and insulin resistance in super versus morbidly obese women. <i>Obesity Surgery</i> , <b>2009</b> , 19, 577-82	3.7	40
23	Involvement of TNF-alpha in abnormal adipocyte and muscle sortilin expression in obese mice and humans. <i>Diabetologia</i> , <b>2009</b> , 52, 932-40	10.3	44
22	Bariatric surgery can correct iron depletion in morbidly obese women: a link with chronic inflammation. <i>Obesity Surgery</i> , <b>2008</b> , 18, 709-14	3.7	58
21	HMG2A is the partner of MDM2 in well-differentiated and dedifferentiated liposarcomas whereas CDK4 belongs to a distinct inconsistent amplicon. <i>International Journal of Cancer</i> , <b>2008</b> , 122, 2233-41	7.5	153
20	C3H/HeJ mice carrying a toll-like receptor 4 mutation are protected against the development of insulin resistance in white adipose tissue in response to a high-fat diet. <i>Diabetologia</i> , <b>2007</b> , 50, 1267-76	10.3	278
19	Hepatocyte growth factor induces glucose uptake in 3T3-L1 adipocytes through A Gab1/phosphatidylinositol 3-kinase/Glut4 pathway. <i>Journal of Biological Chemistry</i> , <b>2007</b> , 282, 10325-32	5.4	32
18	Glucose intolerance and hypoadiponectinemia are already present in lean patients with chronic hepatitis C infected with genotype non-3 viruses. <i>European Journal of Gastroenterology and Hepatology</i> , <b>2007</b> , 19, 671-7	2.2	15
17	Osmotic regulation of cellular glucose uptake. <i>Methods in Enzymology</i> , <b>2007</b> , 428, 343-54	1.7	2
16	Differential effects of IRS1 phosphorylated on Ser307 or Ser632 in the induction of insulin resistance by oxidative stress. <i>Diabetologia</i> , <b>2006</b> , 49, 2463-73	10.3	69
15	Enigma interacts with adaptor protein with PH and SH2 domains to control insulin-induced actin cytoskeleton remodeling and glucose transporter 4 translocation. <i>Molecular Endocrinology</i> , <b>2006</b> , 20, 2864-75		24

14	The inflammatory C-reactive protein is increased in both liver and adipose tissue in severely obese patients independently from metabolic syndrome, Type 2 diabetes, and NASH. <i>American Journal of Gastroenterology</i> , <b>2006</b> , 101, 1824-33	0.7	137
13	Increased adipose tissue expression of hepcidin in severe obesity is independent from diabetes and NASH. <i>Gastroenterology</i> , <b>2006</b> , 131, 788-96	13.3	365
12	Positive and negative regulation of insulin signaling through IRS-1 phosphorylation. <i>Biochimie</i> , <b>2005</b> , 87, 99-109	4.6	646
11	MAP kinases and mTOR mediate insulin-induced phosphorylation of insulin receptor substrate-1 on serine residues 307, 612 and 632. <i>Diabetologia</i> , <b>2003</b> , 46, 1532-42	10.3	119
10	Reduced activation of phosphatidylinositol-3 kinase and increased serine 636 phosphorylation of insulin receptor substrate-1 in primary culture of skeletal muscle cells from patients with type 2 diabetes. <i>Diabetes</i> , <b>2003</b> , 52, 1319-25	0.9	241
9	Hyperosmotic stress inhibits insulin receptor substrate-1 function by distinct mechanisms in 3T3-L1 adipocytes. <i>Journal of Biological Chemistry</i> , <b>2003</b> , 278, 26550-7	5.4	58
8	A Crk-II/TC10 signaling pathway is required for osmotic shock-stimulated glucose transport. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 43980-6	5.4	26
7	Gab1 phosphorylation: a novel mechanism for negative regulation of HGF receptor signaling. <i>Oncogene</i> , <b>2001</b> , 20, 156-66	9.2	39
6	Differential requirement of the last C-terminal tail of Met receptor for cell transformation and invasiveness. <i>Oncogene</i> , <b>2001</b> , 20, 5493-502	9.2	5
5	Sustained recruitment of phospholipase C-gamma to Gab1 is required for HGF-induced branching tubulogenesis. <i>Oncogene</i> , <b>2000</b> , 19, 1509-18	9.2	145
4	MET(PRC) mutations in the Ron receptor result in upregulation of tyrosine kinase activity and acquisition of oncogenic potential. <i>Journal of Cellular Physiology</i> , <b>1999</b> , 181, 507-14	7	22
3	Interaction of Janus kinases JAK-1 and JAK-2 with the insulin receptor and the insulin-like growth factor-1 receptor. <i>Endocrinology</i> , <b>1998</b> , 139, 884-93	4.8	98
2	Role of the insulin receptor C-terminal acidic domain in the modulation of the receptor kinase by polybasic effectors. <i>FEBS Journal</i> , <b>1996</b> , 241, 186-92		4
1	Insulin receptor-induced phosphorylation of cellular and synthetic substrates is regulated by the receptor beta-subunit C-terminus		2