

# Sabine Werner

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/1147954/sabine-werner-publications-by-citations.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118  
papers

16,040  
citations

50  
h-index

126  
g-index

139  
ext. papers

18,129  
ext. citations

9.6  
avg, IF

6.75  
L-index

| #   | Paper   | IF   | Citations |
|-----|---|------|-----------|
| 118 | Wound repair and regeneration. <i>Nature</i> , <b>2008</b> , 453, 314-21  | 50.4 | 3599      |
| 117 | Regulation of wound healing by growth factors and cytokines. <i>Physiological Reviews</i> , <b>2003</b> , 83, 835-70  | 47.9 | 2436      |
| 116 | Keratinocyte-fibroblast interactions in wound healing. <i>Journal of Investigative Dermatology</i> , <b>2007</b> , 127, 998-1008  | 4.3  | 777       |
| 115 | Active caspase-1 is a regulator of unconventional protein secretion. <i>Cell</i> , <b>2008</b> , 132, 818-31  | 56.2 | 682       |
| 114 | Cancer as an overhealing wound: an old hypothesis revisited. <i>Nature Reviews Molecular Cell Biology</i> , <b>2008</b> , 9, 628-38   | 48.7 | 653       |
| 113 | The function of KGF in morphogenesis of epithelium and reepithelialization of wounds. <i>Science</i> , <b>1994</b> , 266, 819-22  | 33.3 | 526       |
| 112 | Regulation of vascular endothelial growth factor expression in cultured keratinocytes. Implications for normal and impaired wound healing. <i>Journal of Biological Chemistry</i> , <b>1995</b> , 270, 12607-13               | 5.4  | 519       |
| 111 | Oxidative stress in normal and impaired wound repair. <i>Pharmacological Research</i> , <b>2008</b> , 58, 165-71  | 10.2 | 450       |
| 110 | The inflammasome mediates UVB-induced activation and secretion of interleukin-1beta by keratinocytes. <i>Current Biology</i> , <b>2007</b> , 17, 1140-5   | 6.3  | 417       |
| 109 | Differential regulation of pro-inflammatory cytokines during wound healing in normal and glucocorticoid-treated mice. <i>Cytokine</i> , <b>1996</b> , 8, 548-56   | 4    | 391       |
| 108 | Nrf2 transcription factor, a novel target of keratinocyte growth factor action which regulates gene expression and inflammation in the healing skin wound. <i>Molecular and Cellular Biology</i> , <b>2002</b> , 22, 5492-505 | 4.8  | 314       |
| 107 | Psoriasisiform dermatitis is driven by IL-36-mediated DC-keratinocyte crosstalk. <i>Journal of Clinical Investigation</i> , <b>2012</b> , 122, 3965-76  | 15.9 | 278       |
| 106 | c-Met is essential for wound healing in the skin. <i>Journal of Cell Biology</i> , <b>2007</b> , 177, 151-62  | 7.3  | 243       |
| 105 | Impaired liver regeneration in Nrf2 knockout mice: role of ROS-mediated insulin/IGF-1 resistance. <i>EMBO Journal</i> , <b>2008</b> , 27, 212-23  | 13   | 203       |
| 104 | Roles of activin in tissue repair, fibrosis, and inflammatory disease. <i>Cytokine and Growth Factor Reviews</i> , <b>2006</b> , 17, 157-71   | 17.9 | 173       |
| 103 | Regulation of liver regeneration by growth factors and cytokines. <i>EMBO Molecular Medicine</i> , <b>2010</b> , 2, 294-305   | 12   | 166       |
| 102 | Strong induction of activin expression after injury suggests an important role of activin in wound repair. <i>Developmental Biology</i> , <b>1996</b> , 173, 490-8  | 3.1  | 163       |

|     |   |      |     |
|-----|---|------|-----|
| 101 | The Nrf2 transcription factor protects from toxin-induced liver injury and fibrosis. <i>Laboratory Investigation</i> , <b>2008</b> , 88, 1068-78                                  | 5.9  | 161 |
| 100 | Activating mutations of the tyrosine kinase receptor FGFR3 are associated with benign skin tumors in mice and humans. <i>Human Molecular Genetics</i> , <b>2005</b> , 14, 1153-60 | 5.6  | 151 |
| 99  | Transcriptional control of wound repair. <i>Annual Review of Cell and Developmental Biology</i> , <b>2007</b> , 23, 69-92   | 2.6  | 133 |
| 98  | Nrf2 links epidermal barrier function with antioxidant defense. <i>EMBO Molecular Medicine</i> , <b>2012</b> , 4, 364-72  | 2.2  | 122 |
| 97  | Fibroblast growth factors: key players in regeneration and tissue repair. <i>Development (Cambridge)</i> , <b>2017</b> , 144, 4047-4060   | 6.6  | 119 |
| 96  | Different types of ROS-scavenging enzymes are expressed during cutaneous wound repair. <i>Experimental Cell Research</i> , <b>1999</b> , 247, 484-94                              | 4.2  | 117 |
| 95  | Nrf2--A regulator of keratinocyte redox signaling. <i>Free Radical Biology and Medicine</i> , <b>2015</b> , 88, 243-252   | 7.8  | 114 |
| 94  | Nrf2 establishes a glutathione-mediated gradient of UVB cytoprotection in the epidermis. <i>Genes and Development</i> , <b>2010</b> , 24, 1045-58                                 | 12.6 | 112 |
| 93  | Mouse fibroblast growth factor 10: cDNA cloning, protein characterization, and regulation of mRNA expression. <i>Oncogene</i> , <b>1997</b> , 15, 2211-8                          | 9.2  | 111 |
| 92  | Nrf transcription factors in keratinocytes are essential for skin tumor prevention but not for wound healing. <i>Molecular and Cellular Biology</i> , <b>2006</b> , 26, 3773-84   | 4.8  | 105 |
| 91  | The role of fibroblast growth factor receptor 2b in skin homeostasis and cancer development. <i>EMBO Journal</i> , <b>2007</b> , 26, 1268-78                                      | 13   | 102 |
| 90  | Fibroblast growth factor receptors 1 and 2 in keratinocytes control the epidermal barrier and cutaneous homeostasis. <i>Journal of Cell Biology</i> , <b>2010</b> , 188, 935-52   | 7.3  | 101 |
| 89  | Fibroblast growth factors and neuroprotection. <i>Advances in Experimental Medicine and Biology</i> , <b>2002</b> , 513, 335-51   | 3.6  | 91  |
| 88  | Peroxiredoxin 6 is a potent cytoprotective enzyme in the epidermis. <i>American Journal of Pathology</i> , <b>2006</b> , 169, 1194-205  | 5.8  | 89  |
| 87  | Activin a promotes the TGF-beta-induced conversion of CD4+CD25- T cells into Foxp3+ induced regulatory T cells. <i>Journal of Immunology</i> , <b>2009</b> , 182, 4633-40         | 5.3  | 85  |
| 86  | Nrf2-Mediated Fibroblast Reprogramming Drives Cellular Senescence by Targeting the Matrisome. <i>Developmental Cell</i> , <b>2018</b> , 46, 145-161.e10                           | 10.2 | 80  |
| 85  | Glucocorticoid-regulated gene expression during cutaneous wound repair. <i>Vitamins and Hormones</i> , <b>2000</b> , 59, 217-39   | 2.5  | 76  |
| 84  | The bright and the dark sides of activin in wound healing and cancer. <i>Journal of Cell Science</i> , <b>2012</b> , 125, 3929-37   | 5.3  | 72  |

|    |   |      |    |
|----|---|------|----|
| 83 | Activation of Nrf2 in keratinocytes causes chloracne (MADISH)-like skin disease in mice. <i>EMBO Molecular Medicine</i> , <b>2014</b> , 6, 442-57   | 12   | 71 |
| 82 | FGF receptors 1 and 2 are key regulators of keratinocyte migration in vitro and in wounded skin. <i>Journal of Cell Science</i> , <b>2012</b> , 125, 5690-701   | 5.3  | 70 |
| 81 | Fibroblast growth factor receptor signalling is crucial for liver homeostasis and regeneration. <i>Oncogene</i> , <b>2003</b> , 22, 4380-8  | 9.2  | 70 |
| 80 | Activin controls skin morphogenesis and wound repair predominantly via stromal cells and in a concentration-dependent manner via keratinocytes. <i>American Journal of Pathology</i> , <b>2005</b> , 167, 733-47                                      | 5.8  | 67 |
| 79 | Mast cells are dispensable for normal and activin-promoted wound healing and skin carcinogenesis. <i>Journal of Immunology</i> , <b>2013</b> , 191, 6147-55   | 5.3  | 62 |
| 78 | Activated hepatic stellate cells express keratinocyte growth factor in chronic liver disease. <i>American Journal of Pathology</i> , <b>2004</b> , 165, 1233-41   | 5.8  | 62 |
| 77 | Serum growth factors and proinflammatory cytokines are potent inducers of activin expression in cultured fibroblasts and keratinocytes. <i>Experimental Cell Research</i> , <b>1996</b> , 228, 106-13   | 4.2  | 62 |
| 76 | Activated Nrf2 impairs liver regeneration in mice by activation of genes involved in cell-cycle control and apoptosis. <i>Hepatology</i> , <b>2014</b> , 60, 670-8  | 11.2 | 61 |
| 75 | A Glutathione-Nrf2-Thioredoxin Cross-Talk Ensures Keratinocyte Survival and Efficient Wound Repair. <i>PLoS Genetics</i> , <b>2016</b> , 12, e1005800   | 6    | 58 |
| 74 | HMGB1 promotes ductular reaction and tumorigenesis in autophagy-deficient livers. <i>Journal of Clinical Investigation</i> , <b>2018</b> , 128, 2419-2435   | 15.9 | 56 |
| 73 | A role for endogenous glucocorticoids in wound repair. <i>EMBO Reports</i> , <b>2002</b> , 3, 575-82  | 6.5  | 54 |
| 72 | Expression of inflammasome proteins and inflammasome activation occurs in human, but not in murine keratinocytes. <i>Cell Death and Disease</i> , <b>2018</b> , 9, 24   | 9.8  | 53 |
| 71 | Control of hepatocyte proliferation and survival by Fgf receptors is essential for liver regeneration in mice. <i>Gut</i> , <b>2015</b> , 64, 1444-53   | 19.2 | 51 |
| 70 | Knockdown and knockout of $\beta$ -integrin in hepatocytes impairs liver regeneration through inhibition of growth factor signalling. <i>Nature Communications</i> , <b>2014</b> , 5, 3862  | 17.4 | 51 |
| 69 | Identification of novel AP-1 target genes in fibroblasts regulated during cutaneous wound healing. <i>Oncogene</i> , <b>2004</b> , 23, 7005-17  | 9.2  | 50 |
| 68 | Loss of serum response factor in keratinocytes results in hyperproliferative skin disease in mice. <i>Journal of Clinical Investigation</i> , <b>2009</b> , 119, 899-910  | 15.9 | 48 |
| 67 | Dual role of the antioxidant enzyme peroxiredoxin 6 in skin carcinogenesis. <i>Cancer Research</i> , <b>2013</b> , 73, 3460-9   | 10.1 | 47 |
| 66 | Down-regulation of connective tissue growth factor and type I collagen mRNA expression by connective tissue growth factor antisense oligonucleotide during experimental liver fibrosis. <i>Wound Repair and Regeneration</i> , <b>2004</b> , 12, 60-6 | 3.6  | 47 |

|    |   |      |    |
|----|---|------|----|
| 65 | Amniotic fluid activates the nrf2/keap1 pathway to repair an epidermal barrier defect in utero. <i>Developmental Cell</i> , <b>2012</b> , 23, 1238-46   | 10.2 | 46 |
| 64 | A novel Nrf2-miR-29-desmocollin-2 axis regulates desmosome function in keratinocytes. <i>Nature Communications</i> , <b>2014</b> , 5, 5099  | 17.4 | 45 |
| 63 | Activin enhances skin tumourigenesis and malignant progression by inducing a pro-tumourigenic immune cell response. <i>Nature Communications</i> , <b>2011</b> , 2, 576   | 17.4 | 42 |
| 62 | FGF receptors 1 and 2 control chemically induced injury and compound detoxification in regenerating livers of mice. <i>Gastroenterology</i> , <b>2010</b> , 139, 1385-96  | 13.3 | 41 |
| 61 | Opposing effects of Nrf2 and Nrf2-activating compounds on the NLRP3 inflammasome independent of Nrf2-mediated gene expression. <i>European Journal of Immunology</i> , <b>2017</b> , 47, 806-817  | 6.1  | 39 |
| 60 | Electrophilic chemicals but not UV irradiation or reactive oxygen species activate Nrf2 in keratinocytes in vitro and in vivo. <i>Journal of Investigative Dermatology</i> , <b>2007</b> , 127, 646-53  | 4.3  | 39 |
| 59 | Beta1 integrin-mediated adhesion signalling is essential for epidermal progenitor cell expansion. <i>PLoS ONE</i> , <b>2009</b> , 4, e5488  | 3.7  | 38 |
| 58 | The commensal skin microbiota triggers type I IFN-dependent innate repair responses in injured skin. <i>Nature Immunology</i> , <b>2020</b> , 21, 1034-1045   | 19.1 | 38 |
| 57 | NRF2 and microRNAs: new but awaited relations. <i>Biochemical Society Transactions</i> , <b>2015</b> , 43, 595-601  | 5.1  | 36 |
| 56 | Nrf2 Activation Promotes Keratinocyte Survival during Early Skin Carcinogenesis via Metabolic Alterations. <i>Cancer Research</i> , <b>2015</b> , 75, 4817-29   | 10.1 | 36 |
| 55 | The cytoprotective Nrf2 transcription factor controls insulin receptor signaling in the regenerating liver. <i>Cell Cycle</i> , <b>2008</b> , 7, 874-8  | 4.7  | 35 |
| 54 | Activation of the Nrf2-ARE pathway in hepatocytes protects against steatosis in nutritionally induced non-alcoholic steatohepatitis in mice. <i>Toxicological Sciences</i> , <b>2014</b> , 142, 361-74  | 4.4  | 30 |
| 53 | Transcriptional regulation of wound inflammation. <i>Seminars in Immunology</i> , <b>2014</b> , 26, 321-8   | 10.7 | 27 |
| 52 | Sulforaphane homologues: Enantiodivergent synthesis of both enantiomers, activation of the Nrf2 transcription factor and selective cytotoxic activity. <i>European Journal of Medicinal Chemistry</i> , <b>2014</b> , 87, 552-63                            | 6.8  | 26 |
| 51 | NF- $\kappa$ B/RelA and Nrf2 cooperate to maintain hepatocyte integrity and to prevent development of hepatocellular adenoma. <i>Journal of Hepatology</i> , <b>2016</b> , 64, 94-102   | 13.4 | 23 |
| 50 | Regulatory T cells are required for normal and activin-promoted wound repair in mice. <i>European Journal of Immunology</i> , <b>2018</b> , 48, 1001-1013   | 6.1  | 23 |
| 49 | Identification of UV-protective activators of nuclear factor erythroid-derived 2-related factor 2 (Nrf2) by combining a chemical library screen with computer-based virtual screening. <i>Journal of Biological Chemistry</i> , <b>2012</b> , 287, 33001-13 | 5.4  | 23 |
| 48 | Keratinocyte-derived follistatin regulates epidermal homeostasis and wound repair. <i>Laboratory Investigation</i> , <b>2009</b> , 89, 131-41   | 5.9  | 22 |

|    |  |      |    |
|----|--|------|----|
| 47 | Langerhans cells are strongly reduced in the skin of transgenic mice overexpressing follistatin in the epidermis. <i>European Journal of Cell Biology</i> , <b>2005</b> , 84, 733-41   | 6.1  | 22 |
| 46 | A paracrine activin A-mDia2 axis promotes squamous carcinogenesis via fibroblast reprogramming. <i>EMBO Molecular Medicine</i> , <b>2020</b> , 12, e11466  | 12   | 21 |
| 45 | Laminin 5 in the keratinocyte basement membrane is required for epidermal-dermal intercommunication. <i>Matrix Biology</i> , <b>2016</b> , 56, 24-41   | 11.4 | 20 |
| 44 | Activin-mediated alterations of the fibroblast transcriptome and matrisome control the biomechanical properties of skin wounds. <i>Nature Communications</i> , <b>2020</b> , 11, 2604  | 17.4 | 20 |
| 43 | A novel enhancer of the wound healing process: the fibroblast growth factor-binding protein. <i>American Journal of Pathology</i> , <b>2011</b> , 179, 2144-7  | 5.8  | 19 |
| 42 | Wound healing studies in transgenic and knockout mice. A review. <i>Methods in Molecular Medicine</i> , <b>2003</b> , 78, 191-216  |      | 19 |
| 41 | Overactivation of the nuclear factor (erythroid-derived 2)-like 2-antioxidant response element pathway in hepatocytes decreases hepatic ischemia/reperfusion injury in mice. <i>Liver Transplantation</i> , <b>2016</b> , 22, 91-102 | 4.5  | 18 |
| 40 | Activin promotes skin carcinogenesis by attraction and reprogramming of macrophages. <i>EMBO Molecular Medicine</i> , <b>2017</b> , 9, 27-45   | 12   | 17 |
| 39 | The mechanical fingerprint of murine excisional wounds. <i>Acta Biomaterialia</i> , <b>2018</b> , 65, 226-236  | 10.8 | 17 |
| 38 | Regulation of Wound Healing by the NRF2 Transcription Factor-More Than Cytoprotection. <i>International Journal of Molecular Sciences</i> , <b>2019</b> , 20,  | 6.3  | 16 |
| 37 | Cell-specific Activation of the Nrf2 Antioxidant Pathway Increases Mucosal Inflammation in Acute but Not in Chronic Colitis. <i>Journal of Crohn's and Colitis</i> , <b>2017</b> , 11, 485-499                                       | 1.5  | 16 |
| 36 | Nrf2 is highly expressed in neutrophils, but myeloid cell-derived Nrf2 is dispensable for wound healing in mice. <i>PLoS ONE</i> , <b>2017</b> , 12, e0187162  | 3.7  | 14 |
| 35 | Exosomes for Wound Healing: Purification Optimization and Identification of Bioactive Components. <i>Advanced Science</i> , <b>2020</b> , 7, 2002596   | 13.6 | 14 |
| 34 | Tussilagonone Ameliorates Psoriatic Features in Keratinocytes and Imiquimod-Induced Psoriasis-Like Lesions in Mice via NRF2 Activation. <i>Journal of Investigative Dermatology</i> , <b>2020</b> , 140, 1223-1232.e4                | 4.3  | 13 |
| 33 | Kdm6b and Pmpa1 as Targets of Bioelectrically and Behaviorally Induced Activin A Signaling. <i>Molecular Neurobiology</i> , <b>2016</b> , 53, 4210-4225  | 6.2  | 12 |
| 32 | Comprehensive characterization of myeloid cells during wound healing in healthy and healing-impaired diabetic mice. <i>European Journal of Immunology</i> , <b>2020</b> , 50, 1335-1349  | 6.1  | 12 |
| 31 | Humidity-regulated CLCA2 protects the epidermis from hyperosmotic stress. <i>Science Translational Medicine</i> , <b>2018</b> , 10,  | 17.5 | 12 |
| 30 | Mast cells are dispensable in a genetic mouse model of chronic dermatitis. <i>American Journal of Pathology</i> , <b>2015</b> , 185, 1575-87   | 5.8  | 11 |

|    |  |      |    |
|----|--|------|----|
| 29 | Autocrine and Paracrine Regulation of Keratinocyte Proliferation through a Novel Nrf2-IL-36 $\beta$ Pathway. <i>Journal of Immunology</i> , <b>2016</b> , 196, 4663-70   | 5.3  | 11 |
| 28 | Nrf3 promotes UV-induced keratinocyte apoptosis through suppression of cell adhesion. <i>Cell Death and Differentiation</i> , <b>2018</b> , 25, 1749-1765  | 12.7 | 10 |
| 27 | Targeting metabolism to treat psoriasis. <i>Nature Medicine</i> , <b>2018</b> , 24, 537-539  | 50.5 | 10 |
| 26 | Large-Scale Quantitative Proteomics Identifies the Ubiquitin Ligase Nedd4-1 as an Essential Regulator of Liver Regeneration. <i>Developmental Cell</i> , <b>2017</b> , 42, 616-625.e8                                      | 10.2 | 10 |
| 25 | Wound Repair, Scar Formation, and Cancer: Converging on Activin. <i>Trends in Molecular Medicine</i> , <b>2020</b> , 26, 1107-1117   | 11.5 | 10 |
| 24 | Cell-specific overactivation of nuclear erythroid 2 p45-related factor 2-mediated gene expression in myeloid cells decreases hepatic ischemia/reperfusion injury. <i>Liver Transplantation</i> , <b>2016</b> , 22, 1115-28 | 4.5  | 9  |
| 23 | Long-Term Imaging of Wound Angiogenesis with Large Scale Optoacoustic Microscopy. <i>Advanced Science</i> , <b>2021</b> , 8, 2004226   | 13.6 | 9  |
| 22 | The NLRP1 Inflammasome Pathway Is Silenced in Cutaneous Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , <b>2019</b> , 139, 1788-1797.e6   | 4.3  | 8  |
| 21 | Accumulation and activation of epidermal $\gamma\delta$ T cells in a mouse model of chronic dermatitis is not required for the inflammatory phenotype. <i>European Journal of Immunology</i> , <b>2015</b> , 45, 2517-28   | 6.1  | 7  |
| 20 | Nrf2-Mediated Expansion of Pilosebaceous Cells Accelerates Cutaneous Wound Healing. <i>American Journal of Pathology</i> , <b>2019</b> , 189, 568-579  | 5.8  | 7  |
| 19 | Genetic activation of Nrf2 reduces cutaneous symptoms in a murine model of Netherton syndrome. <i>DMM Disease Models and Mechanisms</i> , <b>2020</b> , 13,  | 4.1  | 5  |
| 18 | A modeling approach to study the effect of cell polarization on keratinocyte migration. <i>PLoS ONE</i> , <b>2015</b> , 10, e0117676   | 3.7  | 5  |
| 17 | Mouse genetics identifies unique and overlapping functions of fibroblast growth factor receptors in keratinocytes. <i>Journal of Cellular and Molecular Medicine</i> , <b>2020</b> , 24, 1774-1785                         | 5.6  | 5  |
| 16 | Low levels of glutathione are sufficient for survival of keratinocytes after UV irradiation and for healing of mouse skin wounds. <i>Archives of Dermatological Research</i> , <b>2016</b> , 308, 443-8                    | 3.3  | 4  |
| 15 | Antagonism of interferon signaling by fibroblast growth factors promotes viral replication. <i>EMBO Molecular Medicine</i> , <b>2020</b> , 12, e11793  | 12   | 4  |
| 14 | Interaction of the NRF2 and p63 transcription factors promotes keratinocyte proliferation in the epidermis. <i>Nucleic Acids Research</i> , <b>2021</b> , 49, 3748-3763  | 20.1 | 4  |
| 13 | A Dual-Acting Nitric Oxide Donor and Phosphodiesterase 5 Inhibitor Promotes Wound Healing in Normal Mice and Mice with Diabetes. <i>Journal of Investigative Dermatology</i> , <b>2021</b> , 141, 415-426                  | 4.3  | 4  |
| 12 | Imaging and targeting LOX-mediated tissue remodeling with a reactive collagen peptide. <i>Nature Chemical Biology</i> , <b>2021</b> , 17, 865-871  | 11.7 | 4  |

|    |  |      |   |
|----|--|------|---|
| 11 | Non-invasive longitudinal imaging of VEGF-induced microvascular alterations in skin wounds..<br><i>Theranostics</i> , <b>2022</b> , 12, 558-573  | 12.1 | 3 |
| 10 | Molecular and cellular mechanisms of tissue repair. <i>Experimental Dermatology</i> , <b>2005</b> , 14, 786-7  | 4    | 2 |
| 9  | Acute and chronic effects of a light-activated FGF receptor in keratinocytes in vitro and in mice. <i>Life Science Alliance</i> , <b>2021</b> , 4,   | 5.8  | 2 |
| 8  | Peroxiredoxin 6 in skin carcinogenesis. <i>Oncoscience</i> , <b>2014</b> , 1, 392-3  | 0.8  | 1 |
| 7  | Stromal-epithelial interactions in skin homeostasis, wound repair and skin cancer. <i>Experimental Dermatology</i> , <b>2008</b> , 17, 882-883   | 4    | 1 |
| 6  | Genotoxic Agents: An Unexpected Effect on Healthy Epithelia. <i>Developmental Cell</i> , <b>2020</b> , 55, 515-517   | 10.2 | 0 |
| 5  | Tissue Repair: Guarding against Friendly Fire. <i>Current Biology</i> , <b>2019</b> , 29, R1191-R1193  | 6.3  | 0 |
| 4  | Fibroblast growth factor receptor 3 in hepatocytes protects from toxin-induced liver injury and fibrosis. <i>iScience</i> , <b>2021</b> , 24, 103143   | 6.1  | 0 |
| 3  | Fibroblast Growth Factors in Epithelial Homeostasis and Repair <b>2017</b> , 187-209   |      |   |
| 2  | A Phase 1 Single Dose Escalation Study of Palifermin Administered Pre-Transplant Conditioning in Subjects Undergoing Matched Unrelated Donor Allogeneic Hematopoietic Stem Cell Transplantation. <i>Blood</i> , <b>2020</b> , 136, 21-21 | 2.2  |   |
| 1  | p62 Promotes Survival and Hepatocarcinogenesis in Mice with Liver-Specific NEMO Ablation. <i>Cancers</i> , <b>2022</b> , 14, 2436  | 6.6  |   |