

# James A Coyer

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11484210/publications.pdf>

Version: 2024-02-01

39  
papers

2,255  
citations

279798

23  
h-index

315739

38  
g-index

41  
all docs

41  
docs citations

41  
times ranked

2375  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Seagrass Methylome Is Associated With Variation in Photosynthetic Performance Among Clonal Shoots. <i>Frontiers in Plant Science</i> , 2020, 11, 571646.	3.6	21
2	Decadal stability in genetic variation and structure in the intertidal seaweed <i>Fucus serratus</i> (Heterokontophyta: Fucaceae). <i>BMC Evolutionary Biology</i> , 2018, 18, 94.	3.2	10
3	Unravelling the complexity of salt marsh <i>Fucus cottonii</i> forms (Phaeophyceae, Fucales). <i>European Journal of Phycology</i> , 2017, 52, 360-370.	2.0	9
4	The fate of the Arctic seaweed <i>Fucus distichus</i> under climate change: an ecological niche modeling approach. <i>Ecology and Evolution</i> , 2016, 6, 1712-1724.	1.9	109
5	Variation in thermal stress response in two populations of the brown seaweed, <i>Fucus distichus</i> , from the Arctic and subarctic intertidal. <i>Royal Society Open Science</i> , 2016, 3, 150429.	2.4	26
6	Climate Oscillations, Range Shifts and Phylogeographic Patterns of North Atlantic Fucaceae. , 2016, , 279-308.		27
7	Seascape drivers of <i>Monocystis pyrifera</i> population genetic structure in the northeast Pacific. <i>Molecular Ecology</i> , 2015, 24, 4866-4885.	3.9	55
8	Genome-wide transcriptomic responses of the seagrasses <i>Zostera marina</i> and <i>Nanozostera noltii</i> under a simulated heatwave confirm functional types. <i>Marine Genomics</i> , 2014, 15, 65-73.	1.1	68
9	Distribution patterns and introduction pathways of the cosmopolitan brown alga <i>Colpomenia peregrina</i> using mt <i>cox3</i> and <i>atp6</i> sequences. <i>Journal of Applied Phycology</i> , 2014, 26, 491-504.	2.8	15
10	Thermal stress resistance of the brown alga <i>Fucus serratus</i> along the North-Atlantic coast: Acclimatization potential to climate change. <i>Marine Genomics</i> , 2014, 13, 27-36.	1.1	57
11	Numerous mitigation transplants of the eelgrass <i>Zostera marina</i> in southern California shuffle genetic diversity and may promote hybridization with <i>Zostera pacifica</i> . <i>Biological Conservation</i> , 2014, 176, 133-143.	4.1	15
12	Climate change impact on seaweed meadow distribution in the North Atlantic rocky intertidal. <i>Ecology and Evolution</i> , 2013, 3, 1356-1373.	1.9	170
13	Reply to Hu and Duan (Mar Biol): Insufficient geographical sampling could severely influence phylogeographic interpretations; comment on "Phylogeography of the seaweed <i>Ishige okamurae</i> (Phaeophyceae): evidence for glacial refugia in the northwest Pacific region" (Lee et al. 2012). <i>Marine Biology</i> . 2013. 160. 1519-1520.	1.5	1
14	Phylogeography of the seaweed <i>Ishige okamurae</i> (Phaeophyceae): evidence for glacial refugia in the northwest Pacific region. <i>Marine Biology</i> , 2012, 159, 1021-1028.	1.5	40
15	Stress Ecology in <i>Fucus</i> : Abiotic, Biotic and Genetic Interactions. <i>Advances in Marine Biology</i> , 2011, 59, 37-105.	1.4	95
16	Trans-Pacific and trans-Arctic pathways of the intertidal macroalga <i>Fucus distichus</i> L. reveal multiple glacial refugia and colonizations from the North Pacific to the North Atlantic. <i>Journal of Biogeography</i> , 2011, 38, 756-771.	3.0	58
17	An evaluation of small-scale genetic diversity and the mating system in <i>Zostera noltii</i> on an intertidal sandflat in the Wadden Sea. <i>Annals of Botany</i> , 2011, 107, 127-134.	2.9	33
18	An Expressed Sequence Tag Analysis of the Intertidal Brown Seaweeds <i>Fucus serratus</i> (L.) and <i>F. vesiculosus</i> (L.) (Heterokontophyta, Phaeophyceae) in Response to Abiotic Stressors. <i>Marine Biotechnology</i> , 2010, 12, 195-213.	2.4	77

#	ARTICLE	IF	CITATIONS
19	The phylogeographic architecture of the fucoid seaweed <i>Ascophyllum nodosum</i> : an intertidal marine tree and survivor of more than one glacial-interglacial cycle. <i>Journal of Biogeography</i> , 2010, 37, 842-856.	3.0	93
20	Waterfowl grazing in autumn enhances spring seedling recruitment of intertidal <i>Zostera noltii</i> . <i>Aquatic Botany</i> , 2010, 93, 202-205.	1.6	16
21	Historical invasions of the intertidal zone of Atlantic North America associated with distinctive patterns of trade and emigration. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 8239-8244.	7.1	73
22	Clonal architecture in an intertidal bed of the dwarf eelgrass <i>Zostera noltii</i> in the Northern Wadden Sea: persistence through extreme physical perturbation and the importance of a seed bank. <i>Marine Biology</i> , 2009, 156, 2139-2148.	1.5	22
23	PATERNAL LEAKAGE OF MITOCHONDRIAL DNA IN A <i>FUCUS</i> (PHAEOPHYCEAE) HYBRID ZONE <sup>1</sup> . <i>Journal of Phycology</i> , 2009, 45, 621-624.	2.3	15
24	Being abundant is not enough: a decrease in effective population size over eight generations in a Norwegian population of the seaweed, <i>Fucus serratus</i> . <i>Biology Letters</i> , 2008, 4, 755-757.	2.3	15
25	A fast and inexpensive DNA extraction/purification protocol for brown macroalgae. <i>Molecular Ecology Notes</i> , 2007, 7, 191-193.	1.7	56
26	Convergent adaptation to a marginal habitat by homoploid hybrids and polyploid ecads in the seaweed genus <i>Fucus</i> . <i>Biology Letters</i> , 2006, 2, 405-408.	2.3	54
27	A mtDNA-based phylogeny of the brown algal genus <i>Fucus</i> (Heterokontophyta; Phaeophyta). <i>Molecular Phylogenetics and Evolution</i> , 2006, 39, 209-222.	2.7	83
28	North Atlantic phylogeography and large-scale population differentiation of the seagrass <i>Zostera marina</i> L.. <i>Molecular Ecology</i> , 2004, 13, 1923-1941.	3.9	277
29	Title is missing!. <i>Hydrobiologia</i> , 2003, 501, 149-166.	2.0	63
30	COMMUNITY-WIDE EFFECTS OF NONINDIGENOUS SPECIES ON TEMPERATE ROCKY REEFS. <i>Ecology</i> , 2002, 83, 3182-3193.	3.2	181
31	Asexual propagation in the coral reef macroalga <i>Halimeda</i> (Chlorophyta, Bryopsidales): production, dispersal and attachment of small fragments. <i>Journal of Experimental Marine Biology and Ecology</i> , 2002, 278, 47-65.	1.5	44
32	EVOLUTION OF MACROCYSTIS SPP. (PHAEOPHYCEAE) AS DETERMINED BY ITS1 AND ITS2 SEQUENCES <sup>1</sup> . <i>Journal of Phycology</i> , 2001, 37, 574-585.	2.3	113
33	GENETIC VARIABILITY AND SPATIAL SEPARATION IN THE SEA PALM KELP <i>POSTELSIA PALMAEFORMIS</i> (PHAEOPHYCEAE) AS ASSESSED WITH M13 FINGERPRINTS AND RAPDS <sup>1</sup> . <i>Journal of Phycology</i> , 1997, 33, 561-568.	2.3	64
34	A FIELD-COMPATIBLE METHOD FOR EXTRACTION OF FINGERPRINT-QUALITY DNA FROM MACROCYSTIS PYRIFERA (PHAEOPHYCEAE) <sup>1</sup> . <i>Journal of Phycology</i> , 1995, 31, 177-180.	2.3	9
35	GENETIC VARIABILITY AND PARENTAGE IN MACROCYSTIS PYRIFERA (PHAEOPHYCEAE) USING MULTI-LOCUS DNA FINGERPRINTING <sup>1</sup> . <i>Journal of Phycology</i> , 1995, 31, 819-823.	2.3	8
36	GENETIC VARIABILITY WITHIN A POPULATION AND BETWEEN DIPLOID/HAPLOID TISSUE OF MACROCYSTIS PYRIFERA (PHAEOPHYCEAE) <sup>1</sup> . <i>Journal of Phycology</i> , 1994, 30, 545-552.	2.3	41

#	ARTICLE	IF	CITATIONS
37	Interactions between corals and algae on a temperate zone rocky reef: mediation by sea urchins. Journal of Experimental Marine Biology and Ecology, 1993, 167, 21-37.	1.5	82
38	DISCOVERY OF A FERTILE PELAGOPHYCUSxMACROCYSTIS(PHAEOPHYTA) PUTATIVE HYBRID AND SUBSEQUENT PRODUCTION OF F2 SPOROPHYTES IN THE LABORATORY1. Journal of Phycology, 1992, 28, 127-130.	2.3	16
39	Utilization of purple and red sea urchins ( <i>Strongylocentrotus purpuratus</i> Stimpson and S.) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf laboratory. Journal of Experimental Marine Biology and Ecology, 1987, 105, 21-38.	1.5	15