
Maurice Ombok

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11529311/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	COMMUNITY-WIDE EFFECTS OF PERMETHRIN-TREATED BED NETS ON CHILD MORTALITY AND MALARIA MORBIDITY IN WESTERN KENYA. American Journal of Tropical Medicine and Hygiene, 2003, 68, 121-127.	1.4	450
2	Characteristics of Larval Anopheline (Diptera: Culicidae) Habitats in Western Kenya. Journal of Medical Entomology, 2001, 38, 282-288.	1.8	280
3	Density-Dependent Development of <i>Anopheles gambiae</i> (Diptera: Culicidae) Larvae in Artificial Habitats. Journal of Medical Entomology, 2002, 39, 162-172.	1.8	245
4	Profile: The KEMRI/CDC Health and Demographic Surveillance SystemWestern Kenya. International Journal of Epidemiology, 2012, 41, 977-987.	1.9	199
5	The impact of distance of residence from a peripheral health facility on pediatric health utilisation in rural western Kenya. Tropical Medicine and International Health, 2009, 14, 54-61.	2.3	159
6	Persistently high estimates of late night, indoor exposure to malaria vectors despite high coverage of insecticide treated nets. Parasites and Vectors, 2014, 7, 380.	2.5	106
7	Effect of Permethrin-Impregnated Nets on Exiting Behavior, Blood Feeding Success, and Time of Feeding of Malaria Mosquitoes (Diptera: Culicidae) in Western Kenya. Journal of Medical Entomology, 2001, 38, 531-536.	1.8	92
8	Water, Sanitation and Hygiene Conditions in Kenyan Rural Schools: Are Schools Meeting the Needs of Menstruating Girls?. Water (Switzerland), 2014, 6, 1453-1466.	2.7	55
9	Pyrethroid susceptibility of malaria vectors in four Districts of western Kenya. Parasites and Vectors, 2014, 7, 310.	2.5	54
10	Insecticide-Treated Nets and Protection against Insecticide-Resistant Malaria Vectors in Western Kenya. Emerging Infectious Diseases, 2017, 23, 758-764.	4.3	41
11	The Relationship Between Distance to Water Source and Moderate-to-Severe Diarrhea in the Global Enterics Multi-Center Study in Kenya, 2008–2011. American Journal of Tropical Medicine and Hygiene, 2016, 94, 1143-1149.	1.4	36
12	Diagnostic dose determination and efficacy of chlorfenapyr and clothianidin insecticides against Anopheles malaria vector populations of western Kenya. Malaria Journal, 2019, 18, 243.	2.3	28
13	Host Decoy Trap (HDT) with cattle odour is highly effective for collection of exophagic malaria vectors. Parasites and Vectors, 2018, 11, 533.	2.5	24
14	Geospatial distribution and determinants of child mortality in rural western Kenya 2002-2005. Tropical Medicine and International Health, 2010, 15, 423-433.	2.3	23
15	Geographic distribution of HIV stigma among women of childbearing age in rural Kenya. Aids, 2014, 28, 1665-1672.	2.2	11
16	Microdam Impoundments Provide Suitable Habitat for Larvae of Malaria Vectors: An Observational Study in Western Kenya. Journal of Medical Entomology, 2018, 55, 723-730.	1.8	7
17	Community-based intermittent mass testing and treatment for malaria in an area of high transmission intensity, western Kenya: development of study site infrastructure and lessons learned. Malaria Journal, 2019, 18, 255.	2.3	7
18	Efficacy of extended release formulations of Natularâ,,¢ (spinosad) against larvae and adults of Anopheles mosquitoes in western Kenya. Malaria Journal, 2020, 19, 436.	2.3	5