

Ivana Nedeljkovic

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11532281/publications.pdf>

Version: 2024-02-01

24
papers

563
citations

840776

11
h-index

677142

22
g-index

24
all docs

24
docs citations

24
times ranked

666
citing authors

#	ARTICLE	IF	CITATIONS
1	Aortic Valve Replacement Versus Conservative Treatment in Asymptomatic Severe Aortic Stenosis: The AVATAR Trial. <i>Circulation</i> , 2022, 145, 648-658.	1.6	130
2	Combined low dose dipyridamole-dobutamine stress echocardiography to identify myocardial viability. <i>Journal of the American College of Cardiology</i> , 1996, 27, 1422-1428.	2.8	71
3	The combined exercise stress echocardiography and cardiopulmonary exercise test for identification of masked heart failure with preserved ejection fraction in patients with hypertension. <i>European Journal of Preventive Cardiology</i> , 2016, 23, 71-77.	1.8	58
4	Rationale and design of the Aortic Valve replAcemenT versus conservative treatment in Asymptomatic seveRe aortic stenosis (AVATAR trial): A randomized multicenter controlled event-driven trial. <i>American Heart Journal</i> , 2016, 174, 147-153.	2.7	55
5	Integrated evaluation of relation between coronary lesion features and stress echocardiography results: the importance of coronary lesion morphology. <i>Journal of the American College of Cardiology</i> , 1999, 33, 717-726.	2.8	51
6	Acute insulin resistance in ST-segment elevation myocardial infarction in non-diabetic patients is associated with incomplete myocardial reperfusion and impaired coronary microcirculatory function. <i>Cardiovascular Diabetology</i> , 2014, 13, 73.	6.8	37
7	The Amount of Weight Loss Six Months after Bariatric Surgery: It Makes a Difference. <i>Obesity Facts</i> , 2019, 12, 281-290.	3.4	17
8	Prediction of Myocardial Functional Recovery by Noninvasive Evaluation of Basal and Hyperemic Coronary Flow in Patients with Previous Myocardial Infarction. <i>Journal of the American Society of Echocardiography</i> , 2011, 24, 573-581.	2.8	15
9	Efficiency of ergonovine echocardiography in detecting angiographically assessed coronary vasospasm. <i>American Journal of Cardiology</i> , 2001, 88, 1183-1187.	1.6	14
10	Time-dependent changes of plasma adiponectin concentration in relation to coronary microcirculatory function in patients with acute myocardial infarction treated by primary percutaneous coronary intervention. <i>Journal of Cardiology</i> , 2015, 65, 208-215.	1.9	14
11	Impairment of coronary flow velocity reserve and global longitudinal strain in women with cardiac syndrome X and slow coronary flow. <i>Journal of Cardiology</i> , 2020, 76, 1-8.	1.9	14
12	Diagnostic value of NT-proBNP in identifying impaired coronary flow reserve in asymptomatic moderate or severe aortic stenosis. <i>Biomarkers in Medicine</i> , 2013, 7, 221-227.	1.4	13
13	Low-dose adenosine stress echocardiography: Detection of myocardial viability. <i>Cardiovascular Ultrasound</i> , 2003, 1, 7.	1.6	11
14	Prognostic Value of Preserved Coronary Flow Velocity Reserve by Noninvasive Transthoracic Doppler Echocardiography in Patients With Angiographically Intermediate Left Main Stenosis. <i>Journal of the American Society of Echocardiography</i> , 2019, 32, 74-80.	2.8	11
15	Diabetes mellitus and coronary microvascular function in asymptomatic patients with severe aortic stenosis and nonobstructed coronary arteries. <i>Diabetes and Vascular Disease Research</i> , 2016, 13, 220-227.	2.0	10
16	Valvulo-arterial impedance is the best mortality predictor in asymptomatic aortic stenosis patients. <i>Journal of Heart Valve Disease</i> , 2015, 24, 156-63.	0.5	9
17	Silent coronary artery disease in asymptomatic patients with severe aortic stenosis and normal exercise testing. <i>Coronary Artery Disease</i> , 2020, 31, 166-173.	0.7	7
18	Ergonovine-Induced Changes of Coronary Artery Diameter in Patients with Nonsignificant Coronary Artery Stenosis. <i>Herz</i> , 2007, 32, 329-335.	1.1	6

#	ARTICLE	IF	CITATIONS
19	Coronary flow of the infarct artery assessed by transthoracic Doppler after primary percutaneous coronary intervention predicts final infarct size. <i>International Journal of Cardiovascular Imaging</i> , 2014, 30, 1509-1518.	1.5	6
20	Prediction of a Good Response to Cardiac Resynchronization Therapy in Patients with Severe Dilated Cardiomyopathy: Could Conventional Echocardiography Be the Answer after All?. <i>Echocardiography</i> , 2012, 29, 267-275.	0.9	5
21	Five-Year Outcomes in Bariatric Surgery Patients. <i>Medicina (Lithuania)</i> , 2020, 56, 669.	2.0	5
22	Gauging the response to cardiac resynchronization therapy: The important interplay between predictor variables and definition of a favorable outcome. <i>Echocardiography</i> , 2017, 34, 371-375.	0.9	4
23	Gender specific differences in functional capacity in asymptomatic patients with severe aortic stenosis. <i>Cor Et Vasa</i> , 2021, 63, 333-338.	0.1	0
24	Cardiopulmonary Exercise Test in the Detection of Unexplained Post-COVID-19 Dyspnea. <i>International Heart Journal</i> , 2021, 62, 1164-1170.	1.0	0