

Amy Cohen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11545121/publications.pdf>

Version: 2024-02-01

23
papers

2,029
citations

331259

21
h-index

642321

23
g-index

23
all docs

23
docs citations

23
times ranked

780
citing authors

#	ARTICLE	IF	CITATIONS
1	A Clinical Trial of Active Management of Labor. <i>New England Journal of Medicine</i> , 1995, 333, 745-750.	13.9	277
2	Intrapartum Maternal Fever and Neonatal Outcome. <i>Pediatrics</i> , 2000, 105, 8-13.	1.0	219
3	Uterine rupture during induced or augmented labor in gravid women with one prior cesarean delivery. <i>American Journal of Obstetrics and Gynecology</i> , 1999, 181, 882-886.	0.7	193
4	Rate of uterine rupture during a trial of labor in women with one or two prior cesarean deliveries. <i>American Journal of Obstetrics and Gynecology</i> , 1999, 181, 872-876.	0.7	156
5	Association of epidural analgesia with cesarean delivery in nulliparas. <i>Obstetrics and Gynecology</i> , 1996, 88, 993-1000.	1.2	155
6	Intrapartum Fever and Unexplained Seizures in Term Infants. <i>Pediatrics</i> , 2000, 106, 983-988.	1.0	142
7	Effect of previous vaginal delivery on the risk of uterine rupture during a subsequent trial of labor. <i>American Journal of Obstetrics and Gynecology</i> , 2000, 183, 1184-1186.	0.7	130
8	Outcomes of trial of labor following previous cesarean delivery among women with fetuses weighing >4000 g. <i>American Journal of Obstetrics and Gynecology</i> , 2001, 185, 903-905.	0.7	110
9	Interdelivery interval and risk of symptomatic uterine rupture. <i>Obstetrics and Gynecology</i> , 2001, 97, 175-177.	1.2	96
10	Trial of labor after cesarean delivery: The effect of previous vaginal delivery. <i>American Journal of Obstetrics and Gynecology</i> , 1998, 179, 938-941.	0.7	84
11	Nulliparas' Preferences for Epidural Analgesia: Their Effects on Actual Use in Labor. <i>Birth</i> , 1999, 26, 139-143.	1.1	60
12	Interdelivery Interval and Risk of Symptomatic Uterine Rupture. <i>Obstetrics and Gynecology</i> , 2001, 97, 175-177.	1.2	60
13	Intrapartum uterine rupture and dehiscence in patients with prior lower uterine segment vertical and transverse incisions ¹ . <i>Obstetrics and Gynecology</i> , 1999, 94, 735-740.	1.2	58
14	The association of maternal age and symptomatic uterine rupture during a trial of labor after prior cesarean delivery. <i>Obstetrics and Gynecology</i> , 2002, 99, 585-588.	1.2	54
15	Intrapartum Uterine Rupture and Dehiscence in Patients With Prior Lower Uterine Segment Vertical and Transverse Incisions. <i>Obstetrics and Gynecology</i> , 1999, 94, 735-740.	1.2	42
16	Trial of labor after 40 weeks' gestation in women with prior cesarean. <i>Obstetrics and Gynecology</i> , 2001, 97, 391-393.	1.2	42
17	Oxytocin dose and the risk of uterine rupture in trial of labor after cesarean. <i>Obstetrics and Gynecology</i> , 2001, 97, 381-384.	1.2	36
18	Maternal Epidural Analgesia and Rates of Maternal Antibiotic Treatment in a Low-risk Nulliparous Population. <i>Journal of Perinatology</i> , 2003, 23, 457-461.	0.9	34

#	ARTICLE	IF	CITATIONS
19	Trial of Labor After 40 Weeks¼ Gestation in Women With Prior Cesarean. <i>Obstetrics and Gynecology</i> , 2001, 97, 391-393.	1.2	29
20	Neonatal Outcome after Trial of Labor Compared with Elective Repeat Cesarean Section. <i>Birth</i> , 2003, 30, 83-88.	1.1	23
21	Oxytocin Dose and the Risk of Uterine Rupture in Trial of Labor After Cesarean. <i>Obstetrics and Gynecology</i> , 2001, 97, 381-384.	1.2	22
22	Epidurals and Cesareans: The Jury Is Still Out. <i>Birth</i> , 1999, 26, 196-198.	1.1	6
23	The Association of Maternal Age and Symptomatic Uterine Rupture During a Trial of Labor After Prior Cesarean Delivery. <i>Obstetrics and Gynecology</i> , 2002, 99, 585-588.	1.2	1