Shunichi Fukuhara

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11555338/publications.pdf

Version: 2024-02-01

363 15,192 57 111 papers citations h-index g-index

373 373 373 15533
all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Translation, Adaptation, and Validation of the SF-36 Health Survey for Use in Japan. Journal of Clinical Epidemiology, 1998, 51, 1037-1044.	2.4	1,146
2	Psychometric and Clinical Tests of Validity of the Japanese SF-36 Health Survey. Journal of Clinical Epidemiology, 1998, 51, 1045-1053.	2.4	813
3	Translating Health Status Questionnaires and Evaluating Their Quality. Journal of Clinical Epidemiology, 1998, 51, 913-923.	2.4	761
4	Health-related quality of life as a predictor of mortality and hospitalization: The Dialysis Outcomes and Practice Patterns Study (DOPPS). Kidney International, 2003, 64, 339-349.	2.6	665
5	Health-related quality of life associated with chronic conditions in eight countries: Results from the International Quality of Life Assessment (IQOLA) Project. Quality of Life Research, 2004, 13, 283-298.	1.5	623
6	Nonadherence in hemodialysis: Associations with mortality, hospitalization, and practice patterns in the DOPPS. Kidney International, 2003, 64, 254-262.	2.6	417
7	Tests of Data Quality, Scaling Assumptions, and Reliability of the SF-36 in Eleven Countries. Journal of Clinical Epidemiology, 1998, 51, 1149-1158.	2.4	359
8	Health-related quality of life among dialysis patients on three continents: The Dialysis Outcomes and Practice Patterns Study. Kidney International, 2003, 64, 1903-1910.	2.6	267
9	Development of a Japanese version of the Epworth Sleepiness Scale (JESS) based on Item Response Theory. Sleep Medicine, 2009, 10, 556-565.	0.8	256
10	Physical exercise among participants in the Dialysis Outcomes and Practice Patterns Study (DOPPS): correlates and associated outcomes. Nephrology Dialysis Transplantation, 2010, 25, 3050-3062.	0.4	251
11	Cross-Cultural Comparisons of the Content of SF-36 Translations across 10 Countries. Journal of Clinical Epidemiology, 1998, 51, 925-932.	2.4	250
12	Sleep quality predicts quality of life and mortality risk in haemodialysis patients: Results from the Dialysis Outcomes and Practice Patterns Study (DOPPS). Nephrology Dialysis Transplantation, 2007, 23, 998-1004.	0.4	243
13	Screening for depression in hemodialysis patients: Associations with diagnosis, treatment, and outcomes in the DOPPS. Kidney International, 2004, 66, 2047-2053.	2.6	237
14	Usefulness of five-item and three-item Mental Health Inventories to screen for depressive symptoms in the general population of Japan. Health and Quality of Life Outcomes, 2005, 3, 48.	1.0	209
15	Validation testing of a three-component model of Short Form-36 scores. Journal of Clinical Epidemiology, 2011, 64, 301-308.	2.4	194
16	Psychometric properties of the 25-item National Eye Institute Visual Function Questionnaire (NEI) Tj ETQq0 0 0 0	gBT/Over	·lock 10 Tf 50
17	Systematic Review and Metaanalysis. Chest, 2009, 136, 1576-1585.	0.4	154
18	Effect of a Carbonaceous Oral Adsorbent on the Progression of CKD: A Multicenter, Randomized, Controlled Trial. American Journal of Kidney Diseases, 2009, 54, 459-467.	2.1	142

#	Article	IF	CITATIONS
19	The Associations of Social Support and Other Psychosocial Factors with Mortality and Quality of Life in the Dialysis Outcomes and Practice Patterns Study. Clinical Journal of the American Society of Nephrology: CJASN, 2011, 6, 142-152.	2.2	142
20	Health-related quality of life in the Dialysis Outcomes and Practice Patterns Study (DOPPS). American Journal of Kidney Diseases, 2004, 44, 54-60.	2.1	138
21	Association between hand-grip strength and depressive symptoms: Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). Age and Ageing, 2015, 44, 592-598.	0.7	130
22	Acculturation of attitudes toward end-of-life care. Journal of General Internal Medicine, 2002, 17, 531-539.	1.3	129
23	Travel Time to Dialysis as a Predictor of Health-Related Quality of Life, Adherence, and Mortality: The Dialysis Outcomes and Practice Patterns Study (DOPPS). American Journal of Kidney Diseases, 2008, 51, 641-650.	2.1	123
24	Blood pressure levels and mortality risk among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study. Kidney International, 2012, 82, 570-580.	2.6	120
25	Factors associated with health-related quality of life among hemodialysis patients in the DOPPS. Quality of Life Research, 2007, 16, 545-557.	1.5	119
26	Validation of the Japanese version of the Roland-Morris Disability Questionnaire. Journal of Orthopaedic Science, 2003, 8, 543-548.	0.5	117
27	Relation between sleep quality and quantity, quality of life, and risk of developing diabetes in healthy workers in Japan: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) Study. BMC Public Health, 2007, 7, 129.	1.2	112
28	Association between number of comorbid conditions, depression, and sleep quality using the Pittsburgh Sleep Quality Index: Results from a population-based survey. Sleep Medicine, 2010, 11, 366-371.	0.8	112
29	Functional Dependence and Mortality in the International Dialysis Outcomes and Practice Patterns Study (DOPPS). American Journal of Kidney Diseases, 2016, 67, 283-292.	2.1	110
30	Linguistic Adaptation and Validation of the General Oral Health Assessment Index (GOHAI) in an Elderly Japanese Population. Journal of Public Health Dentistry, 2006, 66, 273-275.	0.5	103
31	Prevalence of lumbar spinal stenosis, using the diagnostic support tool, and correlated factors in Japan: a population-based study. Journal of Orthopaedic Science, 2013, 18, 893-900.	0.5	101
32	Association of mineral metabolism factors with all-cause and cardiovascular mortality in hemodialysis patients: The Japan dialysis outcomes and practice patterns study. Hemodialysis International, 2007, 11, 340-348.	0.4	100
33	Pruritus in hemodialysis patients: Results from the <scp>J</scp> apanese <scp>D</scp> ialysis <scp>O</scp> utcomes and <scp>P</scp> ractice <scp>P</scp> atterns <scp>S</scp> tudy (<scp>JDOPPS</scp>). Hemodialysis International, 2014, 18, 657-667.	0.4	100
34	Sleep-Disordered Breathing in the Usual Lifestyle Setting as Detected with Home Monitoring in a Population of Working Men in Japan. Sleep, 2008, 31, 419-425.	0.6	99
35	Lack of appetite in haemodialysis patients associations with patient characteristics, indicators of nutritional status and outcomes in the international DOPPS. Nephrology Dialysis Transplantation, 2007, 22, 3538-3546.	0.4	93
36	Erythropoiesis-Stimulating Agent Responsiveness and Mortality in Hemodialysis Patients: Results from a Cohort Study From the Dialysis Registry in Japan. American Journal of Kidney Diseases, 2012, 59, 108-116.	2.1	91

#	Article	IF	CITATIONS
37	Testing the Equivalence of Translations of Widely Used Response Choice Labels. Journal of Clinical Epidemiology, 1998, 51, 933-944.	2.4	90
38	Development of a clinical diagnosis support tool to identify patients with lumbar spinal stenosis. European Spine Journal, 2007, 16, 1951-1957.	1.0	89
39	Greater First-Year Survival on Hemodialysis in Facilities in Which Patients Are Provided Earlier and More Frequent Pre-nephrology Visits. Clinical Journal of the American Society of Nephrology: CJASN, 2009, 4, 595-602.	2.2	88
40	Health-related quality of life in the Dialysis Outcomes and Practice Patterns Study (DOPPS). American Journal of Kidney Diseases, 2004, 44, 54-60.	2.1	87
41	Major bleeding events and risk stratification of antithrombotic agents in hemodialysis: results from the DOPPS. Kidney International, 2013, 84, 600-608.	2.6	83
42	Prospective Evaluation of Selection Criteria for Active Surveillance in Japanese Patients with Stage T1cN0M0 Prostate Cancer. Japanese Journal of Clinical Oncology, 2008, 38, 122-128.	0.6	82
43	ACCULTURATION AND END-OF-LIFE DECISION MAKING: COMPARISON OF JAPANESE AND JAPANESE-AMERICAN FOCUS GROUPS. Bioethics, 2007, 21, 251-262.	0.7	78
44	Development of Japanese version of the UCLA Prostate Cancer Index: a pilot validation study. International Journal of Clinical Oncology, 2002, 7, 0306-0311.	1.0	73
45	Multimorbidity patterns in relation to polypharmacy and dosage frequency: a nationwide, cross-sectional study in a Japanese population. Scientific Reports, 2018, 8, 3806.	1.6	73
46	Glycated Hemoglobin and Risk of Death in Diabetic Patients Treated With Hemodialysis: A Meta-analysis. American Journal of Kidney Diseases, 2014, 63, 84-94.	2.1	72
47	Estimating a preference-based index from the Japanese SF-36. Journal of Clinical Epidemiology, 2009, 62, 1323-1331.	2.4	71
48	Japanese version of the Dermatology Life Quality Index: validity and reliability in patients with acne. Health and Quality of Life Outcomes, 2006, 4, 46.	1.0	68
49	A Prospective Study of Passive Smoking and Risk of Diabetes in a Cohort of Workers. Diabetes Care, 2008, 31, 732-734.	4.3	68
50	Associations Between Obstructive Sleep Apnea, Metabolic Syndrome, and Sleep Duration, As Measured With an Actigraph, in an Urban Male Working Population in Japan. Sleep, 2010, 33, 89-95.	0.6	67
51	Health-related quality of life of mothers of children with leukemia in Japan. Quality of Life Research, 2005, 14, 1079-1085.	1.5	66
52	Plasma Thioredoxin, a Novel Oxidative Stress Marker, in Patients with Obstructive Sleep Apnea Before and After Nasal Continuous Positive Airway Pressure. Antioxidants and Redox Signaling, 2008, 10, 715-726.	2.5	66
53	Japanese haemodialysis anaemia management practices and outcomes (1999-2006): results from the DOPPS. Nephrology Dialysis Transplantation, 2008, 23, 3643-3653.	0.4	65
54	Validation of the Japanese version of the quality of recovery score QoR-40. Journal of Anesthesia, 2011, 25, 509-515.	0.7	64

#	Article	IF	CITATIONS
55	Abnormal Mineral Metabolism and Mortality in Hemodialysis Patients With Secondary Hyperparathyroidism: Evidence From Marginal Structural Models Used to Adjust for Time-Dependent Confounding. American Journal of Kidney Diseases, 2014, 63, 979-987.	2.1	64
56	Care burden and depression in caregivers caring for patients with intractable neurological diseases at home in Japan. Journal of the Neurological Sciences, 2009, 276, 148-152.	0.3	63
57	Gender Differences in Effects of Physical Activity on Quality of Life and Resource Utilization. Quality of Life Research, 2006, 15, 537-546.	1.5	59
58	Health-related quality of life of unrelated bone marrow donors in Japan. Blood, 2002, 99, 1995-2001.	0.6	58
59	The influence of locus of control on preferences for information and decision making. Patient Education and Counseling, 2004, 55, 236-240.	1.0	57
60	C-Reactive Protein and Mortality in Hemodialysis Patients: The Dialysis Outcomes and Practice Patterns Study (DOPPS). Nephron Clinical Practice, 2011, 117, c167-c178.	2.3	57
61	Families' and Physicians' Predictions of Dialysis Patients' Preferences Regarding Life-Sustaining Treatments in Japan. American Journal of Kidney Diseases, 2006, 47, 122-130.	2.1	56
62	Vascular Access Care and Treatment Practices Associated with Outcomes of Arteriovenous Fistula: International Comparisons from the Dialysis Outcomes and Practice Patterns Study. Nephron Clinical Practice, 2013, 124, 23-30.	2.3	56
63	Health Related Quality of Life in Japanese Men With Localized Prostate Cancer Treated With Current Multiple Modalities Assessed by a Newly Developed Japanese Version of the Expanded Prostate Cancer Index Composite. Journal of Urology, 2007, 177, 1856-1861.	0.2	53
64	Psychological adjustment has a greater effect on health-related quality of life than on severity of disease in Parkinson's disease. Movement Disorders, 2006, 21, 761-766.	2.2	52
65	Trust in One's Physician: The Role of Ethnic Match, Autonomy, Acculturation, and Religiosity Among Japanese and Japanese Americans. Annals of Family Medicine, 2005, 3, 339-347.	0.9	51
66	Validation of the Burden Index of Caregivers (BIC), a multidimensional short care burden scale from Japan. Health and Quality of Life Outcomes, 2006, 4, 52.	1.0	50
67	Simple four-variable screening tool for identification of patients with sleep-disordered breathing. Sleep, 2009, 32, 939-48.	0.6	50
68	Locomotor dysfunction and risk of cardiovascular disease, quality of life, and medical costs: design of the Locomotive Syndrome and Health Outcome in Aizu Cohort Study (LOHAS) and baseline characteristics of the study population. Journal of Orthopaedic Science, 2012, 17, 261-271.	0.5	49
69	Factors associated with "do not resuscitate―orders and rates of withdrawal from hemodialysis in the international DOPPS. Kidney International, 2005, 68, 1282-1288.	2.6	47
70	PTH-dependence of the effectiveness of cinacalcet in hemodialysis patients with secondary hyperparathyroidism. Scientific Reports, 2016, 6, 19612.	1.6	47
71	Acylated ghrelin level in patients with OSA before and after nasal CPAP treatment. Respirology, 2008, 13, 810-816.	1.3	46
72	Long-term quality of life of donors after living donor liver transplantation. Liver Transplantation, 2012, 18, 1343-1352.	1.3	45

#	Article	IF	Citations
73	Association Between Subjective Sleep Quality and Future Risk of Falls in Older People: Results From LOHAS. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2018, 73, 1205-1211.	1.7	44
74	Predictive value of self-reported patient information for the identification of lumbar spinal stenosis. Family Practice, 2008, 25, 237-244.	0.8	43
75	Association Between Frequency of Pruritic Symptoms and Perceived Psychological Stress. Archives of Dermatology, 2009, 145, 1384-8.	1.7	43
76	Positive Outcomes of High Hemoglobin Target in Patients With Chronic Kidney Disease Not on Dialysis: A Randomized Controlled Study. Therapeutic Apheresis and Dialysis, 2011, 15, 431-440.	0.4	42
77	Changes in anemia management and hemoglobin levels following revision of a bundling policy to incorporate recombinant human erythropoietin. Kidney International, 2011, 79, 340-346.	2.6	41
78	Majority of systematic reviews published in high-impact journals neglected to register the protocols: a meta-epidemiological study. Journal of Clinical Epidemiology, 2017, 84, 54-60.	2.4	41
79	Discrepancy Between Disability and the Severity of Low Back Pain: Demographic, Psychologic, and Employment-Related Factors. Spine, 2006, 31, 931-939.	1.0	39
80	Patient experience of primary care and advance care planning: a multicentre cross-sectional study in Japan. Family Practice, 2017, 34, 206-212.	0.8	39
81	Use of Renin-Angiotensin System Inhibitors Is Associated with Reduction of Fracture Risk in Hemodialysis Patients. PLoS ONE, 2015, 10, e0122691.	1.1	39
82	Mineral Metabolism Management in Hemodialysis Patients with Secondary Hyperparathyroidism in Japan: Baseline Data from the MBD-5D. American Journal of Nephrology, 2011, 33, 427-437.	1.4	38
83	Attitudes of the Japanese public and doctors towards use of archived information and samples without informed consent: Preliminary findings based on focus group interviews. BMC Medical Ethics, 2002, 3, E1.	1.0	37
84	The organization and financing of end-stage renal disease treatment in Japan. International Journal of Health Care Finance and Economics, 2007, 7, 217-231.	1.2	37
85	A Crossâ€sectional Study of Alcohol Drinking and Healthâ€related Quality of Life among Male Workers in Japan. Journal of Occupational Health, 2005, 47, 496-503.	1.0	36
86	Overactive bladder symptom severity is associated with falls in community-dwelling adults: LOHAS study. BMJ Open, 2013, 3, e002413.	0.8	36
87	Reliability, validity, and responsiveness of the Japanese version of the Inflammatory Bowel Disease Questionnaire. Journal of Gastroenterology, 2003, 38, 1138-1143.	2.3	34
88	Consistent Aspirin Use Associated with Improved Arteriovenous Fistula Survival among Incident Hemodialysis Patients in the Dialysis Outcomes and Practice Patterns Study. Clinical Journal of the American Society of Nephrology: CJASN, 2008, 3, 1373-1378.	2.2	34
89	Gender, low Kt/V, and mortality in Japanese hemodialysis patients: Opportunities for improvement through modifiable practices. Hemodialysis International, 2014, 18, 596-606.	0.4	34
90	Focus Group Interviews Examining Attitudes Towards Medical Research Among the Japanese: A Qualitative Study. Bioethics, 2004, 18, 448-470.	0.7	33

#	Article	IF	Citations
91	Health-Related Quality of Life Among Japanese Women With Iron-Deficiency Anemia. Quality of Life Research, 2006, 15, 1559-1563.	1.5	33
92	Health-Related Quality of Life of Predialysis Patients with Chronic Renal Failure. Nephron Clinical Practice, 2006, 105, c1-c8.	2.3	33
93	Associations of Frequency and Duration of Patient-Doctor Contact in Hemodialysis Facilities with Mortality. Journal of the American Society of Nephrology: JASN, 2013, 24, 1493-1502.	3.0	33
94	Revisiting Interdialytic Weight Gain and Mortality Association With Serum Albumin Interactions: The Japanese Dialysis Outcomes and Practice Pattern Study., 2017, 27, 421-429.		32
95	Prevalence of pterygium in a population in Northern Japan: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Acta Ophthalmologica, 2013, 91, e232-6.	0.6	31
96	Influence of urinary urgency and other urinary disturbances on falls in Parkinson's disease. Journal of the Neurological Sciences, 2016, 360, 153-157.	0.3	31
97	Fukushima after the Great East Japan Earthquake: lessons for developing responsive and resilient health systems. Journal of Global Health, 2017, 7, 010501.	1.2	31
98	Prescription Patterns and Mineral Metabolism Abnormalities in the Cinacalcet Era: Results from the MBD-5D Study. Clinical Journal of the American Society of Nephrology: CJASN, 2012, 7, 1473-1480.	2.2	29
99	Prescription of potentially inappropriate medications to elderly hemodialysis patients: prevalence and predictors. Nephrology Dialysis Transplantation, 2015, 30, 498-505.	0.4	29
100	Body Temperature at the Emergency Department as a Predictor of Mortality in Patients With Bacterial Infection. Medicine (United States), 2016, 95, e3628.	0.4	29
101	Both low and high serum ferritin levels predict mortality risk in hemodialysis patients without inflammation. Clinical and Experimental Nephrology, 2017, 21, 685-693.	0.7	29
102	Measuring health-related quality of life in patients with end-stage renal disease: why and how. Nature Clinical Practice Nephrology, 2007, 3, 352-353.	2.0	28
103	Healthâ€related quality of life among communityâ€dwelling patients with intractable neurological diseases and their caregivers in Japan. Psychiatry and Clinical Neurosciences, 2011, 65, 30-38.	1.0	28
104	Decreases in PTH in Japanese Hemodialysis Patients with Secondary Hyperparathyroidism. Clinical Journal of the American Society of Nephrology: CJASN, 2011, 6, 2280-2288.	2.2	28
105	Hypercalcaemia is associated with poor mental health in haemodialysis patients: results from Japan DOPPS. Nephrology Dialysis Transplantation, 2007, 22, 1658-1664.	0.4	27
106	Differences in relationships among sleep apnoea, glucose level, sleep duration and sleepiness between persons with and without type 2 diabetes. Journal of Sleep Research, 2012, 21, 410-418.	1.7	27
107	Identifying Patients with Bacteremia in Community-Hospital Emergency Rooms: A Retrospective Cohort Study. PLoS ONE, 2016, 11, e0148078.	1.1	27
108	Analysis of the evidence-practice gap to facilitate proper medical care for the elderly: investigation, using databases, of utilization measures for National Database of Health Insurance Claims and Specific Health Checkups of Japan (NDB). Environmental Health and Preventive Medicine, 2017, 22, 51.	1.4	27

#	Article	IF	Citations
109	Effect of Patient Experience on Bypassing a Primary Care Gatekeeper: a Multicenter Prospective Cohort Study in Japan. Journal of General Internal Medicine, 2018, 33, 722-728.	1.3	27
110	Factors associated with patient preferences towards deprescribing: a survey of adult patients on prescribed medications. International Journal of Clinical Pharmacy, 2019, 41, 531-537.	1.0	27
111	Gastrointestinal symptoms in a Japanese population: A health diary study. World Journal of Gastroenterology, 2007, 13, 572.	1.4	27
112	Healthâ€related quality of life in Japanese men with localized prostate cancer: Assessment with the SFâ€8. International Journal of Urology, 2008, 15, 524-528.	0.5	26
113	Longitudinal association of fatty pancreas with the incidence of type-2 diabetes in lean individuals: a 6-year computed tomography-based cohort study. Journal of Gastroenterology, 2020, 55, 712-721.	2.3	26
114	Association of coping strategies with mortality and health-related quality of life in hemodialysis patients: The Japan Dialysis Outcomes and Practice Patterns Study. PLoS ONE, 2017, 12, e0180498.	1,1	26
115	Association between ambient air pollution and health-related quality of life in Japan: Ecological study. International Journal of Environmental Health Research, 2005, 15, 383-391.	1.3	25
116	Maintaining high hemoglobin levels improved the left ventricular mass index and quality of life scores in pre-dialysis Japanese chronic kidney disease patients. Clinical and Experimental Nephrology, 2010, 14, 28-35.	0.7	25
117	Effects of Secondary Hyperparathyroidism Treatment on Improvement in Anemia: Results from the MBD-5D Study. PLoS ONE, 2016, 11, e0164865.	1.1	25
118	Use of Prescribed Drugs Among Older People in Japan: Association with Not Having a Regular Physician. Journal of the American Geriatrics Society, 1999, 47, 1425-1429.	1.3	24
119	Dialysis patients[rsquo] preferences regarding cardiopulmonary resuscitation and withdrawal of dialysis in Japan. American Journal of Kidney Diseases, 2001, 37, 1216-1222.	2.1	24
120	Test performance of twoâ€question screening for postpartum depressive symptoms. Pediatrics International, 2009, 51, 48-53.	0.2	24
121	Association between allopurinol and mortality among Japanese hemodialysis patients: results from the DOPPS. International Urology and Nephrology, 2014, 46, 1833-1841.	0.6	24
122	Development of quality indicators for care of chronic kidney disease in the primary care setting using electronic health data: a RAND-modified Delphi method. Clinical and Experimental Nephrology, 2017, 21, 247-256.	0.7	24
123	Chemotherapy in cancer patients undergoing haemodialysis: a nationwide study in Japan. ESMO Open, 2018, 3, e000301.	2.0	24
124	Social Isolation and Patient Experience in Older Adults. Annals of Family Medicine, 2018, 16, 393-398.	0.9	24
125	HEALTH-RELATED QUALITY OF LIFE AMONG RENAL-TRANSPLANT RECIPIENTS IN JAPAN 1. Transplantation, 1999, 68, 1331-1335.	0.5	24
126	Response shift in perception of sleepiness in obstructive sleep apnea-hypopnea syndrome before and after treatment with nasal CPAP. Sleep, 2004, 27, 490-3.	0.6	24

#	Article	IF	CITATIONS
127	Work Performance Assessed by a Newly Developed Japanese Version of the Work Limitation Questionnaire in a General Japanese Adult Population. Journal of Occupational Health, 2014, 56, 124-133.	1.0	23
128	Long-term Trends in Primary Sites of Gastric Adenocarcinoma in Japan and the United States. Journal of Cancer, 2017, 8, 1935-1942.	1.2	23
129	The minimum clinically important difference for the Japanese version of the new Knee Society Score (2011KSS) after total knee arthroplasty. Journal of Orthopaedic Science, 2019, 24, 1053-1057.	0.5	23
130	Dialysis staff encouragement and fluid control adherence in patients on hemodialysis. Nephrology Nursing Journal, 2009, 36, 289-97.	0.1	23
131	Minor Elevation in C-Reactive Protein Levels Predicts Incidence of Erythropoiesis-Stimulating Agent Hyporesponsiveness among Hemodialysis Patients. Nephron, 2015, 131, 123-130.	0.9	22
132	Patients' preferences and factors influencing initial advance care planning discussions' timing: A cross-cultural mixed-methods study. Palliative Medicine, 2020, 34, 906-916.	1.3	22
133	Effect of an Educational Program on Attitudes towards Deceased Organ Donation. Annals of Transplantation, 2015, 20, 269-278.	0.5	22
134	Management of asymptomatic hyperuricaemia in patients with chronic kidney disease by Japanese nephrologists: A questionnaire survey. Nephrology, 2011, 16, 518-521.	0.7	21
135	Mineral and Bone Disorders Outcomes Study for Japanese Chronic Kidney Disease Stage 5D Patients: Rationale and Study Design. Therapeutic Apheresis and Dialysis, 2011, 15, 169-175.	0.4	21
136	Association of blood pressure with allâ€cause mortality and stroke in <scp>J</scp> apanese hemodialysis patients: The <scp>J</scp> apan dialysis outcomes and practice pattern study. Hemodialysis International, 2014, 18, 607-615.	0.4	21
137	Development and Validation of a Symptom Scale to Evaluate Postoperative Patients with Esophagogastric Cancer. Journal of the American College of Surgeons, 2014, 219, 895-903.	0.2	21
138	Postdialysis Hypokalemia and All-Cause Mortality in Patients Undergoing Maintenance Hemodialysis. Clinical Journal of the American Society of Nephrology: CJASN, 2019, 14, 873-881.	2.2	21
139	Higher Internality of Health Locus of Control Is Associated With the Use of Complementary and Alternative Medicine Providers Among Patients Seeking Care for Acute Low-back Pain. Clinical Journal of Pain, 2008, 24, 725-730.	0.8	20
140	Disappearance of Association in Diabetic Patients on Hemodialysis between Anemia and Mortality Risk: The Japan Dialysis Outcomes and Practice Pattern Study. Nephron Clinical Practice, 2012, 120, c91-c100.	2.3	20
141	Relationship between health-related quality of life and clustering of metabolic syndrome diagnostic components. Quality of Life Research, 2012, 21, 1165-1170.	1.5	20
142	Mentoring the next generation of physician-scientists in Japan: a cross-sectional survey of mentees in six academic medical centers. BMC Medical Education, 2015, 15, 54.	1.0	20
143	Low Testosterone Levels and Reduced Kidney Function in Japanese Adult Men: The Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Journal of the American Medical Directors Association, 2016, 17, 371.e1-371.e6.	1.2	20
144	Quality of life after single-incision laparoscopic cholecystectomy: A randomized, clinical trial. Surgery, 2019, 165, 353-359.	1.0	20

#	Article	IF	CITATIONS
145	Effect of coaching on psychological adjustment in patients with spinocerebellar degeneration: a pilot study. Clinical Rehabilitation, 2007, 21, 987-996.	1.0	19
146	Relationships between working status and health or health are utilization among Japanese elderly. Geriatrics and Gerontology International, 2008, 8, 32-40.	0.7	19
147	Cost-effectiveness of administering oral adsorbent AST-120 to patients with diabetes and advance-stage chronic kidney disease. Diabetes Research and Clinical Practice, 2010, 90, 154-159.	1.1	19
148	Gender Difference in Association Between Low Back Pain and Metabolic Syndrome. Spine, 2012, 37, 1130-1137.	1.0	19
149	Chronic hyperglycemia increases the risk of lateral epicondylitis: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study (LOHAS). SpringerPlus, 2015, 4, 407.	1.2	19
150	Unique hemoglobin A1c level distribution and its relationship with mortality in diabetic hemodialysis patients. Kidney International, 2017, 92, 497-503.	2.6	19
151	Associations of employment status and educational levels with mortality and hospitalization in the dialysis outcomes and practice patterns study in Japan. PLoS ONE, 2017, 12, e0170731.	1.1	19
152	Lifestyle and work predictors of fatigue in Japanese manufacturing workers. Occupational Medicine, 2007, 57, 262-269.	0.8	18
153	Depressive Symptoms Predict the Subsequent Risk of Bodily Pain in Dialysis Patients: Japan Dialysis Outcomes and Practice Patterns Study. Pain Medicine, 2009, 10, 883-889.	0.9	18
154	Cost-utility analysis of cataract surgery in Japan: a probabilistic Markov modeling study. Japanese Journal of Ophthalmology, 2013, 57, 391-401.	0.9	18
155	One-Year Period Prevalence of Oral Aphthous Ulcers and Oral Health-Related Quality of Life in Patients with Behçet's Disease. Genetics Research International, 2014, 2014, 1-8.	2.0	18
156	Dietary Patterns and Clinical Outcomes in Hemodialysis Patients in Japan: A Cohort Study. PLoS ONE, 2015, 10, e0116677.	1.1	18
157	The Role of Digital Rectal Examination for Diagnosis of Acute Appendicitis: A Systematic Review and Meta-Analysis. PLoS ONE, 2015, 10, e0136996.	1.1	18
158	Contribution of dysregulated serum magnesium to mortality in hemodialysis patients with secondary hyperparathyroidism: a 3-year cohort study. CKJ: Clinical Kidney Journal, 2015, 8, 744-752.	1.4	18
159	Prognostic utility of serum CRP levels in combination with CURB-65 in patients with clinically suspected sepsis: a decision curve analysis. BMJ Open, 2015, 5, e007049-e007049.	0.8	18
160	Independent association between prediabetes and future pancreatic fat accumulation: a 5-year Japanese cohort study. Journal of Gastroenterology, 2018, 53, 873-882.	2.3	18
161	Development and validation of a concise scale for assessing patient experience of primary care for adults in Japan. Family Practice, 2020, 37, 137-142.	0.8	18
162	Vitality Measured as Self-reported Energy Level and Clinical Outcomes in Hemodialysis Patients: The Japanese Dialysis Outcomes and Practice Pattern Study (J-DOPPS). American Journal of Kidney Diseases, 2019, 73, 486-495.	2.1	18

#	Article	IF	Citations
163	Computed tomography during initial management and mortality among hemodynamically unstable blunt trauma patients: a nationwide retrospective cohort study. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2017, 25, 74.	1.1	17
164	Social participation and the combination of future needs for long-term care and mortality among older Japanese people: a prospective cohort study from the Aichi Gerontological Evaluation Study (AGES). BMJ Open, 2019, 9, e030500.	0.8	17
165	Longitudinal Association Between Subjective Fatigue and Future Falls in Community-Dwelling Older Adults: The Locomotive Syndrome and Health Outcomes in the Aizu Cohort Study (LOHAS). Journal of Aging and Health, 2019, 31, 67-84.	0.9	17
166	Living close to automobile traffic and quality of life in Japan: A population-based survey. International Journal of Environmental Health Research, 2005, 15, 1-9.	1.3	16
167	Keeping vulnerable elderly patients free from pressure ulcer is associated with high caregiver burden in informal caregivers. Journal of Evaluation in Clinical Practice, 2010, 16, 585-589.	0.9	16
168	Assessment of vision-related quality of life among patients with cataracts and the outcomes of cataract surgery using a newly developed visual function questionnaire: the VFQ-J11. Japanese Journal of Ophthalmology, 2014, 58, 415-422.	0.9	16
169	Low Testosterone Levels, Depressive Symptoms, and Falls in Older Men: A Cross-Sectional Study. Journal of the American Medical Directors Association, 2014, 15, 30-35.	1.2	16
170	Relationship Between Kyphotic Posture and Falls in Community-Dwelling Men and Women. Spine, 2016, 41, 1232-1238.	1.0	16
171	Association between pancreatic fat and incidence of metabolic syndrome: a 5â€year Japanese cohort study. Journal of Gastroenterology and Hepatology (Australia), 2018, 33, 2048-2054.	1.4	16
172	Wholeâ∈Body Computed Tomography During Initial Management and Mortality Among Adult Severe Blunt Trauma Patients: A Nationwide Cohort Study. World Journal of Surgery, 2018, 42, 3939-3946.	0.8	16
173	Quality of Life Improvements in Dialysis Patients Receiving Darbepoetin Alfa. Therapeutic Apheresis and Dialysis, 2008, 12, 72-77.	0.4	15
174	Impact of cinacalcet introduction on MBD management: the MBD-5D study in Japan. Kidney International Supplements, 2013, 3, 436-441.	4.6	15
175	A new prognostic index for overall survival in malignant pleural mesothelioma: the rPHS (regimen,) Tj ETQq1 1 0.7	784314 rg 0.6	BT_lOverlock
176	The cardiothoracic ratio and all-cause and cardiovascular disease mortality in patients undergoing maintenance hemodialysis: results of the MBD-5D study. Clinical and Experimental Nephrology, 2017, 21, 797-806.	0.7	15
177	Salt Intake and All-Cause Mortality in Hemodialysis Patients. American Journal of Nephrology, 2018, 48, 87-95.	1.4	15
178	Hospitals' extracorporeal cardiopulmonary resuscitation capabilities and outcomes in out-of-hospital cardiac arrest: A population-based study. Resuscitation, 2019, 136, 85-92.	1.3	15
179	Level of Low Back Pain–Related Disability Is Associated with Risk of Subsequent Falls in an Older Population: Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). Pain Medicine, 2019, 20, 2377-2384.	0.9	15
180	Early rehabilitation in older patients hospitalized with acute decompensated heart failure: A retrospective cohort study. American Heart Journal, 2020, 230, 44-53.	1.2	15

#	Article	IF	CITATIONS
181	Effects of intravenous ascorbic acid on erythropoiesis and quality of life in unselected hemodialysis patients. Journal of Nephrology, 2004, 17, 537-43.	0.9	15
182	Telemedicine in Neurosurgery Using International Digital Telephone Services Between Japan and Malaysia â€"Technical Noteâ€". Neurologia Medico-Chirurgica, 1999, 39, 773-778.	1.0	14
183	Relationships Between Diurnal Blood Pressure Variation, Physical Activity, and Healthâ€Related QOL. Clinical and Experimental Hypertension, 2004, 26, 145-155.	0.5	14
184	Associations between exposure to ambient photochemical oxidants and the vitality or mental health domain of the health related quality of life. Journal of Epidemiology and Community Health, 2006, 60, 173-179.	2.0	14
185	The relation between self-reported body weight and health-related quality of life: a cross-sectional study in Japan. Journal of Public Health, 2011, 33, 518-526.	1.0	14
186	Relationships of Decreased Lung Function with Metabolic Syndrome and Obstructive Sleep Apnea in Japanese Males. Internal Medicine, 2012, 51, 2291-2297.	0.3	14
187	Association between social networks and discussions regarding advance care planning among Japanese older adults. PLoS ONE, 2019, 14, e0213894.	1.1	14
188	Health-related quality of life among dialysis patients in Seattle and Aichi. American Journal of Kidney Diseases, 2001, 37, 987-996.	2.1	13
189	Retinal vessel diameters in a Japanese population: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Acta Ophthalmologica, 2016, 94, e432-41.	0.6	13
190	Influence of family dynamics on burden among family caregivers in aging Japan. Family Practice, 2016, 33, 466-470.	0.8	13
191	Effects of self-monitoring of daily salt intake estimated by a simple electrical device for salt reduction: a cluster randomized trial. Hypertension Research, 2018, 41, 524-530.	1.5	13
192	Gait speed and overactive bladder in the healthy communityâ€dwelling super elderly: The Sukagawa Study. Neurourology and Urodynamics, 2019, 38, 2324-2332.	0.8	13
193	Association of low alanine aminotransferase with loss of independence or death: A 5â€year populationâ€based cohort study. Journal of Gastroenterology and Hepatology (Australia), 2019, 34, 1793-1799.	1.4	13
194	Association between contrast extravasation on computed tomography scans and pseudoaneurysm formation in pediatric blunt splenic and hepatic injury: A multi-institutional observational study. Journal of Pediatric Surgery, 2020, 55, 681-687.	0.8	13
195	Comparison of Primary Care Experience in Hospital-Based Practices and Community-Based Office Practices in Japan. Annals of Family Medicine, 2020, 18, 24-29.	0.9	13
196	Association between religious beliefs and discussions regarding advance care planning: A nationwide survey. Palliative Medicine, 2021, 35, 1856-1864.	1.3	13
197	Violent Patient Behavior Is Associated with Bodily Pain and a High Burden on Informal Caregivers. Journal of General Internal Medicine, 2009, 24, 1085-1088.	1.3	12
198	Effects of the presence of hypertension on the relationship between obstructive sleep apnoea and sleepiness. Journal of Sleep Research, 2011, 20, 538-543.	1.7	12

#	Article	IF	CITATIONS
199	Development of a Short Version of the Visual Function Questionnaire Using Item-Response Theory. PLoS ONE, 2013, 8, e73084.	1.1	12
200	Association of Serum Bicarbonate with Bone Fractures in Hemodialysis Patients: The Mineral and Bone Disorder Outcomes Study for Japanese CKD Stage 5D Patients (MBD-5D). Nephron Clinical Practice, 2014, 128, 79-87.	2.3	12
201	Lumbar Spinal Stenosis–Specific Symptom Scale. Spine, 2014, 39, E1388-E1393.	1.0	12
202	Development and Validation of a Disease-Specific Instrument to Measure Diet-Targeted Quality of Life for Postoperative Patients with Esophagogastric Cancer. Annals of Surgical Oncology, 2015, 22, 848-854.	0.7	12
203	Effects of structured education program on organ donor designation of nursing students and their families: A randomized controlled trial. Clinical Transplantation, 2016, 30, 1513-1519.	0.8	12
204	Effects of Higher Quality of Care on Initiation of Long-term Dialysis in Patients With CKD and Diabetes. American Journal of Kidney Diseases, 2017, 70, 666-674.	2.1	12
205	Impact of nutritional index on the association between phosphorus concentrations and mortality in haemodialysis patients: a cohort study from dialysis outcomes and practice pattern study in Japan. BMJ Open, 2017, 7, e016682.	0.8	12
206	Association between vision-specific quality of life and falls in community-dwelling older adults: LOHAS. PLoS ONE, 2018, 13, e0195806.	1.1	12
207	Association between health-related hope and adherence to prescribed treatment in CKD patients: multicenter cross-sectional study. BMC Nephrology, 2020, 21, 453.	0.8	12
208	A scale for measuring health-related hope: its development and psychometric testing. Annals of Clinical Epidemiology, 2019, 1, 102-119.	0.3	12
209	Sex differences in the change in health-related quality of life associated with low back pain. Quality of Life Research, 2012, 21, 1705-1711.	1.5	11
210	Impact of trajectories of abdominal aortic calcification over 2 years on subsequent mortality: a 10-year longitudinal study. Nephrology Dialysis Transplantation, 2018, 33, 676-683.	0.4	11
211	Inverse Association Between Fatty Liver at Baseline Ultrasonography and Remission of Type 2 Diabetes Over a 2-Year Follow-up Period. Clinical Gastroenterology and Hepatology, 2021, 19, 556-564.e5.	2.4	11
212	Type I Angiotensin II Receptor Blockade Reduces Uremia-Induced Deterioration of Bone Material Properties. Journal of Bone and Mineral Research, 2020, 36, 67-79.	3.1	11
213	Experience as an Informal Caregiver and Discussions Regarding Advance Care Planning in Japan. Journal of Pain and Symptom Management, 2021, 61, 63-70.	0.6	11
214	Association of glucocorticoid doses and emotional health in lupus low disease activity state (LLDAS): a cross-sectional study. Arthritis Research and Therapy, 2021, 23, 79.	1.6	11
215	The Development and Use of Quality-of-Life Measures to Evaluate Health Outcomes in Japan. Pharmacoeconomics, 2002, 20, 17-23.	1.7	10
216	Development and Validation of a Symptom Scale for Lumbar Spinal Stenosis. Spine, 2012, 37, 232-239.	1.0	10

#	Article	IF	CITATIONS
217	Patient-reported disability in the general Japanese population was associated with medical care visits for low back pain, regardless of pain intensity. Journal of Orthopaedic Science, 2015, 20, 742-749.	0.5	10
218	Frequent monitoring of mineral metabolism in hemodialysis patients with secondary hyperparathyroidism: associations with achievement of treatment goals and with adjustments in therapy. Nephrology Dialysis Transplantation, 2016, 32, gfw020.	0.4	10
219	Development of a new risk model for predicting cardiovascular events among hemodialysis patients: Population-based hemodialysis patients from the Japan Dialysis Outcome and Practice Patterns Study (J-DOPPS). PLoS ONE, 2017, 12, e0173468.	1.1	10
220	Salt intake and body weight correlate with higher blood pressure in the very elderly population: The Sukagawa study. Journal of Clinical Hypertension, 2019, 21, 942-949.	1.0	10
221	Validation of the Japanese version of the Quality of Life-Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA). Growth Hormone and IGF Research, 2006, 16, 340-347.	0.5	9
222	Musculoskeletal pain in Japan: prospective health diary study. Rheumatology International, 2007, 28, 7-14.	1.5	9
223	Opinions of Japanese rheumatology physicians regarding clinical practice guidelines. International Journal for Quality in Health Care, 2010, 22, 78-85.	0.9	9
224	Relationship of type of work with health-related quality of life. Quality of Life Research, 2015, 24, 2927-2932.	1.5	9
225	Development and validation of a prediction model for loss of physical function in elderly hemodialysis patients. Nephrology Dialysis Transplantation, 2018, 33, 1452-1458.	0.4	9
226	Association Between the Discrepancy in Self-Reported and Performance-Based Physical Functioning Levels and Risk of Future Falls Among Community-Dwelling Older Adults: The Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). Journal of the American Medical Directors Association, 2019, 20, 195-200.e1.	1,2	9
227	How Psychosocial Factors Contribute to Japanese Older Adults' Initiation of Advance Care Planning Discussions: A Qualitative Study. Journal of Applied Gerontology, 2021, 40, 1180-1188.	1.0	9
228	Effects of low dialysate calcium concentration on health-related quality of life in hemodialysis patients. Clinical and Experimental Nephrology, 2005, 9, 153-157.	0.7	8
229	Darbepoetin alfa (KRN321) is safe and effective when administered subcutaneously once every 2 or 4Âweeks to patients on peritoneal dialysis in Japan. Clinical and Experimental Nephrology, 2011, 15, 884-892.	0.7	8
230	Effects of cyclophosphamide on the prognosis of Japanese patients with renal vasculitis associated with anti-neutrophil cytoplasmic antibody-positive microscopic polyangiitis. Clinical and Experimental Nephrology, 2016, 20, 712-719.	0.7	8
231	Association between parental history of Helicobacter pyloritreatment failure and treatment failure in the offspring. Journal of Gastroenterology and Hepatology (Australia), 2019, 34, 2112-2117.	1.4	8
232	Interdialytic Weight Gain Effects on Hemoglobin Concentration and Cardiovascular Events. Kidney International Reports, 2020, 5, 1670-1678.	0.4	8
233	Healthcare Providers' Perceptions on the Timing of Initial Advance Care Planning Discussions in Japan: a Mixed-Methods Study. Journal of General Internal Medicine, 2021, 36, 2935-2942.	1.3	8
234	Effect of multimorbidity patterns on the decline in health-related quality of life: a nationwide prospective cohort study in Japan. BMJ Open, 2021, 11, e047812.	0.8	8

#	Article	IF	CITATIONS
235	Understanding Measurements of Vitality in Patients with Chronic Kidney Disease: Connecting a Quality-of-Life Scale to Daily Activities. PLoS ONE, 2012, 7, e40455.	1.1	8
236	Development and Validation of a Clinical Prediction Rule for Bacteremia among Maintenance Hemodialysis Patients in Outpatient Settings. PLoS ONE, 2017, 12, e0169975.	1.1	8
237	Results of the international DOPPS hemodialysis study in Japan. Nihon Toseki Igakkai Zasshi, 2004, 37, 1865-1873.	0.2	7
238	Development and Testing of the Japanese Version of the Migraine-Specific Quality of Life Instrument. Quality of Life Research, 2004, 13, 1489-1493.	1.5	7
239	Study design and methods for a clinical trial of an oral carbonaceous adsorbent used to prevent the progression of chronic kidney disease (CAP-KD). Clinical and Experimental Nephrology, 2005, 9, 219-227.	0.7	7
240	A study of the burden of caring for patients with amyotrophic lateral sclerosis (MND) in Japan. British Journal of Neuroscience Nursing, 2006, 2, 38-43.	0.1	7
241	Impact of Early Referral to Nephrologist on Mental Health among Hemodialysis Patients: A Dialysis Outcomes and Practice Patterns Study (DOPPS). Nephron Clinical Practice, 2009, 113, c191-c197.	2.3	7
242	Development and Validation of a Quality of Life Scale Specific for Lumbar Spinal Stenosis. Spine, 2011, 36, E1407-E1414.	1.0	7
243	Increased Mortality Rate after Hospitalization Among Chronic Hemodialysis Patients: A Prospective Cohort Study. Nephron, 2018, 140, 194-202.	0.9	7
244	Application of a two-question screening instrument to detect depressive symptoms in patients with vitiligo: AÂpilot study. Journal of the American Academy of Dermatology, 2011, 64, e69-e70.	0.6	6
245	Attributes of primary care in relation to polypharmacy: a multicenter cross-sectional study in Japan. International Journal for Quality in Health Care, 2017, 29, 378-383.	0.9	6
246	Interactive Effectiveness of Angiotensin-Converting Enzyme Inhibitors and Angiotensin Receptor Blockers or Their Combination on Survival of Hemodialysis Patients. American Journal of Nephrology, 2017, 46, 439-447.	1.4	6
247	Association between statistical significance and time to publication among systematic reviews: a study protocol for a meta-epidemiological investigation. BMJ Open, 2017, 7, e018856.	0.8	6
248	The development of quality indicators for systemic lupus erythematosus using electronic health data: A modified RAND appropriateness method. Modern Rheumatology, 2020, 30, 525-531.	0.9	6
249	Physical multimorbidity patterns and depressive symptoms: a nationwide cross-sectional study in Japan. Family Medicine and Community Health, 2020, 8, e000234.	0.6	6
250	Low Predialysis Plasma Calculated Osmolality Is Associated with Higher All-Cause Mortality: The Japanese Dialysis Outcomes and Practice Patterns Study (J-DOPPS). Nephron, 2020, 144, 138-146.	0.9	6
251	Relationship between serum calcium or phosphate levels and mortality stratified by parathyroid hormone level: an analysis from the MBD-5D study. Clinical and Experimental Nephrology, 2020, 24, 630-637.	0.7	6
252	Knowledge Does Not Correlate with Behavior toward Deceased Organ Donation: A Cross-Sectional Study in Japan. Annals of Transplantation, 2020, 25, e918936.	0.5	6

#	Article	IF	Citations
253	Measurement of Information Seeking Preference of Japanese in Clinical Settings. The Japanese Journal of Health Psychology, 1997, 10, 1-10.	0.1	5
254	Comparison of dental practice income and expenses according to treatment types in the Japanese insurance system. Japanese Dental Science Review, 2010, 46, 4-10.	2.0	5
255	Can pediatricians accurately identify maternal depression at wellâ€child visits?. Pediatrics International, 2010, 52, 284-289.	0.2	5
256	Post-discharge depressive symptoms can predict quality of life in AMI survivors: A prospective cohort study in Japan. International Journal of Cardiology, 2011, 146, 379-384.	0.8	5
257	Dialysis Amyloid Deposition in the Aortic Valve and Its Association with Aortic Stenosis. Blood Purification, 2015, 40, 146-154.	0.9	5
258	Diagnostic utility of appetite loss in addition to existing prediction models for community-acquired pneumonia in the elderly: a prospective diagnostic study in acute care hospitals in Japan. BMJ Open, 2017, 7, e019155.	0.8	5
259	Body Mass Index Change and Hospitalization Risk in Elderly Hemodialysis Patients: Results from Japanese Dialysis Outcomes and Practice Patterns Study. American Journal of Nephrology, 2018, 47, 48-56.	1.4	5
260	Association Between Accumulation of Advanced Glycation End-Products and Hearing Impairment in Community-Dwelling Older People: A Cross-Sectional Sukagawa Study. Journal of the American Medical Directors Association, 2018, 19, 235-239.e1.	1.2	5
261	Usefulness of computed tomography in the diagnosis of acute pyelonephritis in older patients suspected of infection with unknown focus. Acta Radiologica, 2022, 63, 268-277.	0.5	5
262	Cohort profile: A super-elderly population-based cohort in Japan (the Sukagawa Study). International Journal of Epidemiology, 2021, 50, 727-727h.	0.9	5
263	Intraoperative end-tidal carbon dioxide and postoperative mortality in major abdominal surgery: a historical cohort study. Canadian Journal of Anaesthesia, 2021, 68, 1601-1610.	0.7	5
264	Japanese Attitudes toward Advance Care Planning for End-of-Life Decisions. General Medicine, 2001, 2, 3-10.	0.1	5
265	Low Back Pain and Associated Occupational Factors among Nursery School Teachers: A Multicenter, Cross-sectional Study. Annals of Clinical Epidemiology, 2020, 2, 113-120.	0.3	5
266	Low quality of life in men with chronic prostatitis-like symptoms. Prostate Cancer and Prostatic Diseases, 2022, 25, 785-790.	2.0	5
267	Reliability, validity, and responsiveness of the Japanese version of the Neck Pain and Disability Scale. Journal of Orthopaedic Science, 2011, 16, 339-346.	0.5	4
268	Interpersonal Psychosocial Factors Associated With Underreported Dietary Energy Intake in Hemodialysis Patients., 2013, 23, 37-44.		4
269	Clinical Usefulness of the Two-Question Assessment Tool for Depressive Symptoms in Japanese Patients with Chronic Obstructive Pulmonary Disease. Lung, 2013, 191, 101-107.	1.4	4
270	Heart Rate, Responsiveness to Intravenous Immunoglobulin, and Coronary Artery Aneurysms in Kawasaki Disease. Journal of Pediatrics, 2018, 200, 160-165.e5.	0.9	4

#	Article	IF	Citations
271	Public subsidies and the recommendation of child vaccines among primary care physicians: a nationwide cross-sectional study in Japan. BMJ Open, 2018, 8, e020923.	0.8	4
272	Statistical significance did not affect time to publication in non-Cochrane systematic reviews: a metaepidemiological study. Journal of Clinical Epidemiology, 2019, 115, 25-34.	2.4	4
273	Ulcerative colitis outcomes research in Japan: protocol for an observational prospective cohort study of YOURS (YOu and Ulcerative colitis: Registry and Social network). BMJ Open, 2019, 9, e030134.	0.8	4
274	Changes in 3-month mineral and bone disorder patterns were associated with all-cause mortality in prevalent hemodialysis patients with secondary hyperparathyroidism. BMC Nephrology, 2020, 21, 432.	0.8	4
275	Quality of Care in Chronic Kidney Disease and Incidence of End-stage Renal Disease in Older Patients. Medical Care, 2020, 58, 625-631.	1.1	4
276	Urgency urinary incontinence, loss of independence, and increased mortality in older adults: A cohort study. PLoS ONE, 2021, 16, e0245724.	1.1	4
277	Poor oral function is associated with loss of independence or death in functionally independent older adults. PLoS ONE, 2021, 16, e0253559.	1.1	4
278	Usefulness of a mobile phone application for respiratory rate measurement in adult patients. Japan Journal of Nursing Science, 2022, 19, e12481.	0.5	4
279	Contributions of Clinical Epidemiologists and Medical Librarians to Developing Evidence-Based Clinical Practice Guidelines in Japan: A Case of the Treatment of Rheumatoid Arthritis. General Medicine, 2003, 4, 21-28.	0.1	3
280	Prospective Health Diary Study for New Onset Chest Symptoms in the Japanese General Population. Internal Medicine, 2008, 47, 25-31.	0.3	3
281	Evaluation of Newspaper Articles for Coverage of Public Reporting Data: A Case Study of Unadjusted Cancer Survival Data. Japanese Journal of Clinical Oncology, 2013, 43, 95-100.	0.6	3
282	New oral anticoagulants versus warfarin for preventing stroke and systemic embolic events among atrial fibrillation patients with chronic kidney disease. The Cochrane Library, 2014, , .	1.5	3
283	Cardiovascular and Stroke Risk in Japanese Hemodialysis Patients With Atrial Fibrillation. Therapeutic Apheresis and Dialysis, 2016, 20, 608-614.	0.4	3
284	External validation of prognostic indices for overall survival of malignant pleural mesothelioma. Lung Cancer, 2017, 113, 88-92.	0.9	3
285	Development of a profile scoring system for assessing the psychosocial situation of patients with chronic musculoskeletal pain. Journal of Pain Research, 2017, Volume 10, 1853-1859.	0.8	3
286	Association between keeping home records of catheter exit-site and incidence of peritoneal dialysis-related infections. International Urology and Nephrology, 2018, 50, 763-769.	0.6	3
287	Haemoglobin concentration and survival of haemodialysis patients before and after experiencing cardiovascular disease: a cohort study from Japanese dialysis outcomes and practice pattern study (J-DOPPS). BMJ Open, 2019, 9, e031476.	0.8	3
288	Association between work schedules and motivation for lifestyle change in workers with overweight or obesity: a cross-sectional study in Japan. BMJ Open, 2020, 10, e033000.	0.8	3

#	Article	IF	CITATIONS
289	Quality of Life and Disability-free Survival in the Elderly: The Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Journal of Aging and Health, 2021, 33, 197-204.	0.9	3
290	Significance of Overactive Bladder as a Predictor of Falls in Community Dwelling Older Adults: 1-Year Followup of the Sukagawa Study. Journal of Urology, 2021, 205, 219-225.	0.2	3
291	Added value of clinical prediction rules for bacteremia in hemodialysis patients: An external validation study. PLoS ONE, 2021, 16, e0247624.	1.1	3
292	Is "pain before vomiting―useful?: Diagnostic performance of the classic patient history item in acute appendicitis. American Journal of Emergency Medicine, 2021, 41, 84-89.	0.7	3
293	Diagnostic performance of food consumption for bacteraemia in patients admitted with suspected infection: a prospective cohort study. BMJ Open, 2021, 11, e044270.	0.8	3
294	Changes in Drug Utilization After Publication of Clinical Trials and Drug-Related Scandals in Japan: An Interrupted Time Series Analysis, 2005–2017. Journal of Epidemiology, 2021, 31, 410-416.	1.1	3
295	Impact of Primary Tumor Resection on Mortality in Patients with Stage IV Colorectal Cancer with Unresectable Metastases: A Multicenter Retrospective Cohort Study. World Journal of Surgery, 2021, 45, 3230-3239.	0.8	3
296	Shorter dialysis session length was not associated with lower mental health and physical functioning in elderly hemodialysis patients: Results from the Japan Dialysis Outcome and Practice Patterns Study (J-DOPPS). PLoS ONE, 2017, 12, e0184019.	1.1	3
297	Results of the survey on practice patterns including MBD management at dialysis facilities. Nihon Toseki Igakkai Zasshi, 2011, 44, 557-566.	0.2	3
298	Association of Lumbar Spinal Stenosis With Severe Disability and Mortality Among Community-Dwelling Older Adults. Spine, 2020, Publish Ahead of Print, E784-E790.	1.0	3
299	Association of kyphotic posture with loss of independence and mortality in a community-based prospective cohort study: the Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). BMJ Open, 2022, 12, e052421.	0.8	3
300	Impact of lumbar spinal stenosis on metabolic syndrome incidence in community-dwelling adults in Aizu cohort study (LOHAS). Scientific Reports, 2022, 12, .	1.6	3
301	Relationship between Having a Home Doctor and Outpatient Utilization Journal of Epidemiology, 2001, 11, 160-169.	1.1	2
302	Prognostic Factors for a Composite End-Point of Renal Outcomes in Patients With Chronic Kidney Disease. Therapeutic Apheresis and Dialysis, 2006, 10, 72-77.	0.4	2
303	A simple four-grade score for aortic arch calcification by posteroanterior chest X-ray is associated with cardiovascular disease in haemodialysis patients. Nephrology Dialysis Transplantation, 2011, 26, 1747-1748.	0.4	2
304	Opinion of Japanese rheumatology physicians on methods of assessing the quality of rheumatoid arthritis care. Journal of Evaluation in Clinical Practice, 2012, 18, 290-295.	0.9	2
305	Influence of Staff Encouragement on Perceived Burden of Dietary Restriction Among Patients Living Alone. Therapeutic Apheresis and Dialysis, 2016, 20, 623-631.	0.4	2
306	Preferences of young physicians at community hospitals regarding academic research training through graduate school: a cross-sectional research. BMC Research Notes, 2016, 9, 227.	0.6	2

#	Article	IF	Citations
307	Association between daily salt intake of 3â€yearâ€old children and that of their mothers: A crossâ€sectional study. Journal of Clinical Hypertension, 2018, 20, 730-735.	1.0	2
308	Dialysis physicians' referral behaviors for hemodialysis patients suspected of having cancer: A vignette-based questionnaire study. PLoS ONE, 2018, 13, e0202322.	1.1	2
309	Association between post-dialysis hemoglobin level and the survival of vascular access. Journal of Vascular Access, 2018, 19, 69-75.	0.5	2
310	Development and validation of a prediction model for functional decline in older medical inpatients. Archives of Gerontology and Geriatrics, 2018, 77, 184-188.	1.4	2
311	Determinants of incentive preferences for health behavior change in Japan. Health Promotion International, 2019, 34, 760-769.	0.9	2
312	Vehicle configurations associated with anatomical-specific severe injuries resulting from traffic collisions. PLoS ONE, 2019, 14, e0223388.	1.1	2
313	Effectiveness of cinacalcet treatment for secondary hyperparathyroidism on hospitalization: Results from the MBD-5D study. PLoS ONE, 2019, 14, e0216399.	1.1	2
314	Health-related quality of life in Japanese patients with bladder cancer measured by a newly developed Japanese version of the Bladder Cancer Index. International Journal of Clinical Oncology, 2020, 25, 2090-2098.	1.0	2
315	Associations of types of primary care facilities with adult vaccination and cancer screening in Japan. International Journal for Quality in Health Care, 2020, 32, 373-378.	0.9	2
316	The Association between the Establishment of a General Internal Medicine Department and an Increased Number of Blood Cultures in Other Departments: An Interrupted Time Series Analysis. Internal Medicine, 2021, 60, 3729-3735.	0.3	2
317	Impact of Vonoprazan Triple-Drug Blister Packs on H.Âpylori Eradication Rates in Japan: Interrupted Time Series Analysis. Advances in Therapy, 2021, 38, 3937-3947.	1.3	2
318	Association between facility-level adherence to phosphorus management guidelines and mortality in haemodialysis patients: a prospective cohort study. BMJ Open, 2021, 11, e051002.	0.8	2
319	Decline in oral antimicrobial prescription in the outpatient setting after nationwide implementation of financial incentives and provider education: An interrupted time-series analysis. Infection Control and Hospital Epidemiology, 2023, 44, 253-259.	1.0	2
320	Association between high voltage overhead transmission lines and mental health: A cross-sectional study. Bioelectromagnetics, 2006, 27, 473-478.	0.9	1
321	Headache, Mental Health, and Use of Medical Resources: Health Diary Study in Japan. Journal of Health Science, 2008, 54, 30-36.	0.9	1
322	Diabetes in Asia. Lancet, The, 2010, 375, 981-982.	6.3	1
323	Status of Provision of Drug Information to Hypertensive Patients at Community-based Pharmacies. General Medicine, 2011, 12, 75-82.	0.1	1
324	Developing a scale to measure family dynamics related to long-term care, and testing that scale in a multicenter cross-sectional study. BMC Family Practice, 2014, 15, 134.	2.9	1

#	Article	IF	CITATIONS
325	Effect of ABO blood type on the outcomes of patients with metastatic renal cell carcinoma treated with first-line tyrosine kinase inhibitors. Urologic Oncology: Seminars and Original Investigations, 2017, 35, 540.e7-540.e12.	0.8	1
326	Childcare Environment and Japanese Children Who Are Overweight in Early Childhood. Childhood Obesity, 2018, 14, 197-206.	0.8	1
327	Realâ€ife glycemic control in patients with type 2 diabetes treated with insulin therapy: A prospective, longitudinal cohort study (Diabetes Distress and Care Registry at Tenri [DDCRT 9]). Journal of Diabetes Investigation, 2018, 9, 294-302.	1.1	1
328	Dialysisâ€related practice patterns among hemodialysis patients with cancer. Health Science Reports, 2018, 1, e46.	0.6	1
329	Intra-Dialytic Hemoglobin Changes and Cardiovascular Events: A Cohort Study on Dialysis Outcomes and Practice Patterns in Japan. American Journal of Nephrology, 2019, 50, 272-280.	1.4	1
330	Population-level associations of achievement of targets for bone-mineral markers with survival in haemodialysis patients with mildly elevated intact-PTH levels: a case-cohort study. Scientific Reports, 2019, 9, 11301.	1.6	1
331	The association between physician's affiliation and patients' adherence to their antihypertensive medication and pharmaceutical knowledge. Journal of General and Family Medicine, 2019, 20, 19-24.	0.3	1
332	Length of hospital stay is associated with a decline in activities of daily living in hemodialysis patients: a prospective cohort study. BMC Nephrology, 2020, 21, 9.	0.8	1
333	Screening tool for identifying adults with excessive salt intake among community-dwelling adults: a population-based cohort study. American Journal of Clinical Nutrition, 2020, 111, 814-820.	2.2	1
334	Negative impact of being underweight on the outcomes of single-session shockwave lithotripsy in patients with upper urinary tract calculi: a retrospective cohort study. World Journal of Urology, 2021, 39, 571-577.	1.2	1
335	Association between the use of physical restraint and functional decline among older inpatients admitted with pneumonia in an acute care hospital: A retrospective cohort study. Archives of Gerontology and Geriatrics, 2021, 94, 104330.	1.4	1
336	Prevalence and factors associated with uncorrected presbyopia in a rural population of Japan: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Japanese Journal of Ophthalmology, 2021, 65, 724-730.	0.9	1
337	Dose-response relationship between spino-pelvic alignment determined by sagittal modifiers and back pain-specific quality of life. European Spine Journal, 2021, 30, 3019-3027.	1.0	1
338	Validation Study of Gastrointestinal Symptom Rating Scale for Postoperative Patients with Upper Gastrointestinal Disease. Japanese Journal of Gastroenterological Surgery, 2014, 47, 365-373.	0.0	1
339	Cohort profile: Alliance for Quality Assessment in Healthcare-Dialysis (AQuAH-D) prospective cohort study of patients on haemodialysis in Japan. BMJ Open, 2022, 12, e054427.	0.8	1
340	Understanding by the General Public of Newspaper Reports on Publicly Reported Cancer Survival in Japan: A Randomized Controlled Trial. Japanese Journal of Clinical Oncology, 2014, 44, 651-660.	0.6	0
341	SP582ASSOCIATIONS OF EMPLOYMENT STATUS AND EDUCATIONAL LEVELS WITH MORTALITYAND HOSPITALIZATION IN THE DIALYSIS OUTCOMES AND PRACTICE PATTERNS STUDY IN JAPAN. Nephrology Dialysis Transplantation, 2016, 31, i288-i288.	0.4	0
342	MP618CLINICAL PREDICTION RULE FOR BACTEREMIA AMONG MAINTENANCE HEMODIALYSIS PATIENTS INOUTPATIENT SETTINGS. Nephrology Dialysis Transplantation, 2016, 31, i547-i547.	0.4	0

#	Article	IF	Citations
343	Detecting central-venous oxygen desaturation without a central-venous catheter: Utility of the difference between invasively and noninvasively measured blood pressure. Journal of Critical Care, 2016, 33, 257-261.	1.0	0
344	Response to the letter by Dr. Tomoyuki Kawada regarding our manuscript: Independent association between prediabetes and future pancreatic fat accumulation: a 5-year Japanese cohort study. Journal of Gastroenterology, 2018, 53, 891-892.	2.3	0
345	The Authors Reply. Kidney International, 2018, 94, 221-222.	2.6	O
346	FP365 QUALITY OF CARE AND INCIDENCE OF END-STAGE RENAL DISEASE IN THE ELDERLY CHRONIC KIDNEY DISEASE PATIENTS: COHORT STUDY FROM THE NATIONAL DATABASE OF HEALTH INSURANCE CLAIMS OF JAPAN. Nephrology Dialysis Transplantation, 2018, 33, i154-i155.	0.4	0
347	Longitudinal association between mental health and future antibiotic prescriptions in healthy adults: Results from the LOHAS. PLoS ONE, 2020, 15, e0240236.	1.1	0
348	Association of advanced glycation end-product accumulation with overactive bladder in community-dwelling elderly: AÂcross-sectional Sukagawa study. Asian Journal of Urology, 2021, 8, 189-196.	0.5	0
349	Association between single limb standing test results and healthcare costs among community-dwelling older adults. Archives of Gerontology and Geriatrics, 2021, 92, 104256.	1.4	0
350	Comparison of Health-Related Quality of Life Between Japanese and American Patients with Bladder Cancer as Measured by a Newly Developed Japanese Version of the Bladder Cancer Index. Bladder Cancer, 2021, 7, 61-69.	0.2	0
351	Corneal Eccentricity in a Rural Japanese Population: The Locomotive Syndrome and Health Outcome in Aizu Cohort Study (LOHAS). Ophthalmic Epidemiology, 2021, , 1-6.	0.8	0
352	400Overactive bladder with/without urinary incontinence and falls in the community-dwelling elderly. International Journal of Epidemiology, 2021, 50, .	0.9	0
353	Results of a survey on practice patterns at the end of a study involving CKD-MBD management at dialysis facilities. Nihon Toseki Igakkai Zasshi, 2013, 46, 193-200.	0.2	0
354	Japanese physicians' views of ethical problems in dialysis therapy and advance directives Nihon Toseki Igakkai Zasshi, 1998, 31, 1221-1225.	0.2	0
355	Descriptive epidemiology of prescriptions of drugs listed in Screening Tool for Older Person's appropriate Prescriptions in Japanese (STOPP-J) in older adults: results from National Database of Health Insurance Claims and Specific Health Checkups of Japan (NDB). Proceedings for Annual Meeting of the Japanese Pharmacological Society, 2020, 93, 2-AS1-3.	0.0	0
356	The prevalence of uncorrected refractive error in Japan: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Japanese Journal of Ophthalmology, 2022, 66, 199-204.	0.9	0
357	Title is missing!. , 2020, 15, e0240236.		0
358	Title is missing!. , 2020, 15, e0240236.		0
359	Title is missing!. , 2020, 15, e0240236.		0
360	Title is missing!. , 2020, 15, e0240236.		O

#	Article	IF	CITATIONS
361	Title is missing!. , 2020, 15, e0240236.		O
362	Title is missing!. , 2020, 15, e0240236.		0
363	Preoperative Use of Alpha-1 Receptor Blockers in Male Patients Undergoing Extracorporeal Shock Wave Lithotripsy for a Ureteral Calculus. Acta Medica Okayama, 2020, 74, 293-299.	0.1	0