

Shunichi Fukuhara

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11555338/publications.pdf>

Version: 2024-02-01

363
papers

15,192
citations

24978

57
h-index

23472

111
g-index

373
all docs

373
docs citations

373
times ranked

15533
citing authors

#	ARTICLE	IF	CITATIONS
1	Decline in oral antimicrobial prescription in the outpatient setting after nationwide implementation of financial incentives and provider education: An interrupted time-series analysis. <i>Infection Control and Hospital Epidemiology</i> , 2023, 44, 253-259.	1.0	2
2	Usefulness of computed tomography in the diagnosis of acute pyelonephritis in older patients suspected of infection with unknown focus. <i>Acta Radiologica</i> , 2022, 63, 268-277.	0.5	5
3	The prevalence of uncorrected refractive error in Japan: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. <i>Japanese Journal of Ophthalmology</i> , 2022, 66, 199-204.	0.9	0
4	Cohort profile: Alliance for Quality Assessment in Healthcare-Dialysis (AQuAH-D) prospective cohort study of patients on haemodialysis in Japan. <i>BMJ Open</i> , 2022, 12, e054427.	0.8	1
5	Association of kyphotic posture with loss of independence and mortality in a community-based prospective cohort study: the Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). <i>BMJ Open</i> , 2022, 12, e052421.	0.8	3
6	Usefulness of a mobile phone application for respiratory rate measurement in adult patients. <i>Japan Journal of Nursing Science</i> , 2022, 19, e12481.	0.5	4
7	Low quality of life in men with chronic prostatitis-like symptoms. <i>Prostate Cancer and Prostatic Diseases</i> , 2022, 25, 785-790.	2.0	5
8	Impact of lumbar spinal stenosis on metabolic syndrome incidence in community-dwelling adults in Aizu cohort study (LOHAS). <i>Scientific Reports</i> , 2022, 12, .	1.6	3
9	How Psychosocial Factors Contribute to Japanese Older Adults'™ Initiation of Advance Care Planning Discussions: A Qualitative Study. <i>Journal of Applied Gerontology</i> , 2021, 40, 1180-1188.	1.0	9
10	Negative impact of being underweight on the outcomes of single-session shockwave lithotripsy in patients with upper urinary tract calculi: a retrospective cohort study. <i>World Journal of Urology</i> , 2021, 39, 571-577.	1.2	1
11	Inverse Association Between Fatty Liver at Baseline Ultrasonography and Remission of Type 2 Diabetes Over a 2-Year Follow-up Period. <i>Clinical Gastroenterology and Hepatology</i> , 2021, 19, 556-564.e5.	2.4	11
12	Association of advanced glycation end-product accumulation with overactive bladder in community-dwelling elderly: A cross-sectional Sukagawa study. <i>Asian Journal of Urology</i> , 2021, 8, 189-196.	0.5	0
13	Association between single limb standing test results and healthcare costs among community-dwelling older adults. <i>Archives of Gerontology and Geriatrics</i> , 2021, 92, 104256.	1.4	0
14	Experience as an Informal Caregiver and Discussions Regarding Advance Care Planning in Japan. <i>Journal of Pain and Symptom Management</i> , 2021, 61, 63-70.	0.6	11
15	Quality of Life and Disability-free Survival in the Elderly: The Locomotive Syndrome and Health Outcome in Aizu Cohort Study. <i>Journal of Aging and Health</i> , 2021, 33, 197-204.	0.9	3
16	Urgency urinary incontinence, loss of independence, and increased mortality in older adults: A cohort study. <i>PLoS ONE</i> , 2021, 16, e0245724.	1.1	4
17	The Association between the Establishment of a General Internal Medicine Department and an Increased Number of Blood Cultures in Other Departments: An Interrupted Time Series Analysis. <i>Internal Medicine</i> , 2021, 60, 3729-3735.	0.3	2
18	Significance of Overactive Bladder as a Predictor of Falls in Community Dwelling Older Adults: 1-Year Followup of the Sukagawa Study. <i>Journal of Urology</i> , 2021, 205, 219-225.	0.2	3

#	ARTICLE	IF	CITATIONS
19	Added value of clinical prediction rules for bacteremia in hemodialysis patients: An external validation study. <i>PLoS ONE</i> , 2021, 16, e0247624.	1.1	3
20	Cohort profile: A super-elderly population-based cohort in Japan (the Sukagawa Study). <i>International Journal of Epidemiology</i> , 2021, 50, 727-727h.	0.9	5
21	Healthcare Providers'™ Perceptions on the Timing of Initial Advance Care Planning Discussions in Japan: a Mixed-Methods Study. <i>Journal of General Internal Medicine</i> , 2021, 36, 2935-2942.	1.3	8
22	Is "œpain before vomiting"•useful?: Diagnostic performance of the classic patient history item in acute appendicitis. <i>American Journal of Emergency Medicine</i> , 2021, 41, 84-89.	0.7	3
23	Association of glucocorticoid doses and emotional health in lupus low disease activity state (LLDAS): a cross-sectional study. <i>Arthritis Research and Therapy</i> , 2021, 23, 79.	1.6	11
24	Comparison of Health-Related Quality of Life Between Japanese and American Patients with Bladder Cancer as Measured by a Newly Developed Japanese Version of the Bladder Cancer Index. <i>Bladder Cancer</i> , 2021, 7, 61-69.	0.2	0
25	Association between the use of physical restraint and functional decline among older inpatients admitted with pneumonia in an acute care hospital: A retrospective cohort study. <i>Archives of Gerontology and Geriatrics</i> , 2021, 94, 104330.	1.4	1
26	Diagnostic performance of food consumption for bacteraemia in patients admitted with suspected infection: a prospective cohort study. <i>BMJ Open</i> , 2021, 11, e044270.	0.8	3
27	Impact of Vonoprazan Triple-Drug Blister Packs on H.Âpylori Eradication Rates in Japan: Interrupted Time Series Analysis. <i>Advances in Therapy</i> , 2021, 38, 3937-3947.	1.3	2
28	Effect of multimorbidity patterns on the decline in health-related quality of life: a nationwide prospective cohort study in Japan. <i>BMJ Open</i> , 2021, 11, e047812.	0.8	8
29	Poor oral function is associated with loss of independence or death in functionally independent older adults. <i>PLoS ONE</i> , 2021, 16, e0253559.	1.1	4
30	Changes in Drug Utilization After Publication of Clinical Trials and Drug-Related Scandals in Japan: An Interrupted Time Series Analysis, 2005â€“2017. <i>Journal of Epidemiology</i> , 2021, 31, 410-416.	1.1	3
31	Impact of Primary Tumor Resection on Mortality in Patients with Stage IV Colorectal Cancer with Unresectable Metastases: A Multicenter Retrospective Cohort Study. <i>World Journal of Surgery</i> , 2021, 45, 3230-3239.	0.8	3
32	Prevalence and factors associated with uncorrected presbyopia in a rural population of Japan: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. <i>Japanese Journal of Ophthalmology</i> , 2021, 65, 724-730.	0.9	1
33	Association between religious beliefs and discussions regarding advance care planning: A nationwide survey. <i>Palliative Medicine</i> , 2021, 35, 1856-1864.	1.3	13
34	Intraoperative end-tidal carbon dioxide and postoperative mortality in major abdominal surgery: a historical cohort study. <i>Canadian Journal of Anaesthesia</i> , 2021, 68, 1601-1610.	0.7	5
35	Corneal Eccentricity in a Rural Japanese Population: The Locomotive Syndrome and Health Outcome in Aizu Cohort Study (LOHAS). <i>Ophthalmic Epidemiology</i> , 2021, , 1-6.	0.8	0
36	Dose-response relationship between spino-pelvic alignment determined by sagittal modifiers and back pain-specific quality of life. <i>European Spine Journal</i> , 2021, 30, 3019-3027.	1.0	1

#	ARTICLE	IF	CITATIONS
37	400Overactive bladder with/without urinary incontinence and falls in the community-dwelling elderly. <i>International Journal of Epidemiology</i> , 2021, 50, .	0.9	0
38	Association between facility-level adherence to phosphorus management guidelines and mortality in haemodialysis patients: a prospective cohort study. <i>BMJ Open</i> , 2021, 11, e051002.	0.8	2
39	Development and validation of a concise scale for assessing patient experience of primary care for adults in Japan. <i>Family Practice</i> , 2020, 37, 137-142.	0.8	18
40	The development of quality indicators for systemic lupus erythematosus using electronic health data: A modified RAND appropriateness method. <i>Modern Rheumatology</i> , 2020, 30, 525-531.	0.9	6
41	Association between contrast extravasation on computed tomography scans and pseudoaneurysm formation in pediatric blunt splenic and hepatic injury: A multi-institutional observational study. <i>Journal of Pediatric Surgery</i> , 2020, 55, 681-687.	0.8	13
42	Length of hospital stay is associated with a decline in activities of daily living in hemodialysis patients: a prospective cohort study. <i>BMC Nephrology</i> , 2020, 21, 9.	0.8	1
43	Early rehabilitation in older patients hospitalized with acute decompensated heart failure: A retrospective cohort study. <i>American Heart Journal</i> , 2020, 230, 44-53.	1.2	15
44	Longitudinal association between mental health and future antibiotic prescriptions in healthy adults: Results from the LOHAS. <i>PLoS ONE</i> , 2020, 15, e0240236.	1.1	0
45	Interdialytic Weight Gain Effects on Hemoglobin Concentration and Cardiovascular Events. <i>Kidney International Reports</i> , 2020, 5, 1670-1678.	0.4	8
46	Changes in 3-month mineral and bone disorder patterns were associated with all-cause mortality in prevalent hemodialysis patients with secondary hyperparathyroidism. <i>BMC Nephrology</i> , 2020, 21, 432.	0.8	4
47	Quality of Care in Chronic Kidney Disease and Incidence of End-stage Renal Disease in Older Patients. <i>Medical Care</i> , 2020, 58, 625-631.	1.1	4
48	Physical multimorbidity patterns and depressive symptoms: a nationwide cross-sectional study in Japan. <i>Family Medicine and Community Health</i> , 2020, 8, e000234.	0.6	6
49	Health-related quality of life in Japanese patients with bladder cancer measured by a newly developed Japanese version of the Bladder Cancer Index. <i>International Journal of Clinical Oncology</i> , 2020, 25, 2090-2098.	1.0	2
50	Association between health-related hope and adherence to prescribed treatment in CKD patients: multicenter cross-sectional study. <i>BMC Nephrology</i> , 2020, 21, 453.	0.8	12
51	Association between work schedules and motivation for lifestyle change in workers with overweight or obesity: a cross-sectional study in Japan. <i>BMJ Open</i> , 2020, 10, e033000.	0.8	3
52	Patients' preferences and factors influencing initial advance care planning discussions' timing: A cross-cultural mixed-methods study. <i>Palliative Medicine</i> , 2020, 34, 906-916.	1.3	22
53	Associations of types of primary care facilities with adult vaccination and cancer screening in Japan. <i>International Journal for Quality in Health Care</i> , 2020, 32, 373-378.	0.9	2
54	Comparison of Primary Care Experience in Hospital-Based Practices and Community-Based Office Practices in Japan. <i>Annals of Family Medicine</i> , 2020, 18, 24-29.	0.9	13

#	ARTICLE	IF	CITATIONS
55	Low Predialysis Plasma Calculated Osmolality Is Associated with Higher All-Cause Mortality: The Japanese Dialysis Outcomes and Practice Patterns Study (J-DOPPS). <i>Nephron</i> , 2020, 144, 138-146.	0.9	6
56	Screening tool for identifying adults with excessive salt intake among community-dwelling adults: a population-based cohort study. <i>American Journal of Clinical Nutrition</i> , 2020, 111, 814-820.	2.2	1
57	Longitudinal association of fatty pancreas with the incidence of type-2 diabetes in lean individuals: a 6-year computed tomography-based cohort study. <i>Journal of Gastroenterology</i> , 2020, 55, 712-721.	2.3	26
58	Relationship between serum calcium or phosphate levels and mortality stratified by parathyroid hormone level: an analysis from the MBD-5D study. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 630-637.	0.7	6
59	Type I Angiotensin II Receptor Blockade Reduces Uremia-Induced Deterioration of Bone Material Properties. <i>Journal of Bone and Mineral Research</i> , 2020, 36, 67-79.	3.1	11
60	Knowledge Does Not Correlate with Behavior toward Deceased Organ Donation: A Cross-Sectional Study in Japan. <i>Annals of Transplantation</i> , 2020, 25, e918936.	0.5	6
61	Low Back Pain and Associated Occupational Factors among Nursery School Teachers: A Multicenter, Cross-sectional Study. <i>Annals of Clinical Epidemiology</i> , 2020, 2, 113-120.	0.3	5
62	Association of Lumbar Spinal Stenosis With Severe Disability and Mortality Among Community-Dwelling Older Adults. <i>Spine</i> , 2020, Publish Ahead of Print, E784-E790.	1.0	3
63	Descriptive epidemiology of prescriptions of drugs listed in Screening Tool for Older Person's appropriate Prescriptions in Japanese (STOPP-J) in older adults: results from National Database of Health Insurance Claims and Specific Health Checkups of Japan (NDB). <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2020, 93, 2-AS1-3.	0.0	0
64	Title is missing!. , 2020, 15, e0240236.		0
65	Title is missing!. , 2020, 15, e0240236.		0
66	Title is missing!. , 2020, 15, e0240236.		0
67	Title is missing!. , 2020, 15, e0240236.		0
68	Title is missing!. , 2020, 15, e0240236.		0
69	Title is missing!. , 2020, 15, e0240236.		0
70	Preoperative Use of Alpha-1 Receptor Blockers in Male Patients Undergoing Extracorporeal Shock Wave Lithotripsy for a Ureteral Calculus. <i>Acta Medica Okayama</i> , 2020, 74, 293-299.	0.1	0
71	Determinants of incentive preferences for health behavior change in Japan. <i>Health Promotion International</i> , 2019, 34, 760-769.	0.9	2
72	Gait speed and overactive bladder in the healthy community-dwelling super elderly: The Sukagawa Study. <i>Neurourology and Urodynamics</i> , 2019, 38, 2324-2332.	0.8	13

#	ARTICLE	IF	CITATIONS
73	Statistical significance did not affect time to publication in non-Cochrane systematic reviews: a metaepidemiological study. <i>Journal of Clinical Epidemiology</i> , 2019, 115, 25-34.	2.4	4
74	Salt intake and body weight correlate with higher blood pressure in the very elderly population: The Sukagawa study. <i>Journal of Clinical Hypertension</i> , 2019, 21, 942-949.	1.0	10
75	Association between parental history of <i>Helicobacter pylori</i> treatment failure and treatment failure in the offspring. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2019, 34, 2112-2117.	1.4	8
76	Intra-Dialytic Hemoglobin Changes and Cardiovascular Events: A Cohort Study on Dialysis Outcomes and Practice Patterns in Japan. <i>American Journal of Nephrology</i> , 2019, 50, 272-280.	1.4	1
77	Population-level associations of achievement of targets for bone-mineral markers with survival in haemodialysis patients with mildly elevated intact-PTH levels: a case-cohort study. <i>Scientific Reports</i> , 2019, 9, 11301.	1.6	1
78	Vehicle configurations associated with anatomical-specific severe injuries resulting from traffic collisions. <i>PLoS ONE</i> , 2019, 14, e0223388.	1.1	2
79	Hospitals'™ extracorporeal cardiopulmonary resuscitation capabilities and outcomes in out-of-hospital cardiac arrest: A population-based study. <i>Resuscitation</i> , 2019, 136, 85-92.	1.3	15
80	Effectiveness of cinacalcet treatment for secondary hyperparathyroidism on hospitalization: Results from the MBD-5D study. <i>PLoS ONE</i> , 2019, 14, e0216399.	1.1	2
81	Postdialysis Hypokalemia and All-Cause Mortality in Patients Undergoing Maintenance Hemodialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019, 14, 873-881.	2.2	21
82	Association between social networks and discussions regarding advance care planning among Japanese older adults. <i>PLoS ONE</i> , 2019, 14, e0213894.	1.1	14
83	Level of Low Back Pain-Related Disability Is Associated with Risk of Subsequent Falls in an Older Population: Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). <i>Pain Medicine</i> , 2019, 20, 2377-2384.	0.9	15
84	The association between physician's affiliation and patients'™ adherence to their antihypertensive medication and pharmaceutical knowledge. <i>Journal of General and Family Medicine</i> , 2019, 20, 19-24.	0.3	1
85	Factors associated with patient preferences towards deprescribing: a survey of adult patients on prescribed medications. <i>International Journal of Clinical Pharmacy</i> , 2019, 41, 531-537.	1.0	27
86	Association of low alanine aminotransferase with loss of independence or death: A 5-year population-based cohort study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2019, 34, 1793-1799.	1.4	13
87	Haemoglobin concentration and survival of haemodialysis patients before and after experiencing cardiovascular disease: a cohort study from Japanese dialysis outcomes and practice pattern study (J-DOPPS). <i>BMJ Open</i> , 2019, 9, e031476.	0.8	3
88	Social participation and the combination of future needs for long-term care and mortality among older Japanese people: a prospective cohort study from the Aichi Gerontological Evaluation Study (AGES). <i>BMJ Open</i> , 2019, 9, e030500.	0.8	17
89	Ulcerative colitis outcomes research in Japan: protocol for an observational prospective cohort study of YOURS (YOu and Ulcerative colitis: Registry and Social network). <i>BMJ Open</i> , 2019, 9, e030134.	0.8	4
90	The minimum clinically important difference for the Japanese version of the new Knee Society Score (2011KSS) after total knee arthroplasty. <i>Journal of Orthopaedic Science</i> , 2019, 24, 1053-1057.	0.5	23

#	ARTICLE	IF	CITATIONS
91	Vitality Measured as Self-reported Energy Level and Clinical Outcomes in Hemodialysis Patients: The Japanese Dialysis Outcomes and Practice Pattern Study (J-DOPPS). <i>American Journal of Kidney Diseases</i> , 2019, 73, 486-495.	2.1	18
92	Quality of life after single-incision laparoscopic cholecystectomy: A randomized, clinical trial. <i>Surgery</i> , 2019, 165, 353-359.	1.0	20
93	Association Between the Discrepancy in Self-Reported and Performance-Based Physical Functioning Levels and Risk of Future Falls Among Community-Dwelling Older Adults: The Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). <i>Journal of the American Medical Directors Association</i> , 2019, 20, 195-200.e1.	1.2	9
94	Longitudinal Association Between Subjective Fatigue and Future Falls in Community-Dwelling Older Adults: The Locomotive Syndrome and Health Outcomes in the Aizu Cohort Study (LOHAS). <i>Journal of Aging and Health</i> , 2019, 31, 67-84.	0.9	17
95	A scale for measuring health-related hope: its development and psychometric testing. <i>Annals of Clinical Epidemiology</i> , 2019, 1, 102-119.	0.3	12
96	Childcare Environment and Japanese Children Who Are Overweight in Early Childhood. <i>Childhood Obesity</i> , 2018, 14, 197-206.	0.8	1
97	Association between daily salt intake of 3-year-old children and that of their mothers: A cross-sectional study. <i>Journal of Clinical Hypertension</i> , 2018, 20, 730-735.	1.0	2
98	Body Mass Index Change and Hospitalization Risk in Elderly Hemodialysis Patients: Results from Japanese Dialysis Outcomes and Practice Patterns Study. <i>American Journal of Nephrology</i> , 2018, 47, 48-56.	1.4	5
99	Association between keeping home records of catheter exit-site and incidence of peritoneal dialysis-related infections. <i>International Urology and Nephrology</i> , 2018, 50, 763-769.	0.6	3
100	Effect of Patient Experience on Bypassing a Primary Care Gatekeeper: a Multicenter Prospective Cohort Study in Japan. <i>Journal of General Internal Medicine</i> , 2018, 33, 722-728.	1.3	27
101	Effects of self-monitoring of daily salt intake estimated by a simple electrical device for salt reduction: a cluster randomized trial. <i>Hypertension Research</i> , 2018, 41, 524-530.	1.5	13
102	Association between pancreatic fat and incidence of metabolic syndrome: a 5-year Japanese cohort study. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2018, 33, 2048-2054.	1.4	16
103	Chemotherapy in cancer patients undergoing haemodialysis: a nationwide study in Japan. <i>ESMO Open</i> , 2018, 3, e000301.	2.0	24
104	Multimorbidity patterns in relation to polypharmacy and dosage frequency: a nationwide, cross-sectional study in a Japanese population. <i>Scientific Reports</i> , 2018, 8, 3806.	1.6	73
105	Impact of trajectories of abdominal aortic calcification over 2 years on subsequent mortality: a 10-year longitudinal study. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 676-683.	0.4	11
106	Real-life glycemic control in patients with type 2 diabetes treated with insulin therapy: A prospective, longitudinal cohort study (Diabetes Distress and Care Registry at Tenri [DDCRT 9]). <i>Journal of Diabetes Investigation</i> , 2018, 9, 294-302.	1.1	1
107	Association Between Subjective Sleep Quality and Future Risk of Falls in Older People: Results From LOHAS. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2018, 73, 1205-1211.	1.7	44
108	Association Between Accumulation of Advanced Glycation End-Products and Hearing Impairment in Community-Dwelling Older People: A Cross-Sectional Sukagawa Study. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 235-239.e1.	1.2	5

#	ARTICLE	IF	CITATIONS
109	Independent association between prediabetes and future pancreatic fat accumulation: a 5-year Japanese cohort study. <i>Journal of Gastroenterology</i> , 2018, 53, 873-882.	2.3	18
110	Social Isolation and Patient Experience in Older Adults. <i>Annals of Family Medicine</i> , 2018, 16, 393-398.	0.9	24
111	Dialysis physicians' referral behaviors for hemodialysis patients suspected of having cancer: A vignette-based questionnaire study. <i>PLoS ONE</i> , 2018, 13, e0202322.	1.1	2
112	Association between post-dialysis hemoglobin level and the survival of vascular access. <i>Journal of Vascular Access</i> , 2018, 19, 69-75.	0.5	2
113	Response to the letter by Dr. Tomoyuki Kawada regarding our manuscript: Independent association between prediabetes and future pancreatic fat accumulation: a 5-year Japanese cohort study. <i>Journal of Gastroenterology</i> , 2018, 53, 891-892.	2.3	0
114	Development and validation of a prediction model for functional decline in older medical inpatients. <i>Archives of Gerontology and Geriatrics</i> , 2018, 77, 184-188.	1.4	2
115	Heart Rate, Responsiveness to Intravenous Immunoglobulin, and Coronary Artery Aneurysms in Kawasaki Disease. <i>Journal of Pediatrics</i> , 2018, 200, 160-165.e5.	0.9	4
116	Whole-Body Computed Tomography During Initial Management and Mortality Among Adult Severe Blunt Trauma Patients: A Nationwide Cohort Study. <i>World Journal of Surgery</i> , 2018, 42, 3939-3946.	0.8	16
117	The Authors Reply. <i>Kidney International</i> , 2018, 94, 221-222.	2.6	0
118	Development and validation of a prediction model for loss of physical function in elderly hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 1452-1458.	0.4	9
119	Association between vision-specific quality of life and falls in community-dwelling older adults: LOHAS. <i>PLoS ONE</i> , 2018, 13, e0195806.	1.1	12
120	Public subsidies and the recommendation of child vaccines among primary care physicians: a nationwide cross-sectional study in Japan. <i>BMJ Open</i> , 2018, 8, e020923.	0.8	4
121	Increased Mortality Rate after Hospitalization Among Chronic Hemodialysis Patients: A Prospective Cohort Study. <i>Nephron</i> , 2018, 140, 194-202.	0.9	7
122	Salt Intake and All-Cause Mortality in Hemodialysis Patients. <i>American Journal of Nephrology</i> , 2018, 48, 87-95.	1.4	15
123	FP365 QUALITY OF CARE AND INCIDENCE OF END-STAGE RENAL DISEASE IN THE ELDERLY CHRONIC KIDNEY DISEASE PATIENTS: COHORT STUDY FROM THE NATIONAL DATABASE OF HEALTH INSURANCE CLAIMS OF JAPAN. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i154-i155.	0.4	0
124	Dialysis-related practice patterns among hemodialysis patients with cancer. <i>Health Science Reports</i> , 2018, 1, e46.	0.6	1
125	Development of quality indicators for care of chronic kidney disease in the primary care setting using electronic health data: a RAND-modified Delphi method. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 247-256.	0.7	24
126	Unique hemoglobin A1c level distribution and its relationship with mortality in diabetic hemodialysis patients. <i>Kidney International</i> , 2017, 92, 497-503.	2.6	19

#	ARTICLE	IF	CITATIONS
127	Attributes of primary care in relation to polypharmacy: a multicenter cross-sectional study in Japan. <i>International Journal for Quality in Health Care</i> , 2017, 29, 378-383.	0.9	6
128	The cardiothoracic ratio and all-cause and cardiovascular disease mortality in patients undergoing maintenance hemodialysis: results of the MBD-5D study. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 797-806.	0.7	15
129	Effect of ABO blood type on the outcomes of patients with metastatic renal cell carcinoma treated with first-line tyrosine kinase inhibitors. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017, 35, 540.e7-540.e12.	0.8	1
130	Majority of systematic reviews published in high-impact journals neglected to register the protocols: a meta-epidemiological study. <i>Journal of Clinical Epidemiology</i> , 2017, 84, 54-60.	2.4	41
131	External validation of prognostic indices for overall survival of malignant pleural mesothelioma. <i>Lung Cancer</i> , 2017, 113, 88-92.	0.9	3
132	Patient experience of primary care and advance care planning: a multicentre cross-sectional study in Japan. <i>Family Practice</i> , 2017, 34, 206-212.	0.8	39
133	Effects of Higher Quality of Care on Initiation of Long-term Dialysis in Patients With CKD and Diabetes. <i>American Journal of Kidney Diseases</i> , 2017, 70, 666-674.	2.1	12
134	Impact of nutritional index on the association between phosphorus concentrations and mortality in haemodialysis patients: a cohort study from dialysis outcomes and practice pattern study in Japan. <i>BMJ Open</i> , 2017, 7, e016682.	0.8	12
135	Diagnostic utility of appetite loss in addition to existing prediction models for community-acquired pneumonia in the elderly: a prospective diagnostic study in acute care hospitals in Japan. <i>BMJ Open</i> , 2017, 7, e019155.	0.8	5
136	Interactive Effectiveness of Angiotensin-Converting Enzyme Inhibitors and Angiotensin Receptor Blockers or Their Combination on Survival of Hemodialysis Patients. <i>American Journal of Nephrology</i> , 2017, 46, 439-447.	1.4	6
137	Association between statistical significance and time to publication among systematic reviews: a study protocol for a meta-epidemiological investigation. <i>BMJ Open</i> , 2017, 7, e018856.	0.8	6
138	Revisiting Interdialytic Weight Gain and Mortality Association With Serum Albumin Interactions: The Japanese Dialysis Outcomes and Practice Pattern Study. , 2017, 27, 421-429.		32
139	Analysis of the evidence-practice gap to facilitate proper medical care for the elderly: investigation, using databases, of utilization measures for National Database of Health Insurance Claims and Specific Health Checkups of Japan (NDB). <i>Environmental Health and Preventive Medicine</i> , 2017, 22, 51.	1.4	27
140	Both low and high serum ferritin levels predict mortality risk in hemodialysis patients without inflammation. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 685-693.	0.7	29
141	Long-term Trends in Primary Sites of Gastric Adenocarcinoma in Japan and the United States. <i>Journal of Cancer</i> , 2017, 8, 1935-1942.	1.2	23
142	Development of a profile scoring system for assessing the psychosocial situation of patients with chronic musculoskeletal pain. <i>Journal of Pain Research</i> , 2017, Volume 10, 1853-1859.	0.8	3
143	Development of a new risk model for predicting cardiovascular events among hemodialysis patients: Population-based hemodialysis patients from the Japan Dialysis Outcome and Practice Patterns Study (J-DOPPS). <i>PLoS ONE</i> , 2017, 12, e0173468.	1.1	10
144	Computed tomography during initial management and mortality among hemodynamically unstable blunt trauma patients: a nationwide retrospective cohort study. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2017, 25, 74.	1.1	17

#	ARTICLE	IF	CITATIONS
145	Development and Validation of a Clinical Prediction Rule for Bacteremia among Maintenance Hemodialysis Patients in Outpatient Settings. PLoS ONE, 2017, 12, e0169975.	1.1	8
146	Associations of employment status and educational levels with mortality and hospitalization in the dialysis outcomes and practice patterns study in Japan. PLoS ONE, 2017, 12, e0170731.	1.1	19
147	Association of coping strategies with mortality and health-related quality of life in hemodialysis patients: The Japan Dialysis Outcomes and Practice Patterns Study. PLoS ONE, 2017, 12, e0180498.	1.1	26
148	Shorter dialysis session length was not associated with lower mental health and physical functioning in elderly hemodialysis patients: Results from the Japan Dialysis Outcome and Practice Patterns Study (J-DOPPS). PLoS ONE, 2017, 12, e0184019.	1.1	3
149	Fukushima after the Great East Japan Earthquake: lessons for developing responsive and resilient health systems. Journal of Global Health, 2017, 7, 010501.	1.2	31
150	Identifying Patients with Bacteremia in Community-Hospital Emergency Rooms: A Retrospective Cohort Study. PLoS ONE, 2016, 11, e0148078.	1.1	27
151	SP582ASSOCIATIONS OF EMPLOYMENT STATUS AND EDUCATIONAL LEVELS WITH MORTALITYAND HOSPITALIZATION IN THE DIALYSIS OUTCOMES AND PRACTICE PATTERNS STUDY IN JAPAN. Nephrology Dialysis Transplantation, 2016, 31, i288-i288.	0.4	0
152	MP618CLINICAL PREDICTION RULE FOR BACTEREMIA AMONG MAINTENANCE HEMODIALYSIS PATIENTS INOUTPATIENT SETTINGS. Nephrology Dialysis Transplantation, 2016, 31, i547-i547.	0.4	0
153	Body Temperature at the Emergency Department as a Predictor of Mortality in Patients With Bacterial Infection. Medicine (United States), 2016, 95, e3628.	0.4	29
154	Relationship Between Kyphotic Posture and Falls in Community-Dwelling Men and Women. Spine, 2016, 41, 1232-1238.	1.0	16
155	Retinal vessel diameters in a Japanese population: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Acta Ophthalmologica, 2016, 94, e432-41.	0.6	13
156	Cardiovascular and Stroke Risk in Japanese Hemodialysis Patients With Atrial Fibrillation. Therapeutic Apheresis and Dialysis, 2016, 20, 608-614.	0.4	3
157	Influence of family dynamics on burden among family caregivers in aging Japan. Family Practice, 2016, 33, 466-470.	0.8	13
158	PTH-dependence of the effectiveness of cinacalcet in hemodialysis patients with secondary hyperparathyroidism. Scientific Reports, 2016, 6, 19612.	1.6	47
159	Low Testosterone Levels and Reduced Kidney Function in Japanese Adult Men: The Locomotive Syndrome and Health Outcome in Aizu Cohort Study. Journal of the American Medical Directors Association, 2016, 17, 371.e1-371.e6.	1.2	20
160	Effects of structured education program on organ donor designation of nursing students and their families: A randomized controlled trial. Clinical Transplantation, 2016, 30, 1513-1519.	0.8	12
161	Influence of Staff Encouragement on Perceived Burden of Dietary Restriction Among Patients Living Alone. Therapeutic Apheresis and Dialysis, 2016, 20, 623-631.	0.4	2
162	Preferences of young physicians at community hospitals regarding academic research training through graduate school: a cross-sectional research. BMC Research Notes, 2016, 9, 227.	0.6	2

#	ARTICLE	IF	CITATIONS
163	Frequent monitoring of mineral metabolism in hemodialysis patients with secondary hyperparathyroidism: associations with achievement of treatment goals and with adjustments in therapy. <i>Nephrology Dialysis Transplantation</i> , 2016, 32, gfw020.	0.4	10
164	Detecting central-venous oxygen desaturation without a central-venous catheter: Utility of the difference between invasively and noninvasively measured blood pressure. <i>Journal of Critical Care</i> , 2016, 33, 257-261.	1.0	0
165	Influence of urinary urgency and other urinary disturbances on falls in Parkinson's disease. <i>Journal of the Neurological Sciences</i> , 2016, 360, 153-157.	0.3	31
166	Functional Dependence and Mortality in the International Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>American Journal of Kidney Diseases</i> , 2016, 67, 283-292.	2.1	110
167	Effects of cyclophosphamide on the prognosis of Japanese patients with renal vasculitis associated with anti-neutrophil cytoplasmic antibody-positive microscopic polyangiitis. <i>Clinical and Experimental Nephrology</i> , 2016, 20, 712-719.	0.7	8
168	Effects of Secondary Hyperparathyroidism Treatment on Improvement in Anemia: Results from the MBD-5D Study. <i>PLoS ONE</i> , 2016, 11, e0164865.	1.1	25
169	Minor Elevation in C-Reactive Protein Levels Predicts Incidence of Erythropoiesis-Stimulating Agent Hyporesponsiveness among Hemodialysis Patients. <i>Nephron</i> , 2015, 131, 123-130.	0.9	22
170	Chronic hyperglycemia increases the risk of lateral epicondylitis: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study (LOHAS). <i>SpringerPlus</i> , 2015, 4, 407.	1.2	19
171	Dietary Patterns and Clinical Outcomes in Hemodialysis Patients in Japan: A Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0116677.	1.1	18
172	The Role of Digital Rectal Examination for Diagnosis of Acute Appendicitis: A Systematic Review and Meta-Analysis. <i>PLoS ONE</i> , 2015, 10, e0136996.	1.1	18
173	Association between hand-grip strength and depressive symptoms: Locomotive Syndrome and Health Outcomes in Aizu Cohort Study (LOHAS). <i>Age and Ageing</i> , 2015, 44, 592-598.	0.7	130
174	Contribution of dysregulated serum magnesium to mortality in hemodialysis patients with secondary hyperparathyroidism: a 3-year cohort study. <i>CKJ: Clinical Kidney Journal</i> , 2015, 8, 744-752.	1.4	18
175	Dialysis Amyloid Deposition in the Aortic Valve and Its Association with Aortic Stenosis. <i>Blood Purification</i> , 2015, 40, 146-154.	0.9	5
176	Development and Validation of a Disease-Specific Instrument to Measure Diet-Targeted Quality of Life for Postoperative Patients with Esophagogastric Cancer. <i>Annals of Surgical Oncology</i> , 2015, 22, 848-854.	0.7	12
177	Patient-reported disability in the general Japanese population was associated with medical care visits for low back pain, regardless of pain intensity. <i>Journal of Orthopaedic Science</i> , 2015, 20, 742-749.	0.5	10
178	A new prognostic index for overall survival in malignant pleural mesothelioma: the rPHS (regimen,) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50	0.6	15
179	Mentoring the next generation of physician-scientists in Japan: a cross-sectional survey of mentees in six academic medical centers. <i>BMC Medical Education</i> , 2015, 15, 54.	1.0	20
180	Relationship of type of work with health-related quality of life. <i>Quality of Life Research</i> , 2015, 24, 2927-2932.	1.5	9

#	ARTICLE	IF	CITATIONS
181	Prognostic utility of serum CRP levels in combination with CURB-65 in patients with clinically suspected sepsis: a decision curve analysis. <i>BMJ Open</i> , 2015, 5, e007049-e007049.	0.8	18
182	Prescription of potentially inappropriate medications to elderly hemodialysis patients: prevalence and predictors. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 498-505.	0.4	29
183	Effect of an Educational Program on Attitudes towards Deceased Organ Donation. <i>Annals of Transplantation</i> , 2015, 20, 269-278.	0.5	22
184	Use of Renin-Angiotensin System Inhibitors Is Associated with Reduction of Fracture Risk in Hemodialysis Patients. <i>PLoS ONE</i> , 2015, 10, e0122691.	1.1	39
185	Association of Serum Bicarbonate with Bone Fractures in Hemodialysis Patients: The Mineral and Bone Disorder Outcomes Study for Japanese CKD Stage 5D Patients (MBD-5D). <i>Nephron Clinical Practice</i> , 2014, 128, 79-87.	2.3	12
186	New oral anticoagulants versus warfarin for preventing stroke and systemic embolic events among atrial fibrillation patients with chronic kidney disease. <i>The Cochrane Library</i> , 2014, , .	1.5	3
187	Understanding by the General Public of Newspaper Reports on Publicly Reported Cancer Survival in Japan: A Randomized Controlled Trial. <i>Japanese Journal of Clinical Oncology</i> , 2014, 44, 651-660.	0.6	0
188	Assessment of vision-related quality of life among patients with cataracts and the outcomes of cataract surgery using a newly developed visual function questionnaire: the VFQ-J11. <i>Japanese Journal of Ophthalmology</i> , 2014, 58, 415-422.	0.9	16
189	One-Year Period Prevalence of Oral Aphthous Ulcers and Oral Health-Related Quality of Life in Patients with Behçet's Disease. <i>Genetics Research International</i> , 2014, 2014, 1-8.	2.0	18
190	Lumbar Spinal Stenosis-Specific Symptom Scale. <i>Spine</i> , 2014, 39, E1388-E1393.	1.0	12
191	Glycated Hemoglobin and Risk of Death in Diabetic Patients Treated With Hemodialysis: A Meta-analysis. <i>American Journal of Kidney Diseases</i> , 2014, 63, 84-94.	2.1	72
192	Low Testosterone Levels, Depressive Symptoms, and Falls in Older Men: A Cross-Sectional Study. <i>Journal of the American Medical Directors Association</i> , 2014, 15, 30-35.	1.2	16
193	Association of blood pressure with all-cause mortality and stroke in Japanese hemodialysis patients: The Japanese dialysis outcomes and practice pattern study. <i>Hemodialysis International</i> , 2014, 18, 607-615.	0.4	21
194	Gender, low Kt/V, and mortality in Japanese hemodialysis patients: Opportunities for improvement through modifiable practices. <i>Hemodialysis International</i> , 2014, 18, 596-606.	0.4	34
195	Development and Validation of a Symptom Scale to Evaluate Postoperative Patients with Esophagogastric Cancer. <i>Journal of the American College of Surgeons</i> , 2014, 219, 895-903.	0.2	21
196	Pruritus in hemodialysis patients: Results from the Japanese Dialysis Outcomes and Practice Patterns Study (JDOPPS). <i>Hemodialysis International</i> , 2014, 18, 657-667.	0.4	100
197	Developing a scale to measure family dynamics related to long-term care, and testing that scale in a multicenter cross-sectional study. <i>BMC Family Practice</i> , 2014, 15, 134.	2.9	1
198	Association between allopurinol and mortality among Japanese hemodialysis patients: results from the DOPPS. <i>International Urology and Nephrology</i> , 2014, 46, 1833-1841.	0.6	24

#	ARTICLE	IF	CITATIONS
199	Abnormal Mineral Metabolism and Mortality in Hemodialysis Patients With Secondary Hyperparathyroidism: Evidence From Marginal Structural Models Used to Adjust for Time-Dependent Confounding. <i>American Journal of Kidney Diseases</i> , 2014, 63, 979-987.	2.1	64
200	Work Performance Assessed by a Newly Developed Japanese Version of the Work Limitation Questionnaire in a General Japanese Adult Population. <i>Journal of Occupational Health</i> , 2014, 56, 124-133.	1.0	23
201	Validation Study of Gastrointestinal Symptom Rating Scale for Postoperative Patients with Upper Gastrointestinal Disease. <i>Japanese Journal of Gastroenterological Surgery</i> , 2014, 47, 365-373.	0.0	1
202	Cost-utility analysis of cataract surgery in Japan: a probabilistic Markov modeling study. <i>Japanese Journal of Ophthalmology</i> , 2013, 57, 391-401.	0.9	18
203	Interpersonal Psychosocial Factors Associated With Underreported Dietary Energy Intake in Hemodialysis Patients. , 2013, 23, 37-44.		4
204	Prevalence of lumbar spinal stenosis, using the diagnostic support tool, and correlated factors in Japan: a population-based study. <i>Journal of Orthopaedic Science</i> , 2013, 18, 893-900.	0.5	101
205	Clinical Usefulness of the Two-Question Assessment Tool for Depressive Symptoms in Japanese Patients with Chronic Obstructive Pulmonary Disease. <i>Lung</i> , 2013, 191, 101-107.	1.4	4
206	Overactive bladder symptom severity is associated with falls in community-dwelling adults: LOHAS study. <i>BMJ Open</i> , 2013, 3, e002413.	0.8	36
207	Associations of Frequency and Duration of Patient-Doctor Contact in Hemodialysis Facilities with Mortality. <i>Journal of the American Society of Nephrology: JASN</i> , 2013, 24, 1493-1502.	3.0	33
208	Vascular Access Care and Treatment Practices Associated with Outcomes of Arteriovenous Fistula: International Comparisons from the Dialysis Outcomes and Practice Patterns Study. <i>Nephron Clinical Practice</i> , 2013, 124, 23-30.	2.3	56
209	Prevalence of pterygium in a population in Northern Japan: the Locomotive Syndrome and Health Outcome in Aizu Cohort Study. <i>Acta Ophthalmologica</i> , 2013, 91, e232-6.	0.6	31
210	Major bleeding events and risk stratification of antithrombotic agents in hemodialysis: results from the DOPPS. <i>Kidney International</i> , 2013, 84, 600-608.	2.6	83
211	Impact of cinacalcet introduction on MBD management: the MBD-5D study in Japan. <i>Kidney International Supplements</i> , 2013, 3, 436-441.	4.6	15
212	Evaluation of Newspaper Articles for Coverage of Public Reporting Data: A Case Study of Unadjusted Cancer Survival Data. <i>Japanese Journal of Clinical Oncology</i> , 2013, 43, 95-100.	0.6	3
213	Development of a Short Version of the Visual Function Questionnaire Using Item-Response Theory. <i>PLoS ONE</i> , 2013, 8, e73084.	1.1	12
214	Results of a survey on practice patterns at the end of a study involving CKD-MBD management at dialysis facilities. <i>Nihon Toseki Igakkai Zasshi</i> , 2013, 46, 193-200.	0.2	0
215	Prescription Patterns and Mineral Metabolism Abnormalities in the Cinacalcet Era: Results from the MBD-5D Study. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012, 7, 1473-1480.	2.2	29
216	Disappearance of Association in Diabetic Patients on Hemodialysis between Anemia and Mortality Risk: The Japan Dialysis Outcomes and Practice Pattern Study. <i>Nephron Clinical Practice</i> , 2012, 120, c91-c100.	2.3	20

#	ARTICLE	IF	CITATIONS
217	Gender Difference in Association Between Low Back Pain and Metabolic Syndrome. <i>Spine</i> , 2012, 37, 1130-1137.	1.0	19
218	Development and Validation of a Symptom Scale for Lumbar Spinal Stenosis. <i>Spine</i> , 2012, 37, 232-239.	1.0	10
219	Relationships of Decreased Lung Function with Metabolic Syndrome and Obstructive Sleep Apnea in Japanese Males. <i>Internal Medicine</i> , 2012, 51, 2291-2297.	0.3	14
220	Blood pressure levels and mortality risk among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study. <i>Kidney International</i> , 2012, 82, 570-580.	2.6	120
221	Long-term quality of life of donors after living donor liver transplantation. <i>Liver Transplantation</i> , 2012, 18, 1343-1352.	1.3	45
222	Relationship between health-related quality of life and clustering of metabolic syndrome diagnostic components. <i>Quality of Life Research</i> , 2012, 21, 1165-1170.	1.5	20
223	Sex differences in the change in health-related quality of life associated with low back pain. <i>Quality of Life Research</i> , 2012, 21, 1705-1711.	1.5	11
224	Locomotor dysfunction and risk of cardiovascular disease, quality of life, and medical costs: design of the Locomotive Syndrome and Health Outcome in Aizu Cohort Study (LOHAS) and baseline characteristics of the study population. <i>Journal of Orthopaedic Science</i> , 2012, 17, 261-271.	0.5	49
225	Erythropoiesis-Stimulating Agent Responsiveness and Mortality in Hemodialysis Patients: Results from a Cohort Study From the Dialysis Registry in Japan. <i>American Journal of Kidney Diseases</i> , 2012, 59, 108-116.	2.1	91
226	Opinion of Japanese rheumatology physicians on methods of assessing the quality of rheumatoid arthritis care. <i>Journal of Evaluation in Clinical Practice</i> , 2012, 18, 290-295.	0.9	2
227	Differences in relationships among sleep apnoea, glucose level, sleep duration and sleepiness between persons with and without type 2 diabetes. <i>Journal of Sleep Research</i> , 2012, 21, 410-418.	1.7	27
228	Understanding Measurements of Vitality in Patients with Chronic Kidney Disease: Connecting a Quality-of-Life Scale to Daily Activities. <i>PLoS ONE</i> , 2012, 7, e40455.	1.1	8
229	Validation testing of a three-component model of Short Form-36 scores. <i>Journal of Clinical Epidemiology</i> , 2011, 64, 301-308.	2.4	194
230	Application of a two-question screening instrument to detect depressive symptoms in patients with vitiligo: A pilot study. <i>Journal of the American Academy of Dermatology</i> , 2011, 64, e69-e70.	0.6	6
231	Post-discharge depressive symptoms can predict quality of life in AMI survivors: A prospective cohort study in Japan. <i>International Journal of Cardiology</i> , 2011, 146, 379-384.	0.8	5
232	Status of Provision of Drug Information to Hypertensive Patients at Community-based Pharmacies. <i>General Medicine</i> , 2011, 12, 75-82.	0.1	1
233	Development and Validation of a Quality of Life Scale Specific for Lumbar Spinal Stenosis. <i>Spine</i> , 2011, 36, E1407-E1414.	1.0	7
234	Management of asymptomatic hyperuricaemia in patients with chronic kidney disease by Japanese nephrologists: A questionnaire survey. <i>Nephrology</i> , 2011, 16, 518-521.	0.7	21

#	ARTICLE	IF	CITATIONS
235	Health-related quality of life among community-dwelling patients with intractable neurological diseases and their caregivers in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2011, 65, 30-38.	1.0	28
236	Effects of the presence of hypertension on the relationship between obstructive sleep apnoea and sleepiness. <i>Journal of Sleep Research</i> , 2011, 20, 538-543.	1.7	12
237	Mineral and Bone Disorders Outcomes Study for Japanese Chronic Kidney Disease Stage 5D Patients: Rationale and Study Design. <i>Therapeutic Apheresis and Dialysis</i> , 2011, 15, 169-175.	0.4	21
238	Positive Outcomes of High Hemoglobin Target in Patients With Chronic Kidney Disease Not on Dialysis: A Randomized Controlled Study. <i>Therapeutic Apheresis and Dialysis</i> , 2011, 15, 431-440.	0.4	42
239	Validation of the Japanese version of the quality of recovery score QoR-40. <i>Journal of Anesthesia</i> , 2011, 25, 509-515.	0.7	64
240	Reliability, validity, and responsiveness of the Japanese version of the Neck Pain and Disability Scale. <i>Journal of Orthopaedic Science</i> , 2011, 16, 339-346.	0.5	4
241	Darbepoetin alfa (KRN321) is safe and effective when administered subcutaneously once every 2 or 4 weeks to patients on peritoneal dialysis in Japan. <i>Clinical and Experimental Nephrology</i> , 2011, 15, 884-892.	0.7	8
242	Changes in anemia management and hemoglobin levels following revision of a bundling policy to incorporate recombinant human erythropoietin. <i>Kidney International</i> , 2011, 79, 340-346.	2.6	41
243	The Associations of Social Support and Other Psychosocial Factors with Mortality and Quality of Life in the Dialysis Outcomes and Practice Patterns Study. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011, 6, 142-152.	2.2	142
244	The relation between self-reported body weight and health-related quality of life: a cross-sectional study in Japan. <i>Journal of Public Health</i> , 2011, 33, 518-526.	1.0	14
245	Decreases in PTH in Japanese Hemodialysis Patients with Secondary Hyperparathyroidism. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011, 6, 2280-2288.	2.2	28
246	C-Reactive Protein and Mortality in Hemodialysis Patients: The Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>Nephron Clinical Practice</i> , 2011, 117, c167-c178.	2.3	57
247	Mineral Metabolism Management in Hemodialysis Patients with Secondary Hyperparathyroidism in Japan: Baseline Data from the MBD-5D. <i>American Journal of Nephrology</i> , 2011, 33, 427-437.	1.4	38
248	A simple four-grade score for aortic arch calcification by posteroanterior chest X-ray is associated with cardiovascular disease in haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 1747-1748.	0.4	2
249	Results of the survey on practice patterns including MBD management at dialysis facilities. <i>Nihon Toseki Igakkai Zasshi</i> , 2011, 44, 557-566.	0.2	3
250	Maintaining high hemoglobin levels improved the left ventricular mass index and quality of life scores in pre-dialysis Japanese chronic kidney disease patients. <i>Clinical and Experimental Nephrology</i> , 2010, 14, 28-35.	0.7	25
251	Comparison of dental practice income and expenses according to treatment types in the Japanese insurance system. <i>Japanese Dental Science Review</i> , 2010, 46, 4-10.	2.0	5
252	Keeping vulnerable elderly patients free from pressure ulcer is associated with high caregiver burden in informal caregivers. <i>Journal of Evaluation in Clinical Practice</i> , 2010, 16, 585-589.	0.9	16

#	ARTICLE	IF	CITATIONS
253	Can pediatricians accurately identify maternal depression at well-child visits?. <i>Pediatrics International</i> , 2010, 52, 284-289.	0.2	5
254	Associations Between Obstructive Sleep Apnea, Metabolic Syndrome, and Sleep Duration, As Measured With an Actigraph, in an Urban Male Working Population in Japan. <i>Sleep</i> , 2010, 33, 89-95.	0.6	67
255	Physical exercise among participants in the Dialysis Outcomes and Practice Patterns Study (DOPPS): correlates and associated outcomes. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 3050-3062.	0.4	251
256	Opinions of Japanese rheumatology physicians regarding clinical practice guidelines. <i>International Journal for Quality in Health Care</i> , 2010, 22, 78-85.	0.9	9
257	Cost-effectiveness of administering oral adsorbent AST-120 to patients with diabetes and advance-stage chronic kidney disease. <i>Diabetes Research and Clinical Practice</i> , 2010, 90, 154-159.	1.1	19
258	Association between number of comorbid conditions, depression, and sleep quality using the Pittsburgh Sleep Quality Index: Results from a population-based survey. <i>Sleep Medicine</i> , 2010, 11, 366-371.	0.8	112
259	Diabetes in Asia. <i>Lancet, The</i> , 2010, 375, 981-982.	6.3	1
260	Greater First-Year Survival on Hemodialysis in Facilities in Which Patients Are Provided Earlier and More Frequent Pre-nephrology Visits. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009, 4, 595-602.	2.2	88
261	Impact of Early Referral to Nephrologist on Mental Health among Hemodialysis Patients: A Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>Nephron Clinical Practice</i> , 2009, 113, c191-c197.	2.3	7
262	Association Between Frequency of Pruritic Symptoms and Perceived Psychological Stress. <i>Archives of Dermatology</i> , 2009, 145, 1384-8.	1.7	43
263	Effect of a Carbonaceous Oral Adsorbent on the Progression of CKD: A Multicenter, Randomized, Controlled Trial. <i>American Journal of Kidney Diseases</i> , 2009, 54, 459-467.	2.1	142
264	Violent Patient Behavior Is Associated with Bodily Pain and a High Burden on Informal Caregivers. <i>Journal of General Internal Medicine</i> , 2009, 24, 1085-1088.	1.3	12
265	Test performance of two-question screening for postpartum depressive symptoms. <i>Pediatrics International</i> , 2009, 51, 48-53.	0.2	24
266	Depressive Symptoms Predict the Subsequent Risk of Bodily Pain in Dialysis Patients: Japan Dialysis Outcomes and Practice Patterns Study. <i>Pain Medicine</i> , 2009, 10, 883-889.	0.9	18
267	Development of a Japanese version of the Epworth Sleepiness Scale (JESS) based on Item Response Theory. <i>Sleep Medicine</i> , 2009, 10, 556-565.	0.8	256
268	Estimating a preference-based index from the Japanese SF-36. <i>Journal of Clinical Epidemiology</i> , 2009, 62, 1323-1331.	2.4	71
269	Care burden and depression in caregivers caring for patients with intractable neurological diseases at home in Japan. <i>Journal of the Neurological Sciences</i> , 2009, 276, 148-152.	0.3	63
270	Systematic Review and Metaanalysis. <i>Chest</i> , 2009, 136, 1576-1585.	0.4	154

#	ARTICLE	IF	CITATIONS
271	Simple four-variable screening tool for identification of patients with sleep-disordered breathing. <i>Sleep</i> , 2009, 32, 939-48.	0.6	50
272	Dialysis staff encouragement and fluid control adherence in patients on hemodialysis. <i>Nephrology Nursing Journal</i> , 2009, 36, 289-97.	0.1	23
273	Quality of Life Improvements in Dialysis Patients Receiving Darbepoetin Alfa. <i>Therapeutic Apheresis and Dialysis</i> , 2008, 12, 72-77.	0.4	15
274	Acylated ghrelin level in patients with OSA before and after nasal CPAP treatment. <i>Respirology</i> , 2008, 13, 810-816.	1.3	46
275	Health-related quality of life in Japanese men with localized prostate cancer: Assessment with the SF-36. <i>International Journal of Urology</i> , 2008, 15, 524-528.	0.5	26
276	Relationships between working status and health or health care utilization among Japanese elderly. <i>Geriatrics and Gerontology International</i> , 2008, 8, 32-40.	0.7	19
277	Travel Time to Dialysis as a Predictor of Health-Related Quality of Life, Adherence, and Mortality: The Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>American Journal of Kidney Diseases</i> , 2008, 51, 641-650.	2.1	123
278	Plasma Thioredoxin, a Novel Oxidative Stress Marker, in Patients with Obstructive Sleep Apnea Before and After Nasal Continuous Positive Airway Pressure. <i>Antioxidants and Redox Signaling</i> , 2008, 10, 715-726.	2.5	66
279	Predictive value of self-reported patient information for the identification of lumbar spinal stenosis. <i>Family Practice</i> , 2008, 25, 237-244.	0.8	43
280	A Prospective Study of Passive Smoking and Risk of Diabetes in a Cohort of Workers. <i>Diabetes Care</i> , 2008, 31, 732-734.	4.3	68
281	Consistent Aspirin Use Associated with Improved Arteriovenous Fistula Survival among Incident Hemodialysis Patients in the Dialysis Outcomes and Practice Patterns Study. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008, 3, 1373-1378.	2.2	34
282	Prospective Evaluation of Selection Criteria for Active Surveillance in Japanese Patients with Stage T1cN0M0 Prostate Cancer. <i>Japanese Journal of Clinical Oncology</i> , 2008, 38, 122-128.	0.6	82
283	Japanese haemodialysis anaemia management practices and outcomes (1999-2006): results from the DOPPS. <i>Nephrology Dialysis Transplantation</i> , 2008, 23, 3643-3653.	0.4	65
284	Headache, Mental Health, and Use of Medical Resources: Health Diary Study in Japan. <i>Journal of Health Science</i> , 2008, 54, 30-36.	0.9	1
285	Sleep-Disordered Breathing in the Usual Lifestyle Setting as Detected with Home Monitoring in a Population of Working Men in Japan. <i>Sleep</i> , 2008, 31, 419-425.	0.6	99
286	Prospective Health Diary Study for New Onset Chest Symptoms in the Japanese General Population. <i>Internal Medicine</i> , 2008, 47, 25-31.	0.3	3
287	Higher Internality of Health Locus of Control Is Associated With the Use of Complementary and Alternative Medicine Providers Among Patients Seeking Care for Acute Low-back Pain. <i>Clinical Journal of Pain</i> , 2008, 24, 725-730.	0.8	20
288	Lifestyle and work predictors of fatigue in Japanese manufacturing workers. <i>Occupational Medicine</i> , 2007, 57, 262-269.	0.8	18

#	ARTICLE	IF	CITATIONS
289	Hypercalcaemia is associated with poor mental health in haemodialysis patients: results from Japan DOPPS. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 1658-1664.	0.4	27
290	Measuring health-related quality of life in patients with end-stage renal disease: why and how. <i>Nature Clinical Practice Nephrology</i> , 2007, 3, 352-353.	2.0	28
291	Lack of appetite in haemodialysis patients associations with patient characteristics, indicators of nutritional status and outcomes in the international DOPPS. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 3538-3546.	0.4	93
292	Effect of coaching on psychological adjustment in patients with spinocerebellar degeneration: a pilot study. <i>Clinical Rehabilitation</i> , 2007, 21, 987-996.	1.0	19
293	Sleep quality predicts quality of life and mortality risk in haemodialysis patients: Results from the Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>Nephrology Dialysis Transplantation</i> , 2007, 23, 998-1004.	0.4	243
294	Health Related Quality of Life in Japanese Men With Localized Prostate Cancer Treated With Current Multiple Modalities Assessed by a Newly Developed Japanese Version of the Expanded Prostate Cancer Index Composite. <i>Journal of Urology</i> , 2007, 177, 1856-1861.	0.2	53
295	ACCULTURATION AND END-OF-LIFE DECISION MAKING: COMPARISON OF JAPANESE AND JAPANESE-AMERICAN FOCUS GROUPS. <i>Bioethics</i> , 2007, 21, 251-262.	0.7	78
296	Association of mineral metabolism factors with all-cause and cardiovascular mortality in hemodialysis patients: The Japan dialysis outcomes and practice patterns study. <i>Hemodialysis International</i> , 2007, 11, 340-348.	0.4	100
297	Relation between sleep quality and quantity, quality of life, and risk of developing diabetes in healthy workers in Japan: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) Study. <i>BMC Public Health</i> , 2007, 7, 129.	1.2	112
298	Musculoskeletal pain in Japan: prospective health diary study. <i>Rheumatology International</i> , 2007, 28, 7-14.	1.5	9
299	Factors associated with health-related quality of life among hemodialysis patients in the DOPPS. <i>Quality of Life Research</i> , 2007, 16, 545-557.	1.5	119
300	The organization and financing of end-stage renal disease treatment in Japan. <i>International Journal of Health Care Finance and Economics</i> , 2007, 7, 217-231.	1.2	37
301	Development of a clinical diagnosis support tool to identify patients with lumbar spinal stenosis. <i>European Spine Journal</i> , 2007, 16, 1951-1957.	1.0	89
302	Gastrointestinal symptoms in a Japanese population: A health diary study. <i>World Journal of Gastroenterology</i> , 2007, 13, 572.	1.4	27
303	Japanese version of the Dermatology Life Quality Index: validity and reliability in patients with acne. <i>Health and Quality of Life Outcomes</i> , 2006, 4, 46.	1.0	68
304	Validation of the Burden Index of Caregivers (BIC), a multidimensional short care burden scale from Japan. <i>Health and Quality of Life Outcomes</i> , 2006, 4, 52.	1.0	50
305	Validation of the Japanese version of the Quality of Life-Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA). <i>Growth Hormone and IGF Research</i> , 2006, 16, 340-347.	0.5	9
306	A study of the burden of caring for patients with amyotrophic lateral sclerosis (MND) in Japan. <i>British Journal of Neuroscience Nursing</i> , 2006, 2, 38-43.	0.1	7

#	ARTICLE	IF	CITATIONS
307	Discrepancy Between Disability and the Severity of Low Back Pain: Demographic, Psychological, and Employment-Related Factors. <i>Spine</i> , 2006, 31, 931-939.	1.0	39
308	Linguistic Adaptation and Validation of the General Oral Health Assessment Index (GOHAI) in an Elderly Japanese Population. <i>Journal of Public Health Dentistry</i> , 2006, 66, 273-275.	0.5	103
309	Prognostic Factors for a Composite End-Point of Renal Outcomes in Patients With Chronic Kidney Disease. <i>Therapeutic Apheresis and Dialysis</i> , 2006, 10, 72-77.	0.4	2
310	Gender Differences in Effects of Physical Activity on Quality of Life and Resource Utilization. <i>Quality of Life Research</i> , 2006, 15, 537-546.	1.5	59
311	Health-Related Quality of Life Among Japanese Women With Iron-Deficiency Anemia. <i>Quality of Life Research</i> , 2006, 15, 1559-1563.	1.5	33
312	Families' and Physicians' Predictions of Dialysis Patients' Preferences Regarding Life-Sustaining Treatments in Japan. <i>American Journal of Kidney Diseases</i> , 2006, 47, 122-130.	2.1	56
313	Association between high voltage overhead transmission lines and mental health: A cross-sectional study. <i>Bioelectromagnetics</i> , 2006, 27, 473-478.	0.9	1
314	Psychological adjustment has a greater effect on health-related quality of life than on severity of disease in Parkinson's disease. <i>Movement Disorders</i> , 2006, 21, 761-766.	2.2	52
315	Health-Related Quality of Life of Predialysis Patients with Chronic Renal Failure. <i>Nephron Clinical Practice</i> , 2006, 105, c1-c8.	2.3	33
316	Associations between exposure to ambient photochemical oxidants and the vitality or mental health domain of the health related quality of life. <i>Journal of Epidemiology and Community Health</i> , 2006, 60, 173-179.	2.0	14
317	A Cross-sectional Study of Alcohol Drinking and Health-related Quality of Life among Male Workers in Japan. <i>Journal of Occupational Health</i> , 2005, 47, 496-503.	1.0	36
318	Factors associated with do not resuscitate orders and rates of withdrawal from hemodialysis in the international DOPPS. <i>Kidney International</i> , 2005, 68, 1282-1288.	2.6	47
319	Effects of low dialysate calcium concentration on health-related quality of life in hemodialysis patients. <i>Clinical and Experimental Nephrology</i> , 2005, 9, 153-157.	0.7	8
320	Study design and methods for a clinical trial of an oral carbonaceous adsorbent used to prevent the progression of chronic kidney disease (CAP-KD). <i>Clinical and Experimental Nephrology</i> , 2005, 9, 219-227.	0.7	7
321	Health-related quality of life of mothers of children with leukemia in Japan. <i>Quality of Life Research</i> , 2005, 14, 1079-1085.	1.5	66
322	Trust in One's Physician: The Role of Ethnic Match, Autonomy, Acculturation, and Religiosity Among Japanese and Japanese Americans. <i>Annals of Family Medicine</i> , 2005, 3, 339-347.	0.9	51
323	Living close to automobile traffic and quality of life in Japan: A population-based survey. <i>International Journal of Environmental Health Research</i> , 2005, 15, 1-9.	1.3	16
324	Association between ambient air pollution and health-related quality of life in Japan: Ecological study. <i>International Journal of Environmental Health Research</i> , 2005, 15, 383-391.	1.3	25

#	ARTICLE	IF	CITATIONS
325	Usefulness of five-item and three-item Mental Health Inventories to screen for depressive symptoms in the general population of Japan. <i>Health and Quality of Life Outcomes</i> , 2005, 3, 48.	1.0	209
326	Psychometric properties of the 25-item National Eye Institute Visual Function Questionnaire (NEI Vt) in Japanese. <i>Optometry</i> , 2004, 75, 163-168.	1.0	163
327	Results of the international DOPPS hemodialysis study in Japan. <i>Nihon Toseki Igakkai Zasshi</i> , 2004, 37, 1865-1873.	0.2	7
328	Relationships Between Diurnal Blood Pressure Variation, Physical Activity, and Health-Related QOL. <i>Clinical and Experimental Hypertension</i> , 2004, 26, 145-155.	0.5	14
329	Focus Group Interviews Examining Attitudes Towards Medical Research Among the Japanese: A Qualitative Study. <i>Bioethics</i> , 2004, 18, 448-470.	0.7	33
330	Screening for depression in hemodialysis patients: Associations with diagnosis, treatment, and outcomes in the DOPPS. <i>Kidney International</i> , 2004, 66, 2047-2053.	2.6	237
331	Health-related quality of life in the Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>American Journal of Kidney Diseases</i> , 2004, 44, 54-60.	2.1	87
332	Health-related quality of life associated with chronic conditions in eight countries: Results from the International Quality of Life Assessment (IQOLA) Project. <i>Quality of Life Research</i> , 2004, 13, 283-298.	1.5	623
333	Development and Testing of the Japanese Version of the Migraine-Specific Quality of Life Instrument. <i>Quality of Life Research</i> , 2004, 13, 1489-1493.	1.5	7
334	The influence of locus of control on preferences for information and decision making. <i>Patient Education and Counseling</i> , 2004, 55, 236-240.	1.0	57
335	Health-related quality of life in the Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>American Journal of Kidney Diseases</i> , 2004, 44, 54-60.	2.1	138
336	Response shift in perception of sleepiness in obstructive sleep apnea-hypopnea syndrome before and after treatment with nasal CPAP. <i>Sleep</i> , 2004, 27, 490-3.	0.6	24
337	Effects of intravenous ascorbic acid on erythropoiesis and quality of life in unselected hemodialysis patients. <i>Journal of Nephrology</i> , 2004, 17, 537-43.	0.9	15
338	Validation of the Japanese version of the Roland-Morris Disability Questionnaire. <i>Journal of Orthopaedic Science</i> , 2003, 8, 543-548.	0.5	117
339	Reliability, validity, and responsiveness of the Japanese version of the Inflammatory Bowel Disease Questionnaire. <i>Journal of Gastroenterology</i> , 2003, 38, 1138-1143.	2.3	34
340	Nonadherence in hemodialysis: Associations with mortality, hospitalization, and practice patterns in the DOPPS. <i>Kidney International</i> , 2003, 64, 254-262.	2.6	417
341	Health-related quality of life as a predictor of mortality and hospitalization: The Dialysis Outcomes and Practice Patterns Study (DOPPS). <i>Kidney International</i> , 2003, 64, 339-349.	2.6	665
342	Health-related quality of life among dialysis patients on three continents: The Dialysis Outcomes and Practice Patterns Study. <i>Kidney International</i> , 2003, 64, 1903-1910.	2.6	267

#	ARTICLE	IF	CITATIONS
343	Contributions of Clinical Epidemiologists and Medical Librarians to Developing Evidence-Based Clinical Practice Guidelines in Japan: A Case of the Treatment of Rheumatoid Arthritis. <i>General Medicine</i> , 2003, 4, 21-28.	0.1	3
344	Health-related quality of life of unrelated bone marrow donors in Japan. <i>Blood</i> , 2002, 99, 1995-2001.	0.6	58
345	The Development and Use of Quality-of-Life Measures to Evaluate Health Outcomes in Japan. <i>Pharmacoeconomics</i> , 2002, 20, 17-23.	1.7	10
346	Development of Japanese version of the UCLA Prostate Cancer Index: a pilot validation study. <i>International Journal of Clinical Oncology</i> , 2002, 7, 0306-0311.	1.0	73
347	Acculturation of attitudes toward end-of-life care. <i>Journal of General Internal Medicine</i> , 2002, 17, 531-539.	1.3	129
348	Attitudes of the Japanese public and doctors towards use of archived information and samples without informed consent: Preliminary findings based on focus group interviews. <i>BMC Medical Ethics</i> , 2002, 3, E1.	1.0	37
349	Health-related quality of life among dialysis patients in Seattle and Aichi. <i>American Journal of Kidney Diseases</i> , 2001, 37, 987-996.	2.1	13
350	Dialysis patients[rsquo] preferences regarding cardiopulmonary resuscitation and withdrawal of dialysis in Japan. <i>American Journal of Kidney Diseases</i> , 2001, 37, 1216-1222.	2.1	24
351	Relationship between Having a Home Doctor and Outpatient Utilization.. <i>Journal of Epidemiology</i> , 2001, 11, 160-169.	1.1	2
352	Japanese Attitudes toward Advance Care Planning for End-of-Life Decisions. <i>General Medicine</i> , 2001, 2, 3-10.	0.1	5
353	Use of Prescribed Drugs Among Older People in Japan: Association with Not Having a Regular Physician. <i>Journal of the American Geriatrics Society</i> , 1999, 47, 1425-1429.	1.3	24
354	Telemedicine in Neurosurgery Using International Digital Telephone Services Between Japan and Malaysia â€”Technical Noteâ€”. <i>Neurologia Medico-Chirurgica</i> , 1999, 39, 773-778.	1.0	14
355	HEALTH-RELATED QUALITY OF LIFE AMONG RENAL-TRANSPLANT RECIPIENTS IN JAPAN1. <i>Transplantation</i> , 1999, 68, 1331-1335.	0.5	24
356	Translating Health Status Questionnaires and Evaluating Their Quality. <i>Journal of Clinical Epidemiology</i> , 1998, 51, 913-923.	2.4	761
357	Cross-Cultural Comparisons of the Content of SF-36 Translations across 10 Countries. <i>Journal of Clinical Epidemiology</i> , 1998, 51, 925-932.	2.4	250
358	Testing the Equivalence of Translations of Widely Used Response Choice Labels. <i>Journal of Clinical Epidemiology</i> , 1998, 51, 933-944.	2.4	90
359	Translation, Adaptation, and Validation of the SF-36 Health Survey for Use in Japan. <i>Journal of Clinical Epidemiology</i> , 1998, 51, 1037-1044.	2.4	1,146
360	Psychometric and Clinical Tests of Validity of the Japanese SF-36 Health Survey. <i>Journal of Clinical Epidemiology</i> , 1998, 51, 1045-1053.	2.4	813

#	ARTICLE	IF	CITATIONS
361	Tests of Data Quality, Scaling Assumptions, and Reliability of the SF-36 in Eleven Countries. Journal of Clinical Epidemiology, 1998, 51, 1149-1158.	2.4	359
362	Japanese physicians' views of ethical problems in dialysis therapy and advance directives.. Nihon Toseki Igakkai Zasshi, 1998, 31, 1221-1225.	0.2	0
363	Measurement of Information Seeking Preference of Japanese in Clinical Settings. The Japanese Journal of Health Psychology, 1997, 10, 1-10.	0.1	5