

# Christos Papamichael

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/11557067/christos-papamichael-publications-by-year.pdf>

**Version:** 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

49  
papers

2,320  
citations

26  
h-index

48  
g-index

50  
ext. papers

2,499  
ext. citations

3.9  
avg, IF

3.95  
L-index

#	Paper	IF	Citations
49	Reactive Vasodilation Predicts Mortality in Primary Systemic Light-Chain Amyloidosis. <i>Circulation Research</i> , <b>2019</b> , 125, 744-758	15.7	18
48	The effect of hypohydration on endothelial function in young healthy adults. <i>European Journal of Nutrition</i> , <b>2017</b> , 56, 1211-1217	5.2	27
47	Differential associations of systolic and diastolic time rate of blood pressure variation with carotid atherosclerosis and plaque echogenicity. <i>Journal of Clinical Hypertension</i> , <b>2017</b> , 19, 1070-1077	2.3	6
46	Eating frequency predicts new onset hypertension and the rate of progression of blood pressure, arterial stiffness, and wave reflections. <i>Journal of Hypertension</i> , <b>2016</b> , 34, 429-37; discussion 437	1.9	4
45	Free androgen index as a predictor of blood pressure progression and accelerated vascular aging in menopause. <i>Atherosclerosis</i> , <b>2016</b> , 247, 177-83	3.1	23
44	Meal patterns in healthy adults: Inverse association of eating frequency with subclinical atherosclerosis indexes. <i>Clinical Nutrition</i> , <b>2015</b> , 34, 302-8	5.9	10
43	Circulating levels of TNF-like cytokine 1A correlate with reflected waves and atherosclerosis extent and may predict cardiac death in patients with stable coronary artery disease. <i>Cytokine</i> , <b>2015</b> , 72, 102-4	4	5
42	Hemodynamic markers and subclinical atherosclerosis in postmenopausal women with primary hyperparathyroidism. <i>Journal of Clinical Endocrinology and Metabolism</i> , <b>2014</b> , 99, 2704-11	5.6	16
41	Subclinical atherosclerosis in menopausal women with low to medium calculated cardiovascular risk. <i>International Journal of Cardiology</i> , <b>2013</b> , 164, 70-6	3.2	30
40	Arterial stiffness is increased in asymptomatic nondiabetic postmenopausal women with a polycystic ovary syndrome phenotype. <i>Journal of Hypertension</i> , <b>2013</b> , 31, 1998-2004	1.9	26
39	Sesame oil consumption exerts a beneficial effect on endothelial function in hypertensive men. <i>European Journal of Preventive Cardiology</i> , <b>2013</b> , 20, 202-8	3.9	17
38	Acute and long-term hemodynamic effects of sesame oil consumption in hypertensive men. <i>Journal of Clinical Hypertension</i> , <b>2012</b> , 14, 630-6	2.3	11
37	Circulating androgen levels are associated with subclinical atherosclerosis and arterial stiffness in healthy recently menopausal women. <i>Metabolism: Clinical and Experimental</i> , <b>2012</b> , 61, 193-201	12.7	65
36	Recently postmenopausal women have the same prevalence of subclinical carotid atherosclerosis as age and traditional risk factor matched men. <i>Atherosclerosis</i> , <b>2012</b> , 221, 508-13	3.1	20
35	Impact of prehypertension on carotid artery intima-media thickening: actual or masked?. <i>Atherosclerosis</i> , <b>2011</b> , 214, 215-9	3.1	16
34	Association of the SHBG gene promoter polymorphism with early markers of atherosclerosis in apparently healthy women. <i>Atherosclerosis</i> , <b>2011</b> , 219, 205-10	3.1	9
33	Homa-IR: A marker of vascular dysfunction in nondiabetic hemodialysis patients?. <i>Blood Purification</i> , <b>2010</b> , 29, 327-8	3.1	1

32	Endogenous estrogen levels are associated with endothelial function in males independently of lipid levels. <i>Endocrine</i> , <b>2010</b> , 37, 329-35	4	17
31	Prolactin and preclinical atherosclerosis in menopausal women with cardiovascular risk factors. <i>Hypertension</i> , <b>2009</b> , 54, 98-105	8.5	82
30	Impact of prehypertension on common carotid artery intima-media thickness and left ventricular mass. <i>Stroke</i> , <b>2009</b> , 40, 1515-8	6.7	57
29	Free thyroxine is an independent predictor of subcutaneous fat in euthyroid individuals. <i>European Journal of Endocrinology</i> , <b>2009</b> , 161, 459-65	6.5	63
28	Circulating levels of TNF-like cytokine 1A (TL1A) and its decoy receptor 3 (DcR3) in rheumatoid arthritis. <i>Clinical Immunology</i> , <b>2008</b> , 129, 249-55	9	90
27	Topography and severity of coronary artery disease in white-coat hypertension. <i>European Journal of Internal Medicine</i> , <b>2008</b> , 19, 280-4	3.9	6
26	Postprandial improvement of endothelial function by red wine and olive oil antioxidants: a synergistic effect of components of the Mediterranean diet. <i>Journal of the American College of Nutrition</i> , <b>2008</b> , 27, 448-53	3.5	47
25	Correlations of sialic acid with markers of inflammation, atherosclerosis and cardiovascular events in hemodialysis patients. <i>Blood Purification</i> , <b>2008</b> , 26, 261-6	3.1	12
24	Treatment with granulocyte colony stimulating factor is associated with improvement in endothelial function. <i>Growth Factors</i> , <b>2008</b> , 26, 117-24	1.6	17
23	Target organ damage in "white coat hypertension" and "masked hypertension". <i>American Journal of Hypertension</i> , <b>2008</b> , 21, 393-9	2.3	98
22	Association of thyroid function with arterial pressure in normotensive and hypertensive euthyroid individuals: A cross-sectional study. <i>Thyroid Research</i> , <b>2008</b> , 1, 3	2.4	19
21	Acute smoking induces endothelial dysfunction in healthy smokers. Is this reversible by red wine/ antioxidant constituents?. <i>Journal of the American College of Nutrition</i> , <b>2007</b> , 26, 10-5	3.5	35
20	Acute smoke-induced endothelial dysfunction is more prolonged in smokers than in non-smokers. <i>International Journal of Cardiology</i> , <b>2007</b> , 120, 404-6	3.2	19
19	Common carotid artery intima-media thickness and the risk of stroke recurrence. <i>Stroke</i> , <b>2006</b> , 37, 1913-6.7		85
18	Divergent effects of rofecoxib on endothelial function and inflammation in acute coronary syndromes. <i>International Journal of Cardiology</i> , <b>2006</b> , 112, 359-66	3.2	7
17	Genetic variations of the endothelial nitric oxide synthase gene are related to increased levels of C-reactive protein and macrophage-colony stimulating-factor in patients with coronary artery disease. <i>Thrombosis and Haemostasis</i> , <b>2006</b> , 96, 520-528	7	8
16	Ambulatory blood pressure monitoring and target organ damage: effects of age and sex. <i>Blood Pressure Monitoring</i> , <b>2006</b> , 11, 9-15	1.3	24
15	Combined acute effects of red wine consumption and cigarette smoking on haemodynamics of young smokers. <i>Journal of Hypertension</i> , <b>2006</b> , 24, 1287-92	1.9	28

14	Adolescent obesity is associated with high ambulatory blood pressure and increased carotid intimal-medial thickness. <i>Journal of Pediatrics</i> , <b>2005</b> , 147, 651-6	3.6	146
13	Acute effects of caffeine on blood pressure and wave reflections in healthy subjects: should we consider monitoring central blood pressure?. <i>International Journal of Cardiology</i> , <b>2005</b> , 98, 425-30	3.2	74
12	Rapid effect of pravastatin on endothelial function and lipid peroxidation in unstable angina. <i>International Journal of Cardiology</i> , <b>2005</b> , 101, 65-70	3.2	33
11	White-coat and masked hypertension in children: association with target-organ damage. <i>Pediatric Nephrology</i> , <b>2005</b> , 20, 1151-5	3.2	191
10	Intima-media thickness score from carotid and femoral arteries predicts the extent of coronary artery disease: intima-media thickness and CAD. <i>International Journal of Cardiovascular Imaging</i> , <b>2005</b> , 21, 495-501	2.5	46
9	Time rate of blood pressure variation is associated with increased common carotid artery intima-media thickness. <i>Hypertension</i> , <b>2005</b> , 45, 505-12	8.5	171
8	Red wine antioxidants counteract acute endothelial dysfunction caused by cigarette smoking in healthy nonsmokers. <i>American Heart Journal</i> , <b>2004</b> , 147, 274	4.9	54
7	Constituents of red wine other than alcohol improve endothelial function in patients with coronary artery disease. <i>Coronary Artery Disease</i> , <b>2004</b> , 15, 485-90	1.4	68
6	Isolated clinic hypertension is not an innocent phenomenon: effect on the carotid artery structure. <i>American Journal of Hypertension</i> , <b>1999</b> , 12, 245-50	2.3	40
5	Effects of acute cigarette smoking on endothelium-dependent arterial dilatation in normal subjects. <i>American Journal of Cardiology</i> , <b>1998</b> , 81, 1225-8	3	86
4	Effect of long-term estrogen therapy on brachial arterial endothelium-dependent vasodilation in women with Raynaud phenomenon secondary to systemic sclerosis. <i>American Journal of Cardiology</i> , <b>1998</b> , 82, 1555-7, A8	3	46
3	Short-term estrogen administration improves abnormal endothelial function in women with systemic sclerosis and Raynaud phenomenon. <i>American Heart Journal</i> , <b>1998</b> , 136, 905-12	4.9	67
2	Flow-mediated, endothelium-dependent vasodilation is impaired in subjects with hypothyroidism, borderline hypothyroidism, and high-normal serum thyrotropin (TSH) values. <i>Thyroid</i> , <b>1997</b> , 7, 411-4	6.2	190
1	Effect of acute cigarette smoking on endothelium-dependent brachial artery dilatation in healthy individuals. <i>American Journal of Cardiology</i> , <b>1997</b> , 79, 529-31	3	90