

Yuichi Inoue

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1156060/yuichi-inoue-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

192
papers

4,313
citations

34
h-index

56
g-index

202
ext. papers

5,248
ext. citations

3.4
avg, IF

5.49
L-index

#	Paper	IF	Citations
192	Associations Between the 2011 Great East Japan Earthquake and Tsunami and the Sleep and Mental Health of Japanese People: A 3-Wave Repeated Survey.. <i>Nature and Science of Sleep</i> , 2022 , 14, 61-73	3.6	0
191	Nightmares in People with COVID-19: Did Coronavirus Infect Our Dreams?. <i>Nature and Science of Sleep</i> , 2022 , 14, 93-108	3.6	6
190	Changes in sleep behavior, sleep problems, and psychological distress/health-related quality of life of young Japanese individuals before and during the COVID-19 pandemic.. <i>Chronobiology International</i> , 2022 , 1-11	3.6	0
189	Sleep Apnea Syndrome (SAS) Clinical Practice Guidelines 2020. <i>Sleep and Biological Rhythms</i> , 2022 , 20, 5	1.3	0
188	Sleep Apnea Syndrome (SAS) Clinical Practice Guidelines 2020.. <i>Respiratory Investigation</i> , 2022 , 60, 3-32	3.4	0
187	Optimal dose determination of enerisant (TS-091) for patients with narcolepsy: two randomized, double-blind, placebo-controlled trials.. <i>BMC Psychiatry</i> , 2022 , 22, 141	4.2	0
186	Efficacy and safety of lemborexant over 12 months in Asian adults with insomnia disorder.. <i>Sleep Medicine: X</i> , 2022 , 4, 100044	3.1	
185	A rare genetic variant in the cleavage site of prepro-orexin is associated with idiopathic hypersomnia.. <i>Npj Genomic Medicine</i> , 2022 , 7, 29	6.2	0
184	Dream-enactment behaviours during the COVID-19 pandemic: an international COVID-19 sleep study.. <i>Journal of Sleep Research</i> , 2022 ,	5.8	1
183	Metabolic-Syndrome-Related Comorbidities in Narcolepsy Spectrum Disorders: A Preliminary Cross-Sectional Study in Japan. <i>International Journal of Environmental Research and Public Health</i> , 2022 , 19, 6285	4.6	0
182	Disturbances in sleep, circadian rhythms and daytime functioning in relation to coronavirus infection and Long-COVID - A multinational ICOS study.. <i>Journal of Sleep Research</i> , 2021 , e13542	5.8	4
181	Sleep-wake rhythm and its association with lifestyle, health-related quality of life and academic performance among Japanese nursing students: a cross-sectional study. <i>BMC Nursing</i> , 2021 , 20, 225	3.2	0
180	Social jetlag among Japanese adolescents: Association with irritable mood, daytime sleepiness, fatigue, and poor academic performance. <i>Chronobiology International</i> , 2021 , 1-12	3.6	3
179	Social Jetlag Changes During the COVID-19 Pandemic as a Predictor of Insomnia - A Multi-National Survey Study. <i>Nature and Science of Sleep</i> , 2021 , 13, 1711-1722	3.6	10
178	Sleep Debt and Social Jetlag Associated with Sleepiness, Mood, and Work Performance among Workers in Japan. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	5
177	Efficacy and safety of modafinil in patients with idiopathic hypersomnia without long sleep time: a multicenter, randomized, double-blind, placebo-controlled, parallel-group comparison study. <i>Sleep Medicine</i> , 2021 , 80, 315-321	4.6	3
176	The association between high risk of sleep apnea, comorbidities, and risk of COVID-19: a population-based international harmonized study. <i>Sleep and Breathing</i> , 2021 , 25, 849-860	3.1	13

175	Long-term effectiveness and safety of lemborexant in adults with insomnia disorder: results from a phase 3 randomized clinical trial. <i>Sleep Medicine</i> , 2021 , 80, 333-342	4.6	13
174	Prevalence, clinical symptoms and polysomnographic findings of REM-related sleep disordered breathing in Japanese population. <i>Sleep Medicine</i> , 2021 , 80, 52-56	4.6	1
173	Prevalence of and factors associated with acute withdrawal symptoms after 24 weeks of eszopiclone treatment in patients with chronic insomnia: a prospective, interventional study. <i>BMC Psychiatry</i> , 2021 , 21, 193	4.2	3
172	Efficacy and safety of lemborexant in adults with insomnia: comparing Japanese and non-Japanese subgroups from the global, phase 3, randomized, double-blind, placebo-controlled SUNRISE 2 study. <i>Journal of Clinical Sleep Medicine</i> , 2021 , 17, 1067-1074	3.1	2
171	Reliability and validity of the Japanese version of the Biological Rhythms Interview of assessment in neuropsychiatry-self report for delayed sleep-wake phase disorder. <i>Sleep Medicine</i> , 2021 , 81, 288-293	4.6	2
170	Association of sleep duration on workdays or free days and social jetlag with job stress. <i>Psychiatry and Clinical Neurosciences</i> , 2021 , 75, 244-249	6.2	4
169	Development and validation of the Benzodiazepine Hypnotics Withdrawal Symptom Scale (BHWSS) based on item response theory. <i>Psychiatry Research</i> , 2021 , 300, 113900	9.9	1
168	The relationship between insomnia symptoms and work productivity among blue-collar and white-collar Japanese workers engaged in construction/civil engineering work: a cross-sectional study. <i>BMC Public Health</i> , 2021 , 21, 1244	4.1	0
167	Current state of hypnotic use disorders: Results of a survey using the Japanese version of Benzodiazepine Dependence Self-Report Questionnaire. <i>Neuropsychopharmacology Reports</i> , 2021 , 41, 14-25	2.2	1
166	Prevalence and associated factors of circadian rhythm sleep-wake disorders and insomnia among visually impaired Japanese individuals. <i>BMC Public Health</i> , 2021 , 21, 31	4.1	0
165	Heart rate change and clinical characteristics in patients with neck myoclonus: An observational study. <i>Clinical Neurophysiology Practice</i> , 2021 , 6, 229-233	3.8	
164	The efficacy of add-on ramelteon and subsequent dose reduction in benzodiazepine derivatives/Z-drugs for the treatment of sleep-related eating disorder and night eating syndrome: a retrospective analysis of consecutive patients. <i>Journal of Clinical Sleep Medicine</i> , 2021 , 17, 1475-1483	3.1	1
163	Evening-types show highest increase of sleep and mental health problems during the COVID-19 pandemic - Multinational study on 19,267 adults. <i>Sleep</i> , 2021 ,	1.1	10
162	We need to do better: A systematic review and meta-analysis of diagnostic test accuracy of restless legs syndrome screening instruments. <i>Sleep Medicine Reviews</i> , 2021 , 58, 101461	10.2	4
161	Difference in background factors between responders to gabapentin enacarbil treatment and responders to placebo: pooled analyses of two randomized, double-blind, placebo-controlled studies in Japanese patients with restless legs syndrome. <i>Sleep Medicine</i> , 2021 , 85, 138-146	4.6	0
160	How our Dreams Changed During the COVID-19 Pandemic: Effects and Correlates of Dream Recall Frequency - a Multinational Study on 19,355 Adults. <i>Nature and Science of Sleep</i> , 2021 , 13, 1573-1591	3.6	14
159	Sleep and daytime problems during the COVID-19 pandemic and effects of coronavirus infection, confinement and financial suffering: a multinational survey using a harmonised questionnaire.. <i>BMJ Open</i> , 2021 , 11, e050672	3	11
158	The effect of short or long sleep duration on quality of life and depression: an internet-based survey in Japan. <i>Sleep Medicine</i> , 2020 , 76, 80-85	4.6	5

157	Prevalence and Associated Factors of Nocturnal Eating Behavior and Sleep-Related Eating Disorder-Like Behavior in Japanese Young Adults: Results of an Internet Survey Using Munich Parasomnia Screening. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	6
156	Impact of sleep problems on daytime function in school life: a cross-sectional study involving Japanese university students. <i>BMC Public Health</i> , 2020 , 20, 371	4.1	8
155	Prevalence and clinical characteristics of REM sleep behavior disorder in Japanese elderly people. <i>Sleep</i> , 2020 , 43,	1.1	10
154	Long-term efficacy and tolerability of lemborexant compared with placebo in adults with insomnia disorder: results from the phase 3 randomized clinical trial SUNRISE 2. <i>Sleep</i> , 2020 , 43,	1.1	36
153	Tracking intermediate performance of vigilant attention using multiple eye metrics. <i>Sleep</i> , 2020 , 43,	1.1	4
152	Gender differences in clinical findings and ßynucleiopathy-related markers in patients with idiopathic REM sleep behavior disorder. <i>Sleep Medicine</i> , 2020 , 66, 216-219	4.6	5
151	A missense variant in PER2 is associated with delayed sleep-wake phase disorder in a Japanese population. <i>Journal of Human Genetics</i> , 2019 , 64, 1219-1225	4.3	11
150	Correlation among clock gene expression rhythms, sleep quality, and meal conditions in delayed sleep-wake phase disorder and night eating syndrome. <i>Chronobiology International</i> , 2019 , 36, 770-783	3.6	3
149	A survey on social jetlag in Japan: a nationwide, cross-sectional internet survey. <i>Sleep and Biological Rhythms</i> , 2019 , 17, 417-422	1.3	14
148	An autopsy case of MM2-thalamic subtype of sporadic Creutzfeldt-Jakob disease with Lewy bodies presenting as a sleep disorder mimicking anti-IgLON5 disease. <i>Journal of the Neurological Sciences</i> , 2019 , 404, 36-39	3.2	4
147	Rotigotine suppresses sleep-related muscle activity augmented by injection of dialysis patientsQ sera in a mouse model of restless legs syndrome. <i>Scientific Reports</i> , 2019 , 9, 16344	4.9	2
146	Effect of Yokukansan for the Treatment of Idiopathic Rapid Eye Movement Sleep Behavior Disorder: A Retrospective Analysis of Consecutive Patients. <i>Journal of Clinical Sleep Medicine</i> , 2019 , 15, 1173-1178	3.1	5
145	Discrepancy in wake-up time on school days and free days is associated with daytime sleepiness, lowered mental/physical health and poor academic performance. <i>Shinrigaku Kenkyu</i> , 2019 , 90, 378-388	0.6	2
144	Development of a new benzodiazepine hypnotics withdrawal symptom scale. <i>Sleep and Biological Rhythms</i> , 2018 , 16, 263-271	1.3	5
143	Efficacy of cognitive behavioral therapy for comorbid insomnia: a meta-analysis. <i>Sleep and Biological Rhythms</i> , 2018 , 16, 21-35	1.3	9
142	Clinical significance of social jetlag in patients with excessive daytime sleepiness. <i>Chronobiology International</i> , 2018 , 35, 1637-1646	3.6	5
141	Effectiveness of cognitive behavioral therapy for pharmacotherapy-resistant chronic insomnia: a multi-center randomized controlled trial in Japan. <i>Sleep Medicine</i> , 2018 , 50, 105-112	4.6	9
140	Treatment of restless legs syndrome: Evidence-based review and implications for clinical practice (Revised 2017). <i>Movement Disorders</i> , 2018 , 33, 1077-1091	7	100

139	Ethnic differences in periodic limb movements during sleep in patients with restless legs syndrome: a preliminary cross-sectional study of Austrian and Japanese clinical population. <i>Sleep and Biological Rhythms</i> , 2018 , 16, 345-349	1.3	4
138	Circadian Rhythm Sleep-Wake Disorders Predict Shorter Time to Relapse of Mood Episodes in Euthymic Patients With Bipolar Disorder: A Prospective 48-Week Study. <i>Journal of Clinical Psychiatry</i> , 2018 , 79,	4.6	25
137	A variant at 9q34.11 is associated with HLA-DQB1*06:02[negative essential hypersomnia. <i>Journal of Human Genetics</i> , 2018 , 63, 1259-1267	4.3	3
136	Lack of association between variable number tandem repeat and circadian rhythm sleep-wake disorders. <i>Human Genome Variation</i> , 2018 , 5, 17	1.8	8
135	Excessive daytime sleepiness in adults with possible attention deficit/hyperactivity disorder (ADHD): a web-based cross-sectional study. <i>Sleep Medicine</i> , 2017 , 32, 4-9	4.6	17
134	Association among changes in sleep-related beliefs, sleep reactivity, and improvement of insomnia following cognitive behavioral therapy. <i>Sleep Medicine</i> , 2017 , 29, 96-97	4.6	10
133	Pareidolias in REM Sleep Behavior Disorder: A Possible Predictive Marker of Lewy Body Diseases?. <i>Sleep</i> , 2017 , 40,	1.1	17
132	Circadian rhythm sleep-wake disorders as predictors for bipolar disorder in patients with remitted mood disorders. <i>Journal of Affective Disorders</i> , 2017 , 220, 57-61	6.6	23
131	Insufficient sleep rather than the apnea-hypopnea index can be associated with sleepiness-related driving problems of Japanese obstructive sleep apnea syndrome patients residing in metropolitan areas. <i>Sleep Medicine</i> , 2017 , 33, 19-22	4.6	6
130	Higher sleep reactivity and insomnia mutually aggravate depressive symptoms: a cross-sectional epidemiological study in Japan. <i>Sleep Medicine</i> , 2017 , 33, 130-133	4.6	5
129	Effects of acute morning and evening exercise on subjective and[object]jective sleep quality in older individuals with insomnia. <i>Sleep Medicine</i> , 2017 , 34, 200-208	4.6	23
128	Validation of a sheet-shaped body vibrometer for screening of obstructive sleep apnea. <i>Drug Discoveries and Therapeutics</i> , 2017 , 11, 126-132	5	6
127	Achievements, challenges, and future perspectives of epidemiologic research in restless legs syndrome (RLS). <i>Sleep Medicine</i> , 2017 , 31, 3-9	4.6	26
126	Relationship between Self-Reported Dietary Nutrient Intake and Self-Reported Sleep Duration among Japanese Adults. <i>Nutrients</i> , 2017 , 9,	6.7	31
125	Prevalence, symptomatic features, and factors associated with sleep disturbance/insomnia in Japanese patients with type-2 diabetes. <i>Neuropsychiatric Disease and Treatment</i> , 2017 , 13, 1873-1880	3.1	25
124	Restless legs syndrome augmentation among Japanese patients receiving pramipexole therapy: Rate and risk factors in a retrospective study. <i>PLoS ONE</i> , 2017 , 12, e0173535	3.7	5
123	Social jetlag affects subjective daytime sleepiness in school-aged children and adolescents: A study using the Japanese version of the Pediatric Daytime Sleepiness Scale (PDSS-J). <i>Chronobiology International</i> , 2016 , 33, 1311-1319	3.6	34
122	Progression of obstructive sleep apnoea syndrome in Japanese patients. <i>Sleep and Breathing</i> , 2016 , 20, 711-8	3.1	1

121	Later sleep schedule and depressive symptoms are associated with usage of multiple kinds of hypnotics. <i>Sleep Medicine</i> , 2016 , 25, 56-62	4.6	2
120	Oropharyngeal Crowding Closely Relates to Aggravation of OSA. <i>Chest</i> , 2016 , 150, 346-52	5.3	12
119	Factors associated with discontinuation of aripiprazole treatment after switching from other antipsychotics in patients with chronic schizophrenia: A prospective observational study. <i>Psychiatry Research</i> , 2016 , 236, 71-74	9.9	10
118	Guidelines for the first-line treatment of restless legs syndrome/Willis-Ekbom disease, prevention and treatment of dopaminergic augmentation: a combined task force of the IRLSSG, EURLSSG, and the RLS-Foundation. <i>Sleep Medicine</i> , 2016 , 21, 1-11	4.6	172
117	Prevalence of and Factors Associated With Sleep-Related Eating Disorder in Psychiatric Outpatients Taking Hypnotics. <i>Journal of Clinical Psychiatry</i> , 2016 , 77, e892-8	4.6	4
116	A Nationwide Cross-Sectional Survey of Sleep-Related Problems in Japanese Visually Impaired Patients: Prevalence and Association with Health-Related Quality of Life. <i>Journal of Clinical Sleep Medicine</i> , 2016 , 12, 1659-1667	3.1	17
115	Comparison of clinical features between primary and drug-induced sleep-related eating disorder. <i>Neuropsychiatric Disease and Treatment</i> , 2016 , 12, 1275-80	3.1	6
114	Prevalence of Circadian Rhythm Sleep-Wake Disorders and Associated Factors in Euthymic Patients with Bipolar Disorder. <i>PLoS ONE</i> , 2016 , 11, e0159578	3.7	33
113	Association between the high-dose use of benzodiazepines and rehospitalization in patients with schizophrenia: a 2-year naturalistic study. <i>Neuropsychiatric Disease and Treatment</i> , 2016 , 12, 3243-3247	3.1	2
112	Findings of the Maintenance of Wakefulness Test and its relationship with response to modafinil therapy for residual excessive daytime sleepiness in obstructive sleep apnea patients adequately treated with nasal continuous positive airway pressure. <i>Sleep Medicine</i> , 2016 , 27-28, 45-48	4.6	3
111	Asian specific feature of sleep habits and sleep disorders. <i>Sleep and Biological Rhythms</i> , 2016 , 14, 123-124	4.3	5
110	Sleep patterns among shift-working flight controllers of the International Space Station: an observational study on the JAXA Flight Control Team. <i>Journal of Physiological Anthropology</i> , 2016 , 35, 19	2.5	11
109	Restless legs syndrome-current therapies and management of augmentation. <i>Nature Reviews Neurology</i> , 2015 , 11, 434-45	15	60
108	Impact of hypnotics use on daytime function and factors associated with usage by female shift work nurses. <i>Sleep Medicine</i> , 2015 , 16, 604-11	4.6	11
107	A preliminary study on the relationships between diurnal melatonin secretion profile and sleep variables in patients emergently admitted to the coronary care unit. <i>Chronobiology International</i> , 2015 , 32, 875-9	3.6	12
106	Restless legs syndrome in hemodialysis patients: Prevalence and association to daytime functioning. <i>Sleep and Biological Rhythms</i> , 2015 , 13, 127-135	1.3	1
105	Differences in electroencephalographic findings among categories of narcolepsy-spectrum disorders. <i>Sleep Medicine</i> , 2015 , 16, 999-1005	4.6	8
104	Sleepiness at the Wheel and Countermeasures: Effects of Caffeine, Napping, and Blue Light 2015 , 271-277		

103	A polymorphism in CCR1/CCR3 is associated with narcolepsy. <i>Brain, Behavior, and Immunity</i> , 2015 , 49, 148-55	16.6	30
102	Daytime symptoms of restless legs syndrome--clinical characteristics and rotigotine effectiveness. <i>Sleep Medicine</i> , 2015 , 16, 871-6	4.6	8
101	New susceptibility variants to narcolepsy identified in HLA class II region. <i>Human Molecular Genetics</i> , 2015 , 24, 891-8	5.6	20
100	Internet-based survey of factors associated with subjective feeling of insomnia, depression, and low health-related quality of life among Japanese adults with sleep difficulty. <i>International Journal of Behavioral Medicine</i> , 2015 , 22, 233-8	2.6	6
99	Prevalence and associated factors of hypnotics dependence among Japanese outpatients with psychiatric disorders. <i>Psychiatry Research</i> , 2015 , 230, 958-63	9.9	9
98	Sleep-related eating disorder and its associated conditions. <i>Psychiatry and Clinical Neurosciences</i> , 2015 , 69, 309-20	6.2	29
97	An association analysis of HLA-DQB1 with narcolepsy without cataplexy and idiopathic hypersomnia with/without long sleep time in a Japanese population. <i>Human Genome Variation</i> , 2015 , 2, 15031	1.8	16
96	Polysomnographic Assessment of Sleep Comorbidities in Drug-Naïve Narcolepsy-Spectrum Disorders--A Japanese Cross-Sectional Study. <i>PLoS ONE</i> , 2015 , 10, e0136988	3.7	8
95	The Prevalence and Characteristics of Primary Headache and Dream-Enacting Behaviour in Japanese Patients with Narcolepsy or Idiopathic Hypersomnia: A Multi-Centre Cross-Sectional Study. <i>PLoS ONE</i> , 2015 , 10, e0139229	3.7	18
94	Japanese version of the Munich Parasomnia Screening: translation and linguistic validation of a screening instrument for parasomnias and nocturnal behaviors. <i>Neuropsychiatric Disease and Treatment</i> , 2015 , 11, 2953-8	3.1	2
93	Patient Background Factors Affecting the Therapeutic Outcomes in Response to Eszopiclone in Adult Patients with Chronic Insomnia: A Post Hoc Analysis of a Double-Blind Phase III Study in Japan. <i>Journal of Clinical Sleep Medicine</i> , 2015 , 11, 1171-8	3.1	1
92	Prevalence and Correlates of Insufficient Sleep Syndrome in Japanese Young Adults: A Web-Based Cross-Sectional Study. <i>Journal of Clinical Sleep Medicine</i> , 2015 , 11, 1163-9	3.1	12
91	The impact of a delayed sleep-wake schedule on depression is greater in women--A web-based cross-sectional study in Japanese young adults. <i>Chronobiology International</i> , 2015 , 32, 952-8	3.6	13
90	Screening of clock gene polymorphisms demonstrates association of a PER3 polymorphism with morningness-eveningness preference and circadian rhythm sleep disorder. <i>Scientific Reports</i> , 2014 , 4, 6309	4.9	82
89	Is maxillary dental arch constriction common in Japanese male adult patients with obstructive sleep apnoea?. <i>European Journal of Orthodontics</i> , 2014 , 36, 403-8	3.3	7
88	Effect of delayed sleep phase during university life on the daytime functioning in work life after graduation. <i>Sleep Medicine</i> , 2014 , 15, 1155-8	4.6	8
87	The validity of the PAM-RL device for evaluating periodic limb movements in sleep and an investigation on night-to-night variability of periodic limb movements during sleep in patients with restless legs syndrome or periodic limb movement disorder using this system. <i>Sleep Medicine</i> , 2014 , 15, 138-43	4.6	17
86	Daytime dysfunction in children with restless legs syndrome. <i>Journal of the Neurological Sciences</i> , 2014 , 336, 232-6	3.2	27

85	Impact of frequency of nightmares comorbid with insomnia on depression in Japanese rural community residents: a cross-sectional study. <i>Sleep Medicine</i> , 2014 , 15, 371-4	4.6	15
84	Selection of response criteria affects the success rate of oral appliance treatment for obstructive sleep apnea. <i>Sleep Medicine</i> , 2014 , 15, 367-70	4.6	22
83	Rapid maxillary expansion for obstructive sleep apnea: a lemon for lemonade?. <i>Journal of Clinical Sleep Medicine</i> , 2014 , 10, 233	3.1	6
82	Upper airway anatomical balance contributes to optimal continuous positive airway pressure for Japanese patients with obstructive sleep apnea syndrome. <i>Journal of Clinical Sleep Medicine</i> , 2014 , 10, 137-42	3.1	6
81	Sleep loss, sleep disorders and driving accidents. <i>Sleep and Biological Rhythms</i> , 2014 , 12, 96-105	1.3	9
80	C9orf72 repeat expansions in rapid eye movement sleep behaviour disorder. <i>Canadian Journal of Neurological Sciences</i> , 2014 , 41, 759-62	1	14
79	Validation of the Japanese version of the Ford Insomnia Response to Stress Test and the association of sleep reactivity with trait anxiety and insomnia. <i>Sleep Medicine</i> , 2014 , 15, 196-202	4.6	29
78	Quantitative assessment of isolated rapid eye movement (REM) sleep without atonia without clinical REM sleep behavior disorder: clinical and research implications. <i>Sleep Medicine</i> , 2014 , 15, 1009-15	4.6	21
77	Factors associated with duration before receiving definitive diagnosis of narcolepsy among Japanese patients affected with the disorder. <i>International Journal of Behavioral Medicine</i> , 2014 , 21, 966-70	2.6	5
76	Reducing dysfunctional beliefs about sleep does not significantly improve insomnia in cognitive behavioral therapy. <i>PLoS ONE</i> , 2014 , 9, e102565	3.7	20
75	Factors associated with long-term use of hypnotics among patients with chronic insomnia. <i>PLoS ONE</i> , 2014 , 9, e113753	3.7	15
74	The long-term treatment of restless legs syndrome/Willis-Ekbom disease: evidence-based guidelines and clinical consensus best practice guidance: a report from the International Restless Legs Syndrome Study Group. <i>Sleep Medicine</i> , 2013 , 14, 675-84	4.6	210
73	Clinical significance of REM sleep behavior disorder in Parkinson disease. <i>Sleep Medicine</i> , 2013 , 14, 131-4	4.6	87
72	Gabapentin enacarbil in Japanese patients with restless legs syndrome: a 12-week, randomized, double-blind, placebo-controlled, parallel-group study. <i>Current Medical Research and Opinion</i> , 2013 , 29, 13-21	2.5	27
71	Efficacy and safety of rotigotine in Japanese patients with restless legs syndrome: a phase 3, multicenter, randomized, placebo-controlled, double-blind, parallel-group study. <i>Sleep Medicine</i> , 2013 , 14, 1085-91	4.6	17
70	Cognitive behavioural therapy with behavioural analysis for pharmacological treatment-resistant chronic insomnia. <i>Psychiatry Research</i> , 2013 , 210, 515-21	9.9	15
69	Short sleep duration, sleep disorders, and traffic accidents. <i>IATSS Research</i> , 2013 , 37, 1-7	4.2	20
68	Current status of the treatment of insomnia. <i>Sleep and Biological Rhythms</i> , 2013 , 11, 1-1	1.3	

67	Efficacy and safety of adjunctive modafinil treatment on residual excessive daytime sleepiness among nasal continuous positive airway pressure-treated Japanese patients with obstructive sleep apnea syndrome: a double-blind placebo-controlled study. <i>Journal of Clinical Sleep Medicine</i> , 2013 , 9, 751-7	3.1	19
66	Efficacy, safety and risk of augmentation of rotigotine for treating restless legs syndrome. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013 , 40, 326-33	5.5	19
65	Reliability, validity, and responsiveness of the Japanese version of International Restless Legs Syndrome Study Group rating scale for restless legs syndrome in a clinical trial setting. <i>Psychiatry and Clinical Neurosciences</i> , 2013 , 67, 412-9	6.2	5
64	Factors associated with shift work disorder in nurses working with rapid-rotation schedules in Japan: the nurses' sleep health project. <i>Chronobiology International</i> , 2013 , 30, 628-36	3.6	87
63	Development and validation of the Japanese version of the Athens Insomnia Scale. <i>Psychiatry and Clinical Neurosciences</i> , 2013 , 67, 420-5	6.2	106
62	History of the researches on rapid eye movement sleep behavior disorder in Japan. <i>Sleep and Biological Rhythms</i> , 2013 , 11, 1-5	1.3	2
61	Differences in rapid eye movement sleep behavior disorder manifestation between synucleinopathies and tauopathies. <i>Sleep and Biological Rhythms</i> , 2013 , 11, 82-87	1.3	1
60	Oropharyngeal crowding and obesity as predictors of oral appliance treatment response to moderate obstructive sleep apnea. <i>Chest</i> , 2013 , 144, 558-563	5.3	37
59	Differences in brain morphological findings between narcolepsy with and without cataplexy. <i>PLoS ONE</i> , 2013 , 8, e81059	3.7	19
58	A two-year follow-up study on the symptoms of sleep disturbances/insomnia and their effects on daytime functioning. <i>Sleep Medicine</i> , 2012 , 13, 1115-21	4.6	24
57	Quality of life in patients with narcolepsy with cataplexy, narcolepsy without cataplexy, and idiopathic hypersomnia without long sleep time: comparison between patients on psychostimulants, drug-naïve patients and the general Japanese population. <i>Sleep Medicine</i> , 2012 , 13, 200-6	4.6	62
56	The factors associated with preferences for napping and drinking coffee as countermeasures for sleepiness at the wheel among Japanese drivers. <i>Sleep Medicine</i> , 2012 , 13, 354-61	4.6	14
55	Effects of nasal continuous positive airway pressure on panic disorder comorbid with obstructive sleep apnea syndrome. <i>Sleep Medicine</i> , 2012 , 13, 156-60	4.6	25
54	Relationship between napping pattern and nocturnal sleep among Japanese nursery school children. <i>Sleep Medicine</i> , 2012 , 13, 107-10	4.6	34
53	Comparison of REM sleep behaviour disorder variables between patients with progressive supranuclear palsy and those with Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , 2012 , 18, 394-6	3.6	29
52	Long-term efficacy and safety of gabapentin enacarbil in Japanese restless legs syndrome patients. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012 , 36, 251-7	5.5	26
51	Factors associated with severity of daytime sleepiness and indications for initiating treatment in patients with periodic limb movements during sleep. <i>Sleep and Biological Rhythms</i> , 2012 , 10, 187-194	1.3	1
50	Differences in findings of nocturnal polysomnography and multiple sleep latency test between narcolepsy and idiopathic hypersomnia. <i>Clinical Neurophysiology</i> , 2012 , 123, 137-41	4.3	36

49	Investigation of autonomic function in idiopathic REM sleep behavior disorder. <i>Journal of Neurology</i> , 2012 , 259, 1056-61	5.5	54
48	Effect of post-learning sleep versus wakefulness on advantageous decision-making: A preliminary study. <i>Sleep and Biological Rhythms</i> , 2012 , 10, 72-74	1.3	2
47	Insomnia as a risk for depression: a longitudinal epidemiologic study on a Japanese rural cohort. <i>Journal of Clinical Psychiatry</i> , 2012 , 73, 377-83	4.6	49
46	Effects of sleep-wake pattern on psychological distress in new recruits. <i>The Proceedings of the Annual Convention of the Japanese Psychological Association</i> , 2012 , 76, 2EVB09-2EVB09	0	
45	Detecting deteriorated vigilance using percentage of eyelid closure time during behavioral maintenance of wakefulness tests. <i>International Journal of Psychophysiology</i> , 2011 , 82, 269-74	2.9	52
44	Authors' Reply to the comments of Miyamoto et al. regarding "Cardiac (123)I-MIBG accumulation in Parkinson's disease differs in association with REM sleep behavior disorder". <i>Parkinsonism and Related Disorders</i> , 2011 , 17, 654	3.6	3
43	Relation between morningness-eveningness score and depressive symptoms among patients with delayed sleep phase syndrome. <i>Sleep Medicine</i> , 2011 , 12, 680-4	4.6	74
42	Prospective survey on the natural course of restless legs syndrome over two years in a closed cohort. <i>Sleep Medicine</i> , 2011 , 12, 821-6	4.6	25
41	Prevalence and clinical characteristics of restless legs syndrome in chronic kidney disease patients. <i>Sleep Medicine</i> , 2011 , 12, 1031-3	4.6	29
40	Correlations among insomnia symptoms, sleep medication use and depressive symptoms. <i>Psychiatry and Clinical Neurosciences</i> , 2011 , 65, 20-9	6.2	23
39	Comparison of the clinical features of rapid eye movement sleep behavior disorder in patients with Parkinson's disease and multiple system atrophy. <i>Psychiatry and Clinical Neurosciences</i> , 2011 , 65, 264-71	6.2	34
38	Changes in respiratory disorder parameters during the night in patients with obstructive sleep apnoea. <i>Respirology</i> , 2011 , 16, 116-23	3.6	7
37	Relationship between clinical characteristics of narcolepsy and CSF orexin-A levels. <i>Journal of Sleep Research</i> , 2011 , 20, 45-9	5.8	27
36	A meta-analysis on the treatment effectiveness of cognitive behavioral therapy for primary insomnia. <i>Sleep and Biological Rhythms</i> , 2011 , 9, 24-34	1.3	169
35	Clinical significance and management of insomnia. <i>Sleep and Biological Rhythms</i> , 2011 , 9, 63-72	1.3	4
34	Questionnaire-based evidence of association between sleepiness while driving and motor vehicle crashes that are subjectively not caused by falling asleep. <i>Sleep and Biological Rhythms</i> , 2011 , 9, 134-143	1.3	10
33	Changes in cerebral hemoglobin indices in obstructive sleep apnea syndrome with nasal continuous positive airway pressure treatment. <i>Sleep and Breathing</i> , 2011 , 15, 487-92	3.1	18
32	Restless legs syndrome (RLS) 2011 ,		1

31	Efficacy, safety and dose-response of pramipexole in Japanese patients with primary restless legs syndrome: randomized trial. <i>Neuropsychobiology</i> , 2011 , 63, 35-42	4	13
30	Short sleep duration and long spells of driving are associated with the occurrence of Japanese drivers' rear-end collisions and single-car accidents. <i>Journal of Sleep Research</i> , 2010 , 19, 310-6	5.8	33
29	Clinical characteristics of Japanese patients with familial obstructive sleep apnoea syndrome. <i>Respirology</i> , 2010 , 15, 93-8	3.6	5
28	Long-term open-label study of pramipexole in patients with primary restless legs syndrome. <i>Journal of the Neurological Sciences</i> , 2010 , 294, 62-6	3.2	30
27	Relationship between (123)I-MIBG scintigrams and REM sleep behavior disorder in Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , 2010 , 16, 683-5	3.6	54
26	Efficacy and safety of pramipexole in Japanese patients with primary restless legs syndrome: A polysomnographic randomized, double-blind, placebo-controlled study. <i>Sleep Medicine</i> , 2010 , 11, 11-6	4.6	24
25	Effects of insomnia and sleep medication on health-related quality of life. <i>Sleep Medicine</i> , 2010 , 11, 452-7	4.6	51
24	Clinical characteristics in two subgroups of obstructive sleep apnea syndrome in the elderly: comparison between cases with elderly and middle-age onset. <i>Chest</i> , 2010 , 137, 1310-5	5.3	22
23	Excessive daytime sleepiness among Japanese public transportation drivers engaged in shiftwork. <i>Journal of Occupational and Environmental Medicine</i> , 2010 , 52, 813-8	2	23
22	Gender differences in the clinical characteristics among Japanese patients with obstructive sleep apnea syndrome. <i>Chest</i> , 2009 , 135, 337-343	5.3	37
21	Pathogenetic heterogeneity of restless legs syndrome in Parkinson's disease. <i>Sleep and Biological Rhythms</i> , 2009 , 7, 31-33	1.3	1
20	Development of a Japanese version of the Epworth Sleepiness Scale (JESS) based on item response theory. <i>Sleep Medicine</i> , 2009 , 10, 556-65	4.6	195
19	Comparison of clinical characteristics among narcolepsy with and without cataplexy and idiopathic hypersomnia without long sleep time, focusing on HLA-DRB1(*)1501/DQB1(*)0602 finding. <i>Sleep Medicine</i> , 2009 , 10, 961-6	4.6	41
18	Validation of the Parkinson's disease sleep scale in Japanese patients: a comparison study using the Pittsburgh Sleep Quality Index, the Epworth Sleepiness Scale and Polysomnography. <i>Journal of the Neurological Sciences</i> , 2009 , 287, 36-40	3.2	37
17	Elevated risk of motor vehicle accident for male drivers with obstructive sleep apnea syndrome in the Tokyo metropolitan area. <i>Tohoku Journal of Experimental Medicine</i> , 2009 , 219, 11-6	2.4	42
16	Email-based epidemiological surveys on restless legs syndrome in Japan. <i>Sleep and Biological Rhythms</i> , 2008 , 6, 139-145	1.3	20
15	Clinical significance and correlates of behaviorally induced insufficient sleep syndrome. <i>Sleep Medicine</i> , 2008 , 9, 851-6	4.6	41
14	The prevalence of probable delayed-sleep-phase syndrome in students from junior high school to university in Tottori, Japan. <i>Tohoku Journal of Experimental Medicine</i> , 2008 , 216, 95-8	2.4	28

13	Clinical characteristics of restless legs syndrome in end-stage renal failure and idiopathic RLS patients. <i>Movement Disorders</i> , 2008 , 23, 811-6; quiz 926	7	47
12	Prevalence of restless legs syndrome in a rural community in Japan. <i>Movement Disorders</i> , 2008 , 23, 2363-9		101
11	Health-related quality of life among drug-naïve patients with narcolepsy with cataplexy, narcolepsy without cataplexy, and idiopathic hypersomnia without long sleep time. <i>Journal of Clinical Sleep Medicine</i> , 2008 , 4, 572-8	3.1	21
10	Health-Related Quality of Life Among Drug-Naïve Patients with Narcolepsy with Cataplexy, Narcolepsy Without Cataplexy, and Idiopathic Hypersomnia Without Long Sleep Time. <i>Journal of Clinical Sleep Medicine</i> , 2008 , 04, 572-578	3.1	51
9	Factors influencing subjective sleepiness in patients with obstructive sleep apnea syndrome. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 558-63	6.2	23
8	Prevalence and clinical characteristics of restless legs syndrome in Japanese patients with Parkinson's disease. <i>Movement Disorders</i> , 2006 , 21, 380-4	7	110
7	Clinical characteristics of Restless legs syndrome in patients with Parkinson's disease. <i>Journal of the Neurological Sciences</i> , 2006 , 250, 39-44	3.2	36
6	Narcolepsy without cataplexy: 2 subtypes based on CSF hypocretin-1/orexin-A findings. <i>Sleep</i> , 2006 , 29, 1439-43	1.1	39
5	Difference in the characteristics of subjective and objective sleepiness between narcolepsy and essential hypersomnia. <i>Psychiatry and Clinical Neurosciences</i> , 2005 , 59, 194-9	6.2	36
4	Periodic leg movements during sleep in Japanese community-dwelling adults based on the assessments of their bed partners. <i>Journal of Epidemiology</i> , 2003 , 13, 259-65	3.4	6
3	Clinical characteristics of essential hypersomnia syndrome. <i>Sleep and Biological Rhythms</i> , 2003 , 1, 229-231.3		5
2	CSF hypocretin-1 (orexin-A) concentrations in narcolepsy with and without cataplexy and idiopathic hypersomnia. <i>Journal of Sleep Research</i> , 2002 , 11, 91-3	5.8	167
1	Genome-wide association study of idiopathic hypersomnia in a Japanese population. <i>Sleep and Biological Rhythms</i> , 1	1.3	1