

# Rui Du

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1162112/publications.pdf>

Version: 2024-02-01

17  
papers

1,271  
citations

1040056

9  
h-index

888059

17  
g-index

19  
all docs

19  
docs citations

19  
times ranked

2080  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Multivariate Effect of Ketamine on PTSD: Systematic Review and Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2022, 13, 813103.	2.6	11
2	Rho GTPase-activating protein 35 suppresses gastric cancer metastasis by regulating cytoskeleton reorganization and epithelial-to-mesenchymal transition. <i>Bioengineered</i> , 2022, 13, 14605-14615.	3.2	4
3	Spinophilin modulates pain through suppressing dendritic spine morphogenesis via negative control of Rac1-ERK signaling in rat spinal dorsal horn. <i>Neurobiology of Disease</i> , 2021, 152, 105302.	4.4	4
4	Ceftriaxone Relieves Trigeminal Neuropathic Pain Through Suppression of Spatiotemporal Synaptic Plasticity via Restoration of Glutamate Transporter 1 in the Medullary Dorsal Horn. <i>Frontiers in Cellular Neuroscience</i> , 2020, 14, 199.	3.7	6
5	Image-Forming Visual Basis of Empathy for Pain in Mice. <i>Neuroscience Bulletin</i> , 2020, 36, 1563-1569.	2.9	7
6	The similar past pain experience evokes both observational contagious pain and consolation in stranger rat observers. <i>Neuroscience Letters</i> , 2020, 722, 134840.	2.1	11
7	Empathic Contagious Pain and Consolation in Laboratory Rodents: Species and Sex Comparisons. <i>Neuroscience Bulletin</i> , 2020, 36, 649-653.	2.9	28
8	Rat Model of Empathy for Pain. <i>Bio-protocol</i> , 2019, 9, e3266.	0.4	7
9	Validating Rat Model of Empathy for Pain: Effects of Pain Expressions in Social Partners. <i>Frontiers in Behavioral Neuroscience</i> , 2018, 12, 242.	2.0	36
10	The T-box transcription factor Brachyury promotes renal interstitial fibrosis by repressing E-cadherin expression. <i>Cell Communication and Signaling</i> , 2014, 12, 76.	6.5	13
11	Hypoxia-induced Bmi1 promotes renal tubular epithelial cellâ€mesenchymal transition and renal fibrosis via PI3K/Akt signal. <i>Molecular Biology of the Cell</i> , 2014, 25, 2650-2659.	2.1	98
12	Association of URG11 and Twist with clinical pathological characteristics and prognosis in patients with IgA nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, 2268-2276.	0.7	6
13	Hypoxia-Induced Down-Regulation of microRNA-34a Promotes EMT by Targeting the Notch Signaling Pathway in Tubular Epithelial Cells. <i>PLoS ONE</i> , 2012, 7, e30771.	2.5	154
14	URG11 promotes gastric cancer growth and invasion by activation of $\beta$ -catenin signalling pathway. <i>Journal of Cellular and Molecular Medicine</i> , 2010, 14, 621-635.	3.6	27
15	URG11 mediates hypoxia-induced epithelial-to-mesenchymal transition by modulation of E-cadherin and $\beta$ -catenin. <i>Biochemical and Biophysical Research Communications</i> , 2010, 391, 135-141.	2.1	22
16	Hypoxia-inducible factor-1 $\alpha$ induces Twist expression in tubular epithelial cells subjected to hypoxia, leading to epithelial-to-mesenchymal transition. <i>Kidney International</i> , 2009, 75, 1278-1287.	5.2	188
17	miRâ€15b and miRâ€16 modulate multidrug resistance by targeting BCL2 in human gastric cancer cells. <i>International Journal of Cancer</i> , 2008, 123, 372-379.	5.1	647