

Eija Rosti-Otajärvi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1162648/publications.pdf>

Version: 2024-02-01

13
papers

502
citations

932766

10
h-index

1125271

13
g-index

13
all docs

13
docs citations

13
times ranked

843
citing authors

#	ARTICLE	IF	CITATIONS
1	EpiTrack is a feasible tool for assessing attention and executive functions in patients with refractory epilepsy. <i>Epilepsy and Behavior</i> , 2021, 115, 107691.	0.9	7
2	Executive Functions May Predict Outcome in Deep Brain Stimulation of Anterior Nucleus of Thalamus for Treatment of Refractory Epilepsy. <i>Frontiers in Neurology</i> , 2018, 9, 324.	1.1	21
3	Validity and reliability of the Finnish version of the Multiple Sclerosis Impact Scale. <i>Brain and Behavior</i> , 2017, 7, e00725.	1.0	14
4	Validity and reliability of the Fatigue Severity Scale in Finnish multiple sclerosis patients. <i>Brain and Behavior</i> , 2017, 7, e00743.	1.0	55
5	Recovery from Mild Traumatic Brain Injury in Previously Healthy Adults. <i>Journal of Neurotrauma</i> , 2016, 33, 766-776.	1.7	143
6	Resilience Is Associated with Outcome from Mild Traumatic Brain Injury. <i>Journal of Neurotrauma</i> , 2015, 32, 942-949.	1.7	72
7	Is neuropsychological rehabilitation effective in multiple sclerosis?. <i>Neurodegenerative Disease Management</i> , 2014, 4, 147-154.	1.2	11
8	Predictors and impact of the working alliance in the neuropsychological rehabilitation of patients with multiple sclerosis. <i>Journal of the Neurological Sciences</i> , 2014, 338, 156-161.	0.3	9
9	Cognitive performance profile in different phenotypes of MS with cognitive complaints. <i>Multiple Sclerosis and Related Disorders</i> , 2014, 3, 463-472.	0.9	15
10	Neuropsychological rehabilitation does not improve cognitive performance but reduces perceived cognitive deficits in patients with multiple sclerosis: a randomised, controlled, multi-centre trial. <i>Multiple Sclerosis Journal</i> , 2014, 20, 99-107.	1.4	64
11	Neuropsychological rehabilitation has beneficial effects on perceived cognitive deficits in multiple sclerosis during nine-month follow-up. <i>Journal of the Neurological Sciences</i> , 2013, 334, 154-160.	0.3	40
12	Patient-related factors may affect the outcome of neuropsychological rehabilitation in multiple sclerosis. <i>Journal of the Neurological Sciences</i> , 2013, 334, 106-111.	0.3	21
13	Behavioural symptoms and impairments in multiple sclerosis: a systematic review and meta-analysis. <i>Multiple Sclerosis Journal</i> , 2013, 19, 31-45.	1.4	30