

Sotiris Amillis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11631364/publications.pdf>

Version: 2024-02-01

32
papers

1,507
citations

361045

20
h-index

433756

31
g-index

36
all docs

36
docs citations

36
times ranked

1937
citing authors

#	ARTICLE	IF	CITATIONS
1	Translocation of nutrient transporters to cell membrane via Golgi bypass in <i>Aspergillus nidulans</i> . EMBO Reports, 2020, 21, e49929.	2.0	15
2	The peroxisomal SspA protein is redundant for purine utilization but essential for peroxisome localization in septal pores in <i>Aspergillus nidulans</i> . Fungal Genetics and Biology, 2019, 132, 103259.	0.9	3
3	A pair of non-optimal codons are necessary for the correct biosynthesis of the <i>Aspergillus nidulans</i> urea transporter, UreA. Royal Society Open Science, 2019, 6, 190773.	1.1	3
4	Structural Lipids Enable the Formation of Functional Oligomers of the Eukaryotic Purine Symporter UapA. Cell Chemical Biology, 2018, 25, 840-848.e4.	2.5	64
5	Secretory Vesicle Polar Sorting, Endosome Recycling and Cytoskeleton Organization Require the AP-1 Complex in <i>Aspergillus nidulans</i> . Genetics, 2018, 209, 1121-1138.	1.2	21
6	Structural Aspects of UapA the H ⁺ -Xanthine/Uric Acid Transporter from <i>Aspergillus nidulans</i> . , 2018, , 1-7.		0
7	Comparative genomics reveals high biological diversity and specific adaptations in the industrially and medically important fungal genus <i>Aspergillus</i> . Genome Biology, 2017, 18, 28.	3.8	417
8	Substrate Specificity of the FurE Transporter Is Determined by Cytoplasmic Terminal Domain Interactions. Genetics, 2017, 207, 1387-1400.	1.2	23
9	Cryptic purine transporters in <i>Aspergillus nidulans</i> reveal the role of specific residues in the evolution of specificity in the NCS1 family. Molecular Microbiology, 2017, 103, 319-332.	1.2	22
10	The AP-2 complex has a specialized clathrin-independent role in apical endocytosis and polar growth in fungi. ELife, 2017, 6, .	2.8	45
11	BsdA ^Δ Bsd2 ^Δ -dependent vacuolar turnover of a misfolded version of the UapA transporter along the secretory pathway: prominent role of selective autophagy. Molecular Microbiology, 2016, 100, 893-911.	1.2	17
12	Structure of eukaryotic purine/H ⁺ symporter UapA suggests a role for homodimerization in transport activity. Nature Communications, 2016, 7, 11336.	5.8	108
13	Characterization of AnNce102 and its role in eisosome stability and sphingolipid biosynthesis. Scientific Reports, 2015, 5, 15200.	1.6	21
14	Oligomerization of the UapA Purine Transporter Is Critical for ER-Exit, Plasma Membrane Localization and Turnover. Journal of Molecular Biology, 2015, 427, 2679-2696.	2.0	36
15	Expression and specificity profile of the major acetate transporter AcpA in <i>Aspergillus nidulans</i> . Fungal Genetics and Biology, 2015, 76, 93-103.	0.9	22
16	Purine utilization proteins in the Eurotiales: Cellular compartmentalization, phylogenetic conservation and divergence. Fungal Genetics and Biology, 2014, 69, 96-108.	0.9	17
17	Modelling and mutational analysis of <i>Aspergillus nidulans</i> UreA, a member of the subfamily of urea/H ⁺ transporters in fungi and plants. Open Biology, 2014, 4, 140070.	1.5	13
18	The arrestin-like protein ArtA is essential for ubiquitination and endocytosis of the UapA transporter in response to both broad-range and specific signals. Molecular Microbiology, 2013, 88, 301-317.	1.2	60

#	ARTICLE	IF	CITATIONS
19	Modeling, Substrate Docking, and Mutational Analysis Identify Residues Essential for the Function and Specificity of a Eukaryotic Purine-Cytosine NCS1 Transporter. <i>Journal of Biological Chemistry</i> , 2012, 287, 36792-36803.	1.6	39
20	<i>Aspergillus nidulans</i> CkiA is an essential casein kinase I required for delivery of amino acid transporters to the plasma membrane. <i>Molecular Microbiology</i> , 2012, 84, 530-549.	1.2	45
21	Completing the purine utilisation pathway of <i>Aspergillus nidulans</i> . <i>Fungal Genetics and Biology</i> , 2011, 48, 840-848.	0.9	29
22	Mutational Analysis and Modeling Reveal Functionally Critical Residues in Transmembrane Segments 1 and 3 of the UapA Transporter. <i>Journal of Molecular Biology</i> , 2011, 411, 567-580.	2.0	25
23	Transport-dependent endocytosis and turnover of a uric acid-xanthine permease. <i>Molecular Microbiology</i> , 2010, 75, 246-260.	1.2	76
24	UreA, the major urea/H ⁺ symporter in <i>Aspergillus nidulans</i> . <i>Fungal Genetics and Biology</i> , 2010, 47, 1023-1033.	0.9	36
25	Convergent evolution and orphan genes in the Fur4p-like family and characterization of a general nucleoside transporter in <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2009, 73, 43-57.	1.2	40
26	Specific Interdomain Synergy in the UapA Transporter Determines Its Unique Specificity for Uric Acid among NAT Carriers. <i>Journal of Molecular Biology</i> , 2008, 382, 1121-1135.	2.0	55
27	Characterization and kinetics of the major purine transporters in <i>Aspergillus fumigatus</i> . <i>Fungal Genetics and Biology</i> , 2008, 45, 459-472.	0.9	21
28	Regulation of expression and kinetic modeling of substrate interactions of a uracil transporter in <i>Aspergillus nidulans</i> . <i>Molecular Membrane Biology</i> , 2007, 24, 206-214.	2.0	32
29	A Novel-type Substrate-selectivity Filter and ER-exit Determinants in the UapA Purine Transporter. <i>Journal of Molecular Biology</i> , 2006, 357, 808-819.	2.0	41
30	The AzgA Purine Transporter of <i>Aspergillus nidulans</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 3132-3141.	1.6	78
31	Transcription of purine transporter genes is activated during the isotropic growth phase of <i>Aspergillus nidulans</i> conidia. <i>Molecular Microbiology</i> , 2004, 52, 205-216.	1.2	42
32	Substitution F569S converts UapA, a specific uric acid-xanthine transporter, into a broad specificity transporter for purine-related solutes. <i>Journal of Molecular Biology</i> , 2001, 313, 765-774.	2.0	38