

# Jiejie Xu

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1164160/jiejie-xu-publications-by-year.pdf>

**Version:** 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

148  
papers

2,447  
citations

24  
h-index

41  
g-index

157  
ext. papers

3,183  
ext. citations

4.9  
avg, IF

4.94  
L-index

#	Paper	IF	Citations
148	ASO Visual Abstract: Stromal Tumor-Associated Macrophage Infiltration Predicts Poor Clinical Outcomes in Muscle-Invasive Bladder Cancer Patients.. <i>Annals of Surgical Oncology</i> , <b>2022</b> , 29, 2504	3.1	
147	Immune inactivation by neuropilin-1 predicts clinical outcome and therapeutic benefit in muscle-invasive bladder cancer.. <i>Cancer Immunology, Immunotherapy</i> , <b>2022</b> , 1	7.4	
146	Stromal Tumor-Associated Macrophage Infiltration Predicts Poor Clinical Outcomes in Muscle-Invasive Bladder Cancer Patients.. <i>Annals of Surgical Oncology</i> , <b>2022</b> , 1	3.1	0
145	CD103CD8 tissue-resident memory T cell infiltration predicts clinical outcome and adjuvant therapeutic benefit in muscle-invasive bladder cancer.. <i>British Journal of Cancer</i> , <b>2022</b> ,	8.7	1
144	Infiltration and Polarization of Tumor-associated Macrophages Predict Prognosis and Therapeutic Benefit in Muscle-Invasive Bladder Cancer. <i>Cancer Immunology, Immunotherapy</i> , <b>2021</b> , 1	7.4	0
143	ASO Author Reflections: Optimization of Tumor Therapy for the Specific Immune Microenvironment of Gastric Cancer. <i>Annals of Surgical Oncology</i> , <b>2021</b> , 28, 6451-6452	3.1	
142	Intratumoral CXCR5CD8T associates with favorable clinical outcomes and immunogenic contexture in gastric cancer. <i>Nature Communications</i> , <b>2021</b> , 12, 3080	17.4	4
141	Latency-associated peptide identifies therapeutically resistant muscle-invasive bladder cancer with poor prognosis. <i>Cancer Immunology, Immunotherapy</i> , <b>2021</b> , 1	7.4	1
140	Blocking siglec-10 tumor-associated macrophages improves anti-tumor immunity and enhances immunotherapy for hepatocellular carcinoma. <i>Experimental Hematology and Oncology</i> , <b>2021</b> , 10, 36	7.8	7
139	Lymphocyte-activation gene 3 expression associates with poor prognosis and immunoevasive contexture in Epstein-Barr virus-positive and MLH1-defective gastric cancer patients. <i>International Journal of Cancer</i> , <b>2021</b> , 148, 759-768	7.5	4
138	Poor clinical outcomes and immunoevasive contexture in CXCL13+CD8+ T cells enriched gastric cancer patients. <i>Oncolmmunology</i> , <b>2021</b> , 10, 1915560	7.2	4
137	Intratumoral CXCL13CD8T cell infiltration determines poor clinical outcomes and immunoevasive contexture in patients with clear cell renal cell carcinoma <b>2021</b> , 9,		20
136	Clinical Outcomes and Immune Metrics in Intratumoral Basophil-Enriched Gastric Cancer Patients. <i>Annals of Surgical Oncology</i> , <b>2021</b> , 28, 6439-6450	3.1	3
135	Poor clinical outcomes and immunoevasive contexture in SIRP1 <sup>+</sup> tumor-associated macrophages enriched muscle-invasive bladder cancer patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2021</b> , 40, 109.e11-109.e11	2.8	0
134	Impact of intratumoural CD73 expression on prognosis and therapeutic response in patients with gastric cancer. <i>European Journal of Cancer</i> , <b>2021</b> , 157, 114-123	7.5	3
133	Immune inactivation by APOBEC3B enrichment predicts response to chemotherapy and survival in gastric cancer. <i>Oncolmmunology</i> , <b>2021</b> , 10, 1975386	7.2	2
132	CCR8 blockade primes anti-tumor immunity through intratumoral regulatory T cells destabilization in muscle-invasive bladder cancer. <i>Cancer Immunology, Immunotherapy</i> , <b>2020</b> , 69, 1855-1867	7.4	11

131	CCR5 blockade inflames antitumor immunity in BAP1-mutant clear cell renal cell carcinoma <b>2020</b> , 8,		4
130	Stromal LAG-3 cells infiltration defines poor prognosis subtype muscle-invasive bladder cancer with immunoevasive contexture <b>2020</b> , 8,		13
129	Lauren classification identifies distinct prognostic value and functional status of intratumoral CD8 T cells in gastric cancer. <i>Cancer Immunology, Immunotherapy</i> , <b>2020</b> , 69, 1327-1336	7.4	13
128	Identification and validation of an immunogenic subtype of gastric cancer with abundant intratumoural CD103CD8 T cells conferring favourable prognosis. <i>British Journal of Cancer</i> , <b>2020</b> , 122, 1525-1534	8.7	15
127	Poor clinical outcomes of intratumoral dendritic cell-specific intercellular adhesion molecule 3-grabbing non-integrin-positive macrophages associated with immune evasion in gastric cancer. <i>European Journal of Cancer</i> , <b>2020</b> , 128, 27-37	7.5	12
126	Blockade of DC-SIGN Tumor-Associated Macrophages Reactivates Antitumor Immunity and Improves Immunotherapy in Muscle-Invasive Bladder Cancer. <i>Cancer Research</i> , <b>2020</b> , 80, 1707-1719	10.1	27
125	Identification and validation of dichotomous immune subtypes based on intratumoral immune cells infiltration in clear cell renal cell carcinoma patients <b>2020</b> , 8,		17
124	Identification and validation of poor prognosis immunoevasive subtype of muscle-invasive bladder cancer with tumor-infiltrating podoplanin cell abundance. <i>Oncolmunology</i> , <b>2020</b> , 9, 1747333	7.2	9
123	Latency-associated Peptide Identifies Immunoevasive Subtype Gastric Cancer With Poor Prognosis and Inferior Chemotherapeutic Responsiveness. <i>Annals of Surgery</i> , <b>2020</b> , 275,	7.8	6
122	Intratumoral IL22-producing cells define immunoevasive subtype muscle-invasive bladder cancer with poor prognosis and superior nivolumab responses. <i>International Journal of Cancer</i> , <b>2020</b> , 146, 542-552	7.5	7
121	PAK1 expression determines poor prognosis and immune evasion in metastatic renal cell carcinoma patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2020</b> , 38, 293-304	2.8	5
120	Tumor-infiltrating podoplanin cells in gastric cancer: clinical outcomes and association with immune contexture. <i>Oncolmunology</i> , <b>2020</b> , 9, 1845038	7.2	1
119	Tumor-infiltrating TNFRSF9 CD8 T cells define different subsets of clear cell renal cell carcinoma with prognosis and immunotherapeutic response. <i>Oncolmunology</i> , <b>2020</b> , 9, 1838141	7.2	8
118	Poor clinical outcomes and immunoevasive contexture in interleukin-9 abundant muscle-invasive bladder cancer. <i>International Journal of Cancer</i> , <b>2020</b> , 147, 3539-3549	7.5	3
117	Intratumoral CCR5 neutrophils identify immunogenic subtype muscle-invasive bladder cancer with favorable prognosis and therapeutic responses. <i>Oncolmunology</i> , <b>2020</b> , 9, 1802176	7.2	2
116	Poliovirus receptor CD155 is up-regulated in muscle-invasive bladder cancer and predicts poor prognosis. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2020</b> , 38, 41.e11-41.e18	2.8	8
115	Tumor-infiltrating CD39CD8 T cells determine poor prognosis and immune evasion in clear cell renal cell carcinoma patients. <i>Cancer Immunology, Immunotherapy</i> , <b>2020</b> , 69, 1565-1576	7.4	38
114	Tumor-infiltrating IL-17A cells determine favorable prognosis and adjuvant chemotherapeutic response in muscle-invasive bladder cancer. <i>Oncolmunology</i> , <b>2020</b> , 9, 1747332	7.2	4

113	Tumour-associated macrophages-derived CXCL8 determines immune evasion through autonomous PD-L1 expression in gastric cancer. <i>Gut</i> , <b>2019</b> , 68, 1764-1773	19.2	118
112	Failure to Cite Related Studies and Report Complete Information on Patients and Tissue Samples. <i>JAMA Surgery</i> , <b>2019</b> , 154, 362-363	5.4	0
111	Tumor-associated macrophages expressing galectin-9 identify immunoevasive subtype muscle-invasive bladder cancer with poor prognosis but favorable adjuvant chemotherapeutic response. <i>Cancer Immunology, Immunotherapy</i> , <b>2019</b> , 68, 2067-2080	7.4	20
110	Tumor infiltrating mast cells determine oncogenic HIF-2 $\alpha$ -conferred immune evasion in clear cell renal cell carcinoma. <i>Cancer Immunology, Immunotherapy</i> , <b>2019</b> , 68, 731-741	7.4	21
109	Tumor-infiltrating neutrophils predict therapeutic benefit of tyrosine kinase inhibitors in metastatic renal cell carcinoma. <i>Oncotarget</i> , <b>2019</b> , 8, e1515611	7.2	3
108	CD19 tumor-infiltrating B-cells prime CD4 T-cell immunity and predict platinum-based chemotherapy efficacy in muscle-invasive bladder cancer. <i>Cancer Immunology, Immunotherapy</i> , <b>2019</b> , 68, 45-56	7.4	22
107	Tumor-associated Macrophage-derived Interleukin-23 Interlinks Kidney Cancer Glutamine Addiction with Immune Evasion. <i>European Urology</i> , <b>2019</b> , 75, 752-763	10.2	61
106	Identification and Validation of Stromal Immunity Predict Survival and Benefit from Adjuvant Chemotherapy in Patients with Muscle-Invasive Bladder Cancer. <i>Clinical Cancer Research</i> , <b>2018</b> , 24, 3069-3078	12.9	83
105	CXCR1 expression predicts benefit from tyrosine kinase inhibitors therapy in patients with metastatic renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2018</b> , 36, 242.e15-242.e21	15.8	221
104	Evaluation of Tumor Pseudocapsule Status and its Prognostic Significance in Renal Cell Carcinoma. <i>Journal of Urology</i> , <b>2018</b> , 199, 915-920	2.5	10
103	Tumor-infiltrating mast cells predict prognosis and gemcitabine-based adjuvant chemotherapeutic benefit in biliary tract cancer patients. <i>BMC Cancer</i> , <b>2018</b> , 18, 313	4.8	6
102	C-C motif chemokine 22 predicts postoperative prognosis and adjuvant chemotherapeutic benefits in patients with stage II/III gastric cancer. <i>Oncotarget</i> , <b>2018</b> , 7, e1433517	7.2	13
101	CXCL13 expression is prognostic and predictive for postoperative adjuvant chemotherapy benefit in patients with gastric cancer. <i>Cancer Immunology, Immunotherapy</i> , <b>2018</b> , 67, 261-269	7.4	28
100	HLA class I expression predicts prognosis and therapeutic benefits from tyrosine kinase inhibitors in metastatic renal-cell carcinoma patients. <i>Cancer Immunology, Immunotherapy</i> , <b>2018</b> , 67, 79-87	7.4	5
99	B4GALT1 expression predicts prognosis and adjuvant chemotherapy benefits in muscle-invasive bladder cancer patients. <i>BMC Cancer</i> , <b>2018</b> , 18, 590	4.8	9
98	Tumor infiltrating CD19 B lymphocytes predict prognostic and therapeutic benefits in metastatic renal cell carcinoma patients treated with tyrosine kinase inhibitors. <i>Oncotarget</i> , <b>2018</b> , 7, e1477461	7.2	86
97	Prognostic significance of ST6GalNAc-1 expression in patients with non-metastatic clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2018</b> , 9, 3112-3120	3.3	3
96	Tumor-infiltrating Neutrophils is Prognostic and Predictive for Postoperative Adjuvant Chemotherapy Benefit in Patients With Gastric Cancer. <i>Annals of Surgery</i> , <b>2018</b> , 267, 311-318	7.8	98

95	Prognostic and Predictive Value of O-methylguanine Methyltransferase for Chemotherapy in Patients with Muscle-Invasive Bladder Cancer. <i>Annals of Surgical Oncology</i> , <b>2018</b> , 25, 342-348	3.1	3
94	Tumor stroma-infiltrating mast cells predict prognosis and adjuvant chemotherapeutic benefits in patients with muscle invasive bladder cancer. <i>Oncolmmunology</i> , <b>2018</b> , 7, e1474317	7.2	39
93	Tumor-infiltrating neutrophils predict prognosis and adjuvant chemotherapeutic benefit in patients with biliary cancer. <i>Cancer Science</i> , <b>2018</b> , 109, 2266-2274	6.9	17
92	Tumor-infiltrating neutrophils predict benefit from adjuvant chemotherapy in patients with muscle invasive bladder cancer. <i>Oncolmmunology</i> , <b>2017</b> , 6, e1293211	7.2	37
91	Low CCL17 expression associates with unfavorable postoperative prognosis of patients with clear cell renal cell carcinoma. <i>BMC Cancer</i> , <b>2017</b> , 17, 117	4.8	5
90	Beta-1,4-galactosyltransferase II predicts poor prognosis of patients with non-metastatic clear-cell renal cell carcinoma. <i>Tumor Biology</i> , <b>2017</b> , 39, 1010428317691417	2.9	3
89	Galectin-9 as a prognostic and predictive biomarker in bladder urothelial carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2017</b> , 35, 349-355	2.8	19
88	Tumor Infiltrating Mast Cells (TIMs) Confers a Marked Survival Advantage in Nonmetastatic Clear-Cell Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , <b>2017</b> , 24, 1435-1442	3.1	24
87	Enhancement of Siglec-8 expression predicts adverse prognosis in patients with clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2017</b> , 35, 607.e1-607.e8	2.8	3
86	Decreased expression of granulocyte-macrophage colony-stimulating factor is associated with adverse clinical outcome in patients with gastric cancer undergoing gastrectomy. <i>Oncology Letters</i> , <b>2017</b> , 14, 4701-4707	2.6	0
85	High expression of Mucin13 associates with grimmer postoperative prognosis of patients with non-metastatic clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 7548-7558	3.3	7
84	High expression of CXC chemokine receptor 6 associates with poor prognosis in patients with clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2017</b> , 35, 675.e17-675.e24	2.8	9
83	Association of O6-Methylguanine-DNA Methyltransferase Protein Expression With Postoperative Prognosis and Adjuvant Chemotherapeutic Benefits Among Patients With Stage II or III Gastric Cancer. <i>JAMA Surgery</i> , <b>2017</b> , 152, e173120	5.4	14
82	Tumor-infiltrating T <sub>H</sub> 1 cells predict prognosis and adjuvant chemotherapeutic benefit in patients with gastric cancer. <i>Oncolmmunology</i> , <b>2017</b> , 6, e1353858	7.2	19
81	Decreased expression of JMJD3 predicts poor prognosis of patients with clear cell renal cell carcinoma. <i>Oncology Letters</i> , <b>2017</b> , 14, 1550-1560	2.6	7
80	High CXC chemokine receptor 1 level represents an independent negative prognosticator in non-metastatic clear-cell renal cell carcinoma patients. <i>Oncolmmunology</i> , <b>2017</b> , 6, e1359450	7.2	5
79	Prognostic value of copper transporter 1 expression in patients with clear cell renal cell carcinoma. <i>Oncology Letters</i> , <b>2017</b> , 14, 5791-5800	2.6	6
78	Prognostic value of CC-chemokine receptor seven expression in patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitor. <i>BMC Cancer</i> , <b>2017</b> , 17, 70	4.8	10

77	CXC chemokine receptor 1 predicts postoperative prognosis and chemotherapeutic benefits for TNM II and III resectable gastric cancer patients. <i>Oncotarget</i> , <b>2017</b> , 8, 20328-20339	3.3	10
76	Low CCL-21 expression associates with unfavorable postoperative prognosis of patients with metastatic renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 25650-25659	3.3	5
75	High NUCB2 expression level represents an independent negative prognostic factor in Chinese cohorts of non-metastatic clear cell renal cell carcinoma patients. <i>Oncotarget</i> , <b>2017</b> , 8, 35244-35254	3.3	7
74	Prognostic role of N-Acetylgalactosaminyltransferase 10 in metastatic renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 14995-15003	3.3	3
73	High mucin 5AC expression predicts adverse postoperative recurrence and survival of patients with clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 59777-59790	3.3	8
72	High expression of FUT3 is linked to poor prognosis in clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 61036-61047	3.3	5
71	IRF5 is associated with adverse postoperative prognosis of patients with non-metastatic clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 44186-44194	3.3	2
70	Prognostic value of granulocyte colony-stimulating factor in patients with non-metastatic clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 69961-69971	3.3	7
69	Stathmin 1 expression predicts prognosis and benefits from adjuvant chemotherapy in patients with gallbladder carcinoma. <i>Oncotarget</i> , <b>2017</b> , 8, 108548-108555	3.3	5
68	High truncated-O-glycan score predicts adverse clinical outcome in patients with localized clear-cell renal cell carcinoma after surgery. <i>Oncotarget</i> , <b>2017</b> , 8, 80083-80092	3.3	
67	An indel polymorphism within pre-miR3131 confers risk for hepatocellular carcinoma. <i>Carcinogenesis</i> , <b>2017</b> , 38, 168-176	4.6	9
66	High Expression of Colony-Stimulating Factor 1 Receptor Associates with Unfavorable Cancer-Specific Survival of Patients with Clear Cell Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , <b>2016</b> , 23, 1044-52	3.1	8
65	Increased expression of C-C motif ligand 2 associates with poor prognosis in patients with gastric cancer after gastrectomy. <i>Tumor Biology</i> , <b>2016</b> , 37, 3285-93	2.9	8
64	Enhancer of zeste homolog 2 (EZH2) promotes tumour cell migration and invasion via epigenetic repression of E-cadherin in renal cell carcinoma. <i>BJU International</i> , <b>2016</b> , 117, 351-62	5.6	64
63	Prognostic Value of SETD2 Expression in Patients with Metastatic Renal Cell Carcinoma Treated with Tyrosine Kinase Inhibitors. <i>Journal of Urology</i> , <b>2016</b> , 196, 1363-1370	2.5	30
62	High Level of Anaphylatoxin C5a Predicts Poor Clinical Outcome in Patients with Clear Cell Renal Cell Carcinoma. <i>Scientific Reports</i> , <b>2016</b> , 6, 29177	4.9	12
61	Dectin-1 predicts adverse postoperative prognosis of patients with clear cell renal cell carcinoma. <i>Scientific Reports</i> , <b>2016</b> , 6, 32657	4.9	5
60	The Presence of Vascular Mimicry Predicts High Risk of Clear Cell Renal Cell Carcinoma after Radical Nephrectomy. <i>Journal of Urology</i> , <b>2016</b> , 196, 335-42	2.5	4



59	Decreased expression of Siglec-8 associates with poor prognosis in patients with gastric cancer after surgical resection. <i>Tumor Biology</i> , <b>2016</b> , 37, 10883-91	2.9	9
58	High expression of chemokine CCL2 is associated with recurrence after surgery in clear-cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2016</b> , 34, 238.e19-26	2.8	11
57	Decreased expression of CTR2 predicts poor prognosis of patients with clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2016</b> , 34, 5.e1-9	2.8	5
56	IL-33 is associated with unfavorable postoperative survival of patients with clear-cell renal cell carcinoma. <i>Tumor Biology</i> , <b>2016</b> , 37, 11127-34	2.9	11
55	Increased expression of interleukin-8 is an independent indicator of poor prognosis in clear-cell renal cell carcinoma. <i>Tumor Biology</i> , <b>2016</b> , 37, 4523-9	2.9	8
54	Prognostic value of preoperative lymphocyte to monocyte ratio in patients with nonmetastatic clear cell renal cell carcinoma. <i>Tumor Biology</i> , <b>2016</b> , 37, 4613-20	2.9	16
53	Interleukin-13 receptor $\alpha$ is associated with poor prognosis in patients with gastric cancer after gastrectomy. <i>Oncotarget</i> , <b>2016</b> , 7, 49281-49288	3.3	14
52	CCL2/CCR2 axis is associated with postoperative survival and recurrence of patients with non-metastatic clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 51525-51534	3.3	23
51	High CLEC-2 expression associates with unfavorable postoperative prognosis of patients with clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 63661-63668	3.3	7
50	A three-molecule score based on Notch pathway predicts poor prognosis in non-metastasis clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 68559-68570	3.3	5
49	Dot1l expression predicts adverse postoperative prognosis of patients with clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 84775-84784	3.3	7
48	Prognostic value of vascular mimicry in patients with urothelial carcinoma of the bladder after radical cystectomy. <i>Oncotarget</i> , <b>2016</b> , 7, 76214-76223	3.3	3
47	Enrichment of C5a-C5aR axis predicts poor postoperative prognosis of patients with clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 80925-80934	3.3	16
46	Positive intratumoral chemokine (C-C motif) receptor 8 expression predicts high recurrence risk of post-operation clear-cell renal cell carcinoma patients. <i>Oncotarget</i> , <b>2016</b> , 7, 8413-21	3.3	5
45	Elevated expression of IFN-inducible CXCR3 ligands predicts poor prognosis in patients with non-metastatic clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 13976-83	3.3	19
44	High expression of C-C chemokine receptor 2 associates with poor overall survival in gastric cancer patients after surgical resection. <i>Oncotarget</i> , <b>2016</b> , 7, 23909-18	3.3	7
43	Granulocyte macrophage colony-stimulating factor predicts postoperative recurrence of clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 24527-36	3.3	6
42	Increased B4GALT1 expression associates with adverse outcome in patients with non-metastatic clear cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 32723-30	3.3	17

41	High expression of galectin-7 associates with poor overall survival in patients with non-metastatic clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 41986-41995	3.3	4
40	Increased expression of MUC3A is associated with poor prognosis in localized clear-cell renal cell carcinoma. <i>Oncotarget</i> , <b>2016</b> , 7, 50017-50026	3.3	12
39	Increased expression of IDO associates with poor postoperative clinical outcome of patients with gastric adenocarcinoma. <i>Scientific Reports</i> , <b>2016</b> , 6, 21319	4.9	50
38	Glycoprotein 130 is associated with adverse postoperative clinical outcomes of patients with late-stage non-metastatic gastric cancer. <i>Scientific Reports</i> , <b>2016</b> , 6, 38364	4.9	4
37	Podoplanin associates with adverse postoperative prognosis of patients with clear cell renal cell carcinoma. <i>Cancer Science</i> , <b>2016</b> , 107, 1243-9	6.9	3
36	Prognostic value of UTX expression in patients with clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2016</b> , 34, 338.e19-27	2.8	5
35	High mucin-7 expression is an independent predictor of adverse clinical outcomes in patients with clear-cell renal cell carcinoma. <i>Tumor Biology</i> , <b>2016</b> , 37, 15193-15201	2.9	10
34	Galectin-8 is associated with recurrence and survival of patients with non-metastatic gastric cancer after surgery. <i>Tumor Biology</i> , <b>2016</b> , 37, 12635-12642	2.9	12
33	β,6-N-acetylglucosaminyltransferase V predicts recurrence and survival of patients with clear-cell renal cell carcinoma after surgical resection. <i>World Journal of Urology</i> , <b>2015</b> , 33, 1791-9	4	9
32	p21-activated kinase 1 predicts recurrence and survival in patients with non-metastatic clear cell renal cell carcinoma. <i>International Journal of Urology</i> , <b>2015</b> , 22, 447-53	2.3	5
31	High APOBEC3B expression is a predictor of recurrence in patients with low-risk clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2015</b> , 33, 340.e1-8	2.8	27
30	p21-Activated kinase 4 predicts early recurrence and poor survival in patients with nonmetastatic clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2015</b> , 33, 205.e13-21	2.8	6
29	Association between indel polymorphism in the promoter region of lncRNA GAS5 and the risk of hepatocellular carcinoma. <i>Carcinogenesis</i> , <b>2015</b> , 36, 1136-43	4.6	90
28	Interleukin-11 receptor predicts post-operative clinical outcome in patients with early-stage clear-cell renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , <b>2015</b> , 45, 202-9	2.8	15
27	Galectin-8 predicts postoperative recurrence of patients with localized T1 clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2015</b> , 33, 112.e1-8	2.8	5
26	Increased expression of colony stimulating factor-1 is a predictor of poor prognosis in patients with clear-cell renal cell carcinoma. <i>BMC Cancer</i> , <b>2015</b> , 15, 67	4.8	22
25	Snail predicts recurrence and survival of patients with localized clear cell renal cell carcinoma after surgical resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2015</b> , 33, 69.e1-10	2.8	12
24	Clinical significance of tumor-derived IL-1β and IL-18 in localized renal cell carcinoma: Associations with recurrence and survival. <i>Urologic Oncology: Seminars and Original Investigations</i> , <b>2015</b> , 33, 68.e9-16	2.8	28



23	EZH2-mediated loss of miR-622 determines CXCR4 activation in hepatocellular carcinoma. <i>Nature Communications</i> , <b>2015</b> , 6, 8494	17.4	79
22	Tumor Suppressive Function of p21-activated Kinase 6 in Hepatocellular Carcinoma. <i>Journal of Biological Chemistry</i> , <b>2015</b> , 290, 28489-28501	5.4	14
21	CXC chemokine receptor 2 is associated with postoperative recurrence and survival of patients with non-metastatic clear-cell renal cell carcinoma. <i>European Journal of Cancer</i> , <b>2015</b> , 51, 1953-61	7.5	23
20	Infiltration of diametrically polarized macrophages predicts overall survival of patients with gastric cancer after surgical resection. <i>Gastric Cancer</i> , <b>2015</b> , 18, 740-50	7.6	100
19	High expression of interleukin-11 is an independent indicator of poor prognosis in clear-cell renal cell carcinoma. <i>Cancer Science</i> , <b>2015</b> , 106, 592-7	6.9	19
18	P2X7 receptor predicts postoperative cancer-specific survival of patients with clear-cell renal cell carcinoma. <i>Cancer Science</i> , <b>2015</b> , 106, 1224-31	6.9	23
17	High expression of Solute Carrier Family 1, member 5 (SLC1A5) is associated with poor prognosis in clear-cell renal cell carcinoma. <i>Scientific Reports</i> , <b>2015</b> , 5, 16954	4.9	33
16	The prognostic value of CXC-chemokine receptor 2 (CXCR2) in gastric cancer patients. <i>BMC Cancer</i> , <b>2015</b> , 15, 766	4.8	22
15	Prognostic significance of ST3GAL-1 expression in patients with clear cell renal cell carcinoma. <i>BMC Cancer</i> , <b>2015</b> , 15, 880	4.8	13
14	High peritumoral Bmi-1 expression is an independent prognosticator of poor prognosis in renal cell carcinoma. <i>Tumor Biology</i> , <b>2015</b> , 36, 8007-14	2.9	3
13	Prognostic significance of $\alpha$ 1,6-N-acetylglucosaminyltransferase V expression in patients with hepatocellular carcinoma. <i>Japanese Journal of Clinical Oncology</i> , <b>2015</b> , 45, 844-53	2.8	11
12	Prognostic value of interleukin-6 and interleukin-6 receptor in organ-confined clear-cell renal cell carcinoma: a 5-year conditional cancer-specific survival analysis. <i>British Journal of Cancer</i> , <b>2015</b> , 113, 1581-9	8.7	22
11	Notch1 predicts recurrence and survival of patients with clear-cell renal cell carcinoma after surgical resection. <i>Urology</i> , <b>2015</b> , 85, 483.e9-483.e14	1.6	6
10	Expression of IL-4 and IL-13 predicts recurrence and survival in localized clear-cell renal cell carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , <b>2015</b> , 8, 1594-603	1.4	14
9	Decreased expression of mucin 18 is associated with unfavorable postoperative prognosis in patients with clear cell renal cell carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , <b>2015</b> , 8, 11005-14	1.4	6
8	Expression of Jagged1 predicts postoperative clinical outcome of patients with gastric cancer. <i>International Journal of Clinical and Experimental Medicine</i> , <b>2015</b> , 8, 14782-92		3
7	Prognostic value of diametrically polarized tumor-associated macrophages in renal cell carcinoma. <i>Annals of Surgical Oncology</i> , <b>2014</b> , 21, 3142-50	3.1	80
6	Discovery of specific metastasis-related N-glycan alterations in epithelial ovarian cancer based on quantitative glycomics. <i>PLoS ONE</i> , <b>2014</b> , 9, e87978	3.7	35

5	GALNT4 predicts clinical outcome in patients with clear cell renal cell carcinoma. <i>Journal of Urology</i> , <b>2014</b> , 192, 1534-41	2.5	10
4	Functional short tandem repeat polymorphism of PTPN11 and susceptibility to hepatocellular carcinoma in Chinese populations. <i>PLoS ONE</i> , <b>2014</b> , 9, e106841	3.7	6
3	Hepatitis B virus X protein confers resistance of hepatoma cells to anoikis by up-regulating and activating p21-activated kinase 1. <i>Gastroenterology</i> , <b>2012</b> , 143, 199-212.e4	13.3	60
2	Hepatitis B virus large surface antigen promotes liver carcinogenesis by activating the Src/PI3K/Akt pathway. <i>Cancer Research</i> , <b>2011</b> , 71, 7547-57	10.1	70
1	Identification of beta-1,4-galactosyltransferase I as a target gene of HBx-induced cell cycle progression of hepatoma cell. <i>Journal of Hepatology</i> , <b>2008</b> , 49, 1029-37	13.4	25