

Priscila Chaverri

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11645948/publications.pdf>

Version: 2024-02-01

47
papers

2,229
citations

257101

24
h-index

223531

46
g-index

48
all docs

48
docs citations

48
times ranked

2441
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Biodeterioration and cellulolytic activity by fungi isolated from a nineteenth-century painting at the National Theatre of Costa Rica. <i>Fungal Biology</i> , 2022, 126, 101-112. | 1.1 | 10 |
| 2 | Tolerance and Biological Removal of Fungicides by <i>Trichoderma</i> Species Isolated From the Endosphere of Wild Rubiaceae Plants. <i>Frontiers in Agronomy</i> , 2022, 3, . | 1.5 | 10 |
| 3 | Fungal communities in feces of the frugivorous bat <i>Ectophylla alba</i> and its highly specialized <i>Ficus colubrinae</i> diet. <i>Animal Microbiome</i> , 2022, 4, 24. | 1.5 | 2 |
| 4 | An Integrative View of the Phyllosphere Mycobiome of Native Rubber Trees in the Brazilian Amazon. <i>Journal of Fungi</i> (Basel, Switzerland), 2022, 8, 373. | 1.5 | 7 |
| 5 | Virulence of native isolates of entomopathogenic fungi (Hypocreales) against the "sweetpotato whitefly" <i>Bemisia tabaci</i> (Hemiptera: Aleyrodidae), including the effects of temperature and fungicides. <i>Journal of Invertebrate Pathology</i> , 2022, 192, 107787. | 1.5 | 5 |
| 6 | Expanding the <i>Trichoderma harzianum</i> species complex: Three new species from Argentine natural and cultivated ecosystems. <i>Mycologia</i> , 2021, 113, 1-20. | 0.8 | 10 |
| 7 | Response of psychrophilic plant endosymbionts to experimental temperature increase. <i>Royal Society Open Science</i> , 2020, 7, 201405. | 1.1 | 1 |
| 8 | Endophytes from Wild Rubber Trees as Antagonists of the Pathogen <i>Corynespora cassiicola</i> . <i>Phytopathology</i> , 2019, 109, 1888-1899. | 1.1 | 20 |
| 9 | Exploration of stem endophytic communities revealed developmental stage as one of the drivers of fungal endophytic community assemblages in two Amazonian hardwood genera. <i>Scientific Reports</i> , 2019, 9, 12685. | 1.6 | 29 |
| 10 | Improving taxonomic accuracy for fungi in public sequence databases: applying "one name one species"™ in well-defined genera with <i>Trichoderma/Hypocrea</i> as a test case. <i>Database: the Journal of Biological Databases and Curation</i> , 2017, 2017, . | 1.4 | 28 |
| 11 | Overlooked competing asexual and sexually typified generic names of Ascomycota with recommendations for their use or protection. <i>IMA Fungus</i> , 2016, 7, 289-308. | 1.7 | 38 |
| 12 | The genus <i>Thelonectria</i> (Nectriaceae, Hypocreales, Ascomycota) and closely related species with cylindrocarpon-like asexual states. <i>Fungal Diversity</i> , 2016, 80, 411-455. | 4.7 | 19 |
| 13 | Unraveling <i>Trichoderma</i> species in the attine ant environment: description of three new taxa. <i>Antonie Van Leeuwenhoek</i> , 2016, 109, 633-651. | 0.7 | 37 |
| 14 | Peptaibol, Secondary Metabolite, and Hydrophobin Pattern of Commercial Biocontrol Agents Formulated with Species of the <i>Trichoderma harzianum</i> Complex. <i>Chemistry and Biodiversity</i> , 2015, 12, 662-684. | 1.0 | 57 |
| 15 | Unexpected diversity of basidiomycetous endophytes in sapwood and leaves of <i>Hevea</i> . <i>Mycologia</i> , 2015, 107, 284-297. | 0.8 | 73 |
| 16 | Wild trees in the Amazon basin harbor a great diversity of beneficial endosymbiotic fungi: is this evidence of protective mutualism?. <i>Fungal Ecology</i> , 2015, 17, 18-29. | 0.7 | 44 |
| 17 | <i>Trichoderma asperellum</i> : A Dominant Endophyte Species in Cacao Grown in Sulawesi with Potential for Controlling Vascular Streak Dieback Disease. <i>Tropical Plant Pathology</i> , 2015, 40, 19-25. | 0.8 | 24 |
| 18 | Systematics of the <i>Trichoderma harzianum</i> species complex and the re-identification of commercial biocontrol strains. <i>Mycologia</i> , 2015, 107, 558-590. | 0.8 | 245 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Phylogeny and taxonomic revision of <i>Thelonectria discophora</i> (Ascomycota, Hypocreales, Nectriaceae) species complex. <i>Fungal Diversity</i> , 2015, 70, 1-29. | 4.7 | 15 |
| 20 | Novel endophytic lineages of <i>Tolyposcladium</i> provide new insights into the ecology and evolution of Cordyceps-like fungi. <i>Mycologia</i> , 2014, 106, 1090-1105. | 0.8 | 33 |
| 21 | Endophytic fungi from Peruvian highland and lowland habitats form distinctive and host plant-specific assemblages. <i>Biodiversity and Conservation</i> , 2013, 22, 999-1016. | 1.2 | 29 |
| 22 | EVOLUTION OF HABITAT PREFERENCE AND NUTRITION MODE IN A COSMOPOLITAN FUNGAL GENUS WITH EVIDENCE OF INTERKINGDOM HOST JUMPS AND MAJOR SHIFTS IN ECOLOGY. <i>Evolution; International Journal of Organic Evolution</i> , 2013, 67, n/a-n/a. | 1.1 | 75 |
| 23 | Genera in Bionectriaceae, Hypocreaceae, and Nectriaceae (Hypocreales) proposed for acceptance or rejection. <i>IMA Fungus</i> , 2013, 4, 41-51. | 1.7 | 121 |
| 24 | Multigene phylogenetic analyses of the <i>Thelonectria coronata</i> and <i>T. veuillotiana</i> species complexes. <i>Mycologia</i> , 2012, 104, 1325-1350. | 0.8 | 15 |
| 25 | <i>Trichoderma stromaticum</i> and its overseas relatives. <i>Mycological Progress</i> , 2012, 11, 215-254. | 0.5 | 27 |
| 26 | Linking ex planta fungi with their endophytic stages: <i>Perisporiopsis</i> , a common leaf litter and soil fungus, is a frequent endophyte of <i>Hevea</i> spp. and other plants. <i>Fungal Ecology</i> , 2011, 4, 94-102. | 0.7 | 28 |
| 27 | <i>Trichoderma amazonicum</i> , a new endophytic species on <i>Hevea brasiliensis</i> and <i>H. guianensis</i> from the Amazon basin. <i>Mycologia</i> , 2011, 103, 139-151. | 0.8 | 79 |
| 28 | Species delimitation in fungal endophyte diversity studies and its implications in ecological and biogeographic inferences. <i>Molecular Ecology</i> , 2011, 20, 3001-3013. | 2.0 | 197 |
| 29 | <i>Perisporiopsis lateritia</i> , a new species on decaying leaves of <i>Hevea</i> spp. from the Amazon basin in Peru. <i>Mycotaxon</i> , 2010, 113, 163-169. | 0.1 | 10 |
| 30 | Diversity of fungal endophytes in leaves and stems of wild rubber trees (<i>Hevea brasiliensis</i>) in Peru. <i>Fungal Ecology</i> , 2010, 3, 240-254. | 0.7 | 267 |
| 31 | The <i>Trichoderma brevicompactum</i> clade: a separate lineage with new species, new peptaibiotics, and mycotoxins. <i>Mycological Progress</i> , 2008, 7, 177-219. | 0.5 | 136 |
| 32 | <i>Regiocrella</i> , a new entomopathogenic genus with a pycnidial anamorph and its phylogenetic placement in the Clavicipitaceae. <i>Mycologia</i> , 2005, 97, 1225-1237. | 0.8 | 25 |
| 33 | The genus <i>Podocrella</i> and its nematode-killing anamorph <i>Harposporium</i> . <i>Mycologia</i> , 2005, 97, 433-443. | 0.8 | 16 |
| 34 | Clarification of the host substrate of <i>Ascopolyporus</i> and description of <i>Ascopolyporus philodendrus</i> sp. nov.. <i>Mycologia</i> , 2005, 97, 710-717. | 0.8 | 13 |
| 35 | A new species of <i>Hypocrella</i> , <i>H. macrostroma</i> , and its phylogenetic relationships to other species with large stromata. <i>Mycological Research</i> , 2005, 109, 1268-1275. | 2.5 | 17 |
| 36 | The genus <i>Podocrella</i> and its nematode-killing anamorph <i>Harposporium</i> . <i>Mycologia</i> , 2005, 97, 433-443. | 0.8 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | <i>Regiocrella</i> , a new entomopathogenic genus with a pycnidial anamorph and its phylogenetic placement in the Clavicipitaceae. <i>Mycologia</i> , 2005, 97, 1225-1237. | 0.8 | 36 |
| 38 | <i>Hypocrea phyllostachydis</i> and its <i>Trichoderma</i> anamorph, a new bambusicolous species from France. <i>Mycological Progress</i> , 2004, 3, 29-36. | 0.5 | 8 |
| 39 | <i>Hypocrea</i> / <i>Trichoderma</i> species with pachybasium-like conidiophores: teleomorphs for <i>T. minutisporum</i> and <i>T. polysporum</i> and their newly discovered relatives. <i>Mycologia</i> , 2004, 96, 310-342. | 0.8 | 33 |
| 40 | Multilocus phylogenetic structure within the <i>Trichoderma harzianum</i> / <i>Hypocrea lixii</i> complex. <i>Molecular Phylogenetics and Evolution</i> , 2003, 27, 302-313. | 1.2 | 137 |
| 41 | <i>Hypocrea</i> / <i>Trichoderma</i> : species with conidiophore elongations and green conidia. <i>Mycologia</i> , 2003, 95, 1100-1140. | 0.8 | 80 |
| 42 | <i>Hypocrea</i> / <i>Trichoderma</i> : species with conidiophore elongations and green conidia. <i>Mycologia</i> , 2003, 95, 1100-40. | 0.8 | 27 |
| 43 | <i>Hypocrea lixii</i> , the teleomorph of <i>Trichoderma harzianum</i> . <i>Mycological Progress</i> , 2002, 1, 283-286. | 0.5 | 36 |
| 44 | Taxonomy and phylogenetic relationships of two species of <i>Hypocrea</i> with <i>Trichoderma</i> anamorphs. <i>Mycological Progress</i> , 2002, 1, 409-428. | 0.5 | 35 |
| 45 | <i>Hypocrea nigrovirens</i> , a new species with a gliocladium-like anamorph. <i>Mycologia</i> , 2001, 93, 758-763. | 0.8 | 4 |
| 46 | <i>Hypocrea virens</i> sp. nov., the teleomorph of <i>Trichoderma virens</i> . <i>Mycologia</i> , 2001, 93, 1113-1124. | 0.8 | 24 |
| 47 | <i>Hypocrea virens</i> sp. nov., the Teleomorph of <i>Trichoderma virens</i> . <i>Mycologia</i> , 2001, 93, 1113. | 0.8 | 29 |