

Anne Peasey

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1165256/publications.pdf>

Version: 2024-02-01

75
papers

5,063
citations

136740

32
h-index

91712

69
g-index

77
all docs

77
docs citations

77
times ranked

10579
citing authors

#	ARTICLE	IF	CITATIONS
1	The interleukin-6 receptor as a target for prevention of coronary heart disease: a mendelian randomisation analysis. <i>Lancet, The</i> , 2012, 379, 1214-1224.	6.3	886
2	HMG-coenzyme A reductase inhibition, type 2 diabetes, and bodyweight: evidence from genetic analysis and randomised trials. <i>Lancet, The</i> , 2015, 385, 351-361.	6.3	562
3	Association between alcohol and cardiovascular disease: Mendelian randomisation analysis based on individual participant data. <i>BMJ, The</i> , 2014, 349, g4164-g4164.	3.0	528
4	Vitamin D and mortality: meta-analysis of individual participant data from a large consortium of cohort studies from Europe and the United States. <i>BMJ, The</i> , 2014, 348, g3656-g3656.	3.0	363
5	Determinants of cardiovascular disease and other non-communicable diseases in Central and Eastern Europe: Rationale and design of the HAPIEE study. <i>BMC Public Health</i> , 2006, 6, 255.	1.2	269
6	Accessibility and use of urban green spaces, and cardiovascular health: findings from a Kaunas cohort study. <i>Environmental Health</i> , 2014, 13, 20.	1.7	225
7	Education and coronary heart disease: mendelian randomisation study. <i>BMJ: British Medical Journal</i> , 2017, 358, j3542.	2.4	191
8	Neighbourhood socioeconomic status and cardiovascular risk factors: a multilevel analysis of nine cities in the Czech Republic and Germany. <i>BMC Public Health</i> , 2007, 7, 255.	1.2	115
9	Leukocyte Telomere Length and All-Cause, Cardiovascular Disease, and Cancer Mortality: Results From Individual-Participant-Data Meta-Analysis of 2 Large Prospective Cohort Studies. <i>American Journal of Epidemiology</i> , 2017, 185, 1317-1326.	1.6	91
10	Dietary habits in three Central and Eastern European countries: the HAPIEE study. <i>BMC Public Health</i> , 2009, 9, 439.	1.2	88
11	Mediterranean diet score and total and cardiovascular mortality in Eastern Europe: the HAPIEE study. <i>European Journal of Nutrition</i> , 2017, 56, 421-429.	4.6	87
12	Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: a meta-analysis of individual participant data. <i>BMC Medicine</i> , 2015, 13, 300.	2.3	83
13	APOE polymorphism and its effect on plasma C-reactive protein levels in a large general population sample. <i>Human Immunology</i> , 2010, 71, 304-308.	1.2	63
14	The association of depressive symptoms with cardiovascular and all-cause mortality in Central and Eastern Europe: Prospective results of the HAPIEE study. <i>European Journal of Preventive Cardiology</i> , 2016, 23, 1839-1847.	0.8	62
15	The <i>FTO</i> Gene and Obesity in a Large Eastern European Population Sample: The HAPIEE Study. <i>Obesity</i> , 2008, 16, 2764-2766.	1.5	61
16	Socio-economic circumstances and food habits in Eastern, Central and Western European populations. <i>Public Health Nutrition</i> , 2011, 14, 678-687.	1.1	61
17	Alcohol, drinking pattern and all-cause, cardiovascular and alcohol-related mortality in Eastern Europe. <i>European Journal of Epidemiology</i> , 2016, 31, 21-30.	2.5	60
18	Self-rated health and all-cause and cause-specific mortality of older adults: Individual data meta-analysis of prospective cohort studies in the CHANCES Consortium. <i>Maturitas</i> , 2017, 103, 37-44.	1.0	58

#	ARTICLE	IF	CITATIONS
19	Socio-economic status over the life-course and depressive symptoms in men and women in Eastern Europe. <i>Journal of Affective Disorders</i> , 2008, 105, 125-136.	2.0	52
20	Life Course Socioeconomic Position and Mid-Late Life Cognitive Function in Eastern Europe. <i>Journals of Gerontology - Series B Psychological Sciences and Social Sciences</i> , 2014, 69, 470-481.	2.4	52
21	Psychometric properties and confirmatory factor analysis of the CASP-19, a measure of quality of life in early old age: the HAPIEE study. <i>Aging and Mental Health</i> , 2015, 19, 595-609.	1.5	51
22	Fruit and vegetable consumption and mortality in Eastern Europe: Longitudinal results from the Health, Alcohol and Psychosocial Factors in Eastern Europe study. <i>European Journal of Preventive Cardiology</i> , 2016, 23, 493-501.	0.8	49
23	Age at natural menopause in three Central and Eastern European urban populations: The HAPIEE study. <i>Maturitas</i> , 2013, 75, 87-93.	1.0	47
24	Childhood socioeconomic circumstances and adult height and leg length in central and eastern Europe. <i>Journal of Epidemiology and Community Health</i> , 2008, 62, 351-357.	2.0	45
25	Psychosocial and socioeconomic determinants of cardiovascular mortality in Eastern Europe: A multicentre prospective cohort study. <i>PLoS Medicine</i> , 2017, 14, e1002459.	3.9	40
26	A Farm to Fork Risk Assessment for the Use of Wastewater in Agriculture in Accra, Ghana. <i>PLoS ONE</i> , 2015, 10, e0142346.	1.1	40
27	The FTO gene polymorphism is associated with end-stage renal disease: two large independent case-control studies in a general population. <i>Nephrology Dialysis Transplantation</i> , 2012, 27, 1030-1035.	0.4	39
28	Healthy diet indicator and mortality in Eastern European populations: prospective evidence from the HAPIEE cohort. <i>European Journal of Clinical Nutrition</i> , 2014, 68, 1346-1352.	1.3	38
29	Alcohol consumption, drinking patterns, and cognitive function in older Eastern European adults. <i>Neurology</i> , 2015, 84, 287-295.	1.5	38
30	Education, marital status, and risk of hip fractures in older men and women: the CHANCES project. <i>Osteoporosis International</i> , 2015, 26, 1733-1746.	1.3	38
31	Socioeconomic inequalities in all-cause mortality in the Czech Republic, Russia, Poland and Lithuania in the 2000s: findings from the HAPIEE Study. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 297-303.	2.0	37
32	Diabetes prevalence, awareness and treatment and their correlates in older persons in urban and rural population in the Astana region, Kazakhstan. <i>Diabetes Research and Clinical Practice</i> , 2016, 112, 6-12.	1.1	37
33	Binge Drinking and Blood Pressure: Cross-Sectional Results of the HAPIEE Study. <i>PLoS ONE</i> , 2013, 8, e65856.	1.1	33
34	Risk Perceptions of Wastewater Use for Urban Agriculture in Accra, Ghana. <i>PLoS ONE</i> , 2016, 11, e0150603.	1.1	33
35	Antioxidant vitamin intake and mortality in three Central and Eastern European urban populations: the HAPIEE study. <i>European Journal of Nutrition</i> , 2016, 55, 547-560.	1.8	32
36	Depressive symptoms and levels of C-reactive protein. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009, 44, 217-222.	1.6	29

#	ARTICLE	IF	CITATIONS
37	MLXIPL variant in individuals with low and high triglyceridemia in white population in Central Europe. <i>Human Genetics</i> , 2008, 124, 553-555.	1.8	28
38	Health Factors and Risk of All-Cause, Cardiovascular, and Coronary Heart Disease Mortality: Findings from the MONICA and HAPIEE Studies in Lithuania. <i>PLoS ONE</i> , 2014, 9, e114283.	1.1	27
39	Alcohol consumption and cognitive performance: a Mendelian randomization study. <i>Addiction</i> , 2014, 109, 1462-1471.	1.7	27
40	Alcohol consumption and physical functioning among middle-aged and older adults in Central and Eastern Europe: Results from the HAPIEE study. <i>Age and Ageing</i> , 2015, 44, 84-89.	0.7	26
41	The prospective relationship between social cohesion and depressive symptoms among older adults from Central and Eastern Europe. <i>Journal of Epidemiology and Community Health</i> , 2019, 73, 117-122.	2.0	24
42	Serum folate, vitamin B-12 and cognitive function in middle and older age: The HAPIEE study. <i>Experimental Gerontology</i> , 2016, 76, 33-38.	1.2	23
43	Work Stress and Subsequent Risk of Internet Addiction Among Information Technology Engineers in Taiwan. <i>Cyberpsychology, Behavior, and Social Networking</i> , 2014, 17, 542-550.	2.1	22
44	Blood-Based Oxidative Stress Markers and Cognitive Performance in Early Old Age: The HAPIEE Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2016, 42, 297-309.	0.7	20
45	A faecal exposure assessment of farm workers in Accra, Ghana: a cross sectional study. <i>BMC Public Health</i> , 2016, 16, 587.	1.2	20
46	Traditional Eastern European diet and mortality: prospective evidence from the HAPIEE study. <i>European Journal of Nutrition</i> , 2021, 60, 1091-1100.	1.8	19
47	Development and validation of two SCORE-based cardiovascular risk prediction models for Eastern Europe: a multicohort study. <i>European Heart Journal</i> , 2020, 41, 3325-3333.	1.0	17
48	Prevalence, awareness, treatment and control of dyslipidemia in older persons in urban and rural population in the Astana region, Kazakhstan. <i>BMC Public Health</i> , 2017, 17, 651.	1.2	16
49	DO LIPIDS CONTRIBUTE TO THE LACK OF CARDIO-PROTECTIVE EFFECT OF BINGE DRINKING: ALCOHOL CONSUMPTION AND LIPIDS IN THREE EASTERN EUROPEAN COUNTRIES. <i>Alcohol and Alcoholism</i> , 2005, 40, 431-435.	0.9	14
50	Association between Year of Birth and Cognitive Functions in Russia and the Czech Republic: Cross-Sectional Results of the HAPIEE Study. <i>Neuroepidemiology</i> , 2009, 33, 231-239.	1.1	12
51	Fruit, vegetable intake and blood pressure trajectories in older age. <i>Journal of Human Hypertension</i> , 2019, 33, 671-678.	1.0	12
52	Psychological well-being and mortality: longitudinal findings from Lithuanian middle-aged and older adults study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2019, 54, 803-811.	1.6	12
53	Impact of perceived control on all-cause and cardiovascular disease mortality in three urban populations of Central and Eastern Europe: the HAPIEE study. <i>Journal of Epidemiology and Community Health</i> , 2017, 71, 771-778.	2.0	11
54	Non-fatal injuries in three Central and Eastern European urban population samples: the HAPIEE study. <i>European Journal of Public Health</i> , 2010, 20, 695-701.	0.1	10

#	ARTICLE	IF	CITATIONS
55	Risk of depressive symptoms before and after the first hospitalisation for cancer: Evidence from a 16-year cohort study in the Czech Republic. <i>Journal of Affective Disorders</i> , 2020, 276, 76-83.	2.0	10
56	Trends in Prevalence of Dyslipidaemias and the Risk of Mortality in Lithuanian Urban Population Aged 45-64 in Relation to the Presence of the Dyslipidaemias and the Other Cardiovascular Risk Factors. <i>PLoS ONE</i> , 2014, 9, e100158.	1.1	10
57	<i>ADH1B</i> Polymorphism, Alcohol Consumption, and Binge Drinking in Slavic Caucasians: Results from the Czech HAPIEE Study. <i>Alcoholism: Clinical and Experimental Research</i> , 2012, 36, 900-905.	1.4	9
58	Lack of an association between left-handedness and <i>APOE</i> polymorphism in a large sample of adults: Results of the Czech HAPIEE study. <i>Laterality</i> , 2013, 18, 513-519.	0.5	9
59	Correlates of depressive symptoms in urban middle-aged and elderly Lithuanians. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014, 49, 1199-1207.	1.6	9
60	Socioeconomic circumstances, health behaviours and functional limitations in older persons in four Central and Eastern European populations. <i>Age and Ageing</i> , 2012, 41, 728-735.	0.7	8
61	Alcohol Consumption and Longitudinal Trajectories of Physical Functioning in Central and Eastern Europe: A 10-Year Follow-up of HAPIEE Study. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 1063-1068.	1.7	8
62	The association between the <i>FTO</i> gene variant and alcohol consumption and binge and problem drinking in different gene-environment background: The HAPIEE study. <i>Gene</i> , 2019, 707, 30-35.	1.0	8
63	FAT MASS AND OBESITY-ASSOCIATED (<i>FTO</i>) GENE AND ALCOHOL INTAKE. <i>Addiction</i> , 2012, 107, 1185-1186.	1.7	7
64	The association between <i>APOA5</i> haplotypes and plasma lipids is not modified by energy or fat intake: The Czech HAPIEE study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014, 24, 243-247.	1.1	7
65	Effort-reward imbalance at work, over-commitment personality and diet quality in Central and Eastern European populations. <i>British Journal of Nutrition</i> , 2016, 115, 1254-1264.	1.2	7
66	Education, material condition and physical functioning trajectories in middle-aged and older adults in Central and Eastern Europe: a cross-country comparison. <i>Journal of Epidemiology and Community Health</i> , 2016, 70, 1128-1135.	2.0	7
67	Mediterranean diet and physical functioning trajectories in Eastern Europe: Findings from the HAPIEE study. <i>PLoS ONE</i> , 2018, 13, e0200460.	1.1	7
68	Anthropometric trends and the risk of cardiovascular disease mortality in a Lithuanian urban population aged 45-64 years. <i>Scandinavian Journal of Public Health</i> , 2015, 43, 882-889.	1.2	6
69	Apolipoprotein E Arg136-Cys mutation and hyperlipidemia in a large central European population sample. <i>Clinica Chimica Acta</i> , 2008, 388, 217-218.	0.5	5
70	Dose-response association between physical activity and metabolic syndrome. <i>Open Medicine (Poland)</i> , 2013, 8, 273-282.	0.6	5
71	The relationship between body mass index and 10-year trajectories of physical functioning in middle-aged and older Russians: Prospective results of the Russian HAPIEE study. <i>Journal of Nutrition, Health and Aging</i> , 2017, 21, 381-388.	1.5	5
72	Social Patterning in Grip Strength, Chair Rise, and Walk Speed in an Aging Population: The Czech HAPIEE Study. <i>Journal of Aging and Physical Activity</i> , 2015, 23, 264-271.	0.5	4

#	ARTICLE	IF	CITATIONS
73	Lack of Association between NYD-SP18 Variant and Obesity. The Health Alcohol and Psychosocial Factors in Eastern Europe Study. <i>Annals of Nutrition and Metabolism</i> , 2016, 68, 244-248.	1.0	2
74	Longitudinal trajectories of blood lipid levels in an ageing population sample of Russian Western-Siberian urban population. <i>PLoS ONE</i> , 2021, 16, e0260229.	1.1	1
75	Social Patterning in Grip Strength, Chair Rise, and Walk Speed in an Aging Population: The Czech HAPIEE Study. <i>Journal of Aging and Physical Activity</i> , 2015, 23, 264-271.	0.5	0