

Christian H Lindh

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1166128/christian-h-lindh-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

200
papers

5,342
citations

40
h-index

64
g-index

215
ext. papers

6,692
ext. citations

6
avg. IF

5.84
L-index

#	Paper	IF	Citations
200	Arsenic exposure and biomarkers for oxidative stress and telomere length in indigenous populations in Bolivia.. <i>Ecotoxicology and Environmental Safety</i> , 2022 , 231, 113194	7	0
199	Perfluoroalkyl substances influence DNA methylation in school-age children highly exposed through drinking water contaminated from firefighting foam: a cohort study in Ronneby, Sweden.. <i>Environmental Epigenetics</i> , 2022 , 8, dvac004	2.4	0
198	From cohorts to molecules: Adverse impacts of endocrine disrupting mixtures.. <i>Science</i> , 2022 , 375, eabe8344	33.4	12
197	A Mixture of Endocrine Disrupting Chemicals Associated with Lower Birth Weight in Children Induces Adipogenesis and DNA Methylation Changes in Human Mesenchymal Stem Cells.. <i>International Journal of Molecular Sciences</i> , 2022 , 23,	6.3	1
196	Underground emissions and miners' personal exposure to diesel and renewable diesel exhaust in a Swedish iron ore mine.. <i>International Archives of Occupational and Environmental Health</i> , 2022 , 1	3.2	
195	Per- and Polyfluoroalkyl Substances and Risk of Myocardial Infarction and Stroke: A Nested Case-Control Study in Sweden.. <i>Environmental Health Perspectives</i> , 2022 , 130, 37007	8.4	1
194	Per- and polyfluoroalkyl substances (PFAS) and male reproductive function in young adulthood; a cross-sectional study.. <i>Environmental Research</i> , 2022 , 212, 113157	7.9	1
193	Determinants of serum half-lives for linear and branched perfluoroalkyl substances after long-term high exposure-A study in Ronneby, Sweden.. <i>Environment International</i> , 2022 , 163, 107198	12.9	3
192	Pyrimethanil and chlorpyrifos air concentrations and pregnant women's urinary metabolites in the Infants' Environmental Health Study (ISA), Costa Rica. <i>Environment International</i> , 2022 , 107328	12.9	0
191	Follicular fluid and blood levels of persistent organic pollutants and reproductive outcomes among women undergoing assisted reproductive technologies.. <i>Environmental Research</i> , 2021 , 112626	7.9	3
190	Exposure to common-use pesticides, manganese, lead, and thyroid function among pregnant women from the Infants' Environmental Health (ISA) study, Costa Rica. <i>Science of the Total Environment</i> , 2021 , 810, 151288	10.2	0
189	Associations between perfluoroalkyl substances and thyroid hormones after high exposure through drinking water. <i>Environmental Research</i> , 2021 , 194, 110647	7.9	4
188	Prenatal exposure to bisphenols and cognitive function in children at 7 years of age in the Swedish SELMA study. <i>Environment International</i> , 2021 , 150, 106433	12.9	6
187	Associations of prenatal exposure to phthalates and one phthalate substitute with anthropometric measures in early life: Results from the German LIFE Child cohort study. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2021 , 35, 101532	6.5	0
186	Prenatal exposures to mixtures of endocrine disrupting chemicals and children's weight trajectory up to age 5.5 in the SELMA study. <i>Scientific Reports</i> , 2021 , 11, 11036	4.9	4
185	RUBIC (ReproUnion Biobank and Infertility Cohort): A binational clinical foundation to study risk factors, life course, and treatment of infertility and infertility-related morbidity. <i>Andrology</i> , 2021 , 9, 1828-1842	4.2	3
184	Biomarkers after Controlled Inhalation Exposure to Exhaust from Hydrogenated Vegetable Oil (HVO). <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	4

183	Low-level exposure to polycyclic aromatic hydrocarbons is associated with reduced lung function among Swedish young adults. <i>Environmental Research</i> , 2021 , 197, 111169	7.9	2
182	Exposure to environmental phthalates during preschool age and obesity from childhood to young adulthood. <i>Environmental Research</i> , 2021 , 192, 110249	7.9	5
181	Human experimental exposure to glyphosate and biomonitoring of young Swedish adults. <i>International Journal of Hygiene and Environmental Health</i> , 2021 , 231, 113657	6.9	11
180	Urinary Bisphenol A, F and S Levels and Semen Quality in Young Adult Danish Men. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	6
179	Serum perfluoroalkyl substances in residents following long-term drinking water contamination from firefighting foam in Ronneby, Sweden. <i>Environment International</i> , 2021 , 147, 106333	12.9	14
178	Exposure to perfluoroalkyl substances in early pregnancy and risk of sporadic first trimester miscarriage. <i>Scientific Reports</i> , 2021 , 11, 3568	4.9	5
177	Prenatal phthalate exposure and early childhood wheeze in the SELMA study. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2021 ,	6.7	4
176	Temporal trends, 2000-2017, of perfluoroalkyl acid (PFAA) concentrations in serum of Swedish adolescents. <i>Environment International</i> , 2021 , 155, 106716	12.9	6
175	Persistent organic pollutants and the size of ovarian reserve in reproductive-aged women. <i>Environment International</i> , 2021 , 155, 106589	12.9	7
174	DNA methylation at GRIN2B partially mediates the association between prenatal bisphenol F exposure and cognitive functions in 7-year-old children in the SELMA study. <i>Environment International</i> , 2021 , 156, 106617	12.9	2
173	Mixtures of persistent organic pollutants are found in vital organs of late gestation human fetuses. <i>Chemosphere</i> , 2021 , 283, 131125	8.4	5
172	Association of phthalate exposure with thyroid function during pregnancy. <i>Environment International</i> , 2021 , 157, 106795	12.9	9
171	Filaggrin Polymorphisms and the Uptake of Chemicals through the Skin-A Human Experimental Study. <i>Environmental Health Perspectives</i> , 2021 , 129, 17002	8.4	6
170	Variability and predictors of weekly pesticide exposure in applicators from organic, sustainable and conventional smallholder farms in Costa Rica. <i>Occupational and Environmental Medicine</i> , 2020 , 77, 40-47	2.1	12
169	Long term transcriptional and behavioral effects in mice developmentally exposed to a mixture of endocrine disruptors associated with delayed human neurodevelopment. <i>Scientific Reports</i> , 2020 , 10, 9367	4.9	10
168	Persistent organic pollutants, pre-pregnancy use of combined oral contraceptives, age, and time-to-pregnancy in the SELMA cohort. <i>Environmental Health</i> , 2020 , 19, 67	6	10
167	Intrauterine exposure to perfluorinated compounds and overweight at age 4: A case-control study. <i>PLoS ONE</i> , 2020 , 15, e0230137	3.7	10
166	Associations between perfluoroalkyl substances and serum lipids in a Swedish adult population with contaminated drinking water. <i>Environmental Health</i> , 2020 , 19, 33	6	32

165	Per- and Polyfluoroalkyl Substances in Early Pregnancy and Risk for Preeclampsia: A Case-Control Study in Southern Sweden. <i>Toxics</i> , 2020 , 8,	4.7	10
164	A Probabilistic Approach to Evaluate the Risk of Decreased Total Triiodothyronine Hormone Levels following Chronic Exposure to PFOS and PFHxS via Contaminated Drinking Water. <i>Environmental Health Perspectives</i> , 2020 , 128, 76001	8.4	3
163	Pregnancy-induced changes in serum concentrations of perfluoroalkyl substances and the influence of kidney function. <i>Environmental Health</i> , 2020 , 19, 80	6	5
162	Prenatal pesticide exposure and respiratory health outcomes in the first year of life: Results from the infants' Environmental Health (ISA) study. <i>International Journal of Hygiene and Environmental Health</i> , 2020 , 225, 113474	6.9	7
161	Concentrations and temporal trends in pesticide biomarkers in urine of Swedish adolescents, 2000-2017. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2020 , 30, 756-767	6.7	17
160	Determinants of serum concentrations of perfluoroalkyl acids (PFAAs) in school children and the contribution of low-level PFAA-contaminated drinking water. <i>Environmental Sciences: Processes and Impacts</i> , 2020 , 22, 930-944	4.3	8
159	Association between serum concentrations of perfluoroalkyl substances (PFAS) and expression of serum microRNAs in a cohort highly exposed to PFAS from drinking water. <i>Environment International</i> , 2020 , 136, 105446	12.9	26
158	Fluorene exposure among PAH-exposed workers is associated with epigenetic markers related to lung cancer. <i>Occupational and Environmental Medicine</i> , 2020 , 77, 488-495	2.1	16
157	Maternal serum levels of perfluoroalkyl substances in early pregnancy and offspring birth weight. <i>Pediatric Research</i> , 2020 , 87, 1093-1099	3.2	36
156	Serum albumin adducts, DNA adducts and micronuclei frequency measured in benzo[a]pyrene-exposed mice for estimation of genotoxic potency. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2020 , 849, 503127	3	2
155	Inflammatory bowel disease and biomarkers of gut inflammation and permeability in a community with high exposure to perfluoroalkyl substances through drinking water. <i>Environmental Research</i> , 2020 , 181, 108923	7.9	18
154	Associations between serum concentrations of perfluoroalkyl substances and DNA methylation in women exposed through drinking water: A pilot study in Ronneby, Sweden. <i>Environment International</i> , 2020 , 145, 106148	12.9	7
153	Serum Half-Lives for Short- and Long-Chain Perfluoroalkyl Acids after Ceasing Exposure from Drinking Water Contaminated by Firefighting Foam. <i>Environmental Health Perspectives</i> , 2020 , 128, 77004	8.4	66
152	Exposure to polychlorinated compounds and cryptorchidism; A nested case-control study. <i>PLoS ONE</i> , 2020 , 15, e0236394	3.7	6
151	Contrasting prenatal nutrition and environmental exposures in association with birth weight and cognitive function in children at 7 years. <i>BMJ Nutrition, Prevention and Health</i> , 2020 , 3, 162-171	6.7	1
150	Fetal Programming of Semen Quality (FEPOS) Cohort - A DNBC Male-Offspring Cohort. <i>Clinical Epidemiology</i> , 2020 , 12, 757-770	5.9	4
149	Pesticide exposure among Bolivian farmers: associations between worker protection and exposure biomarkers. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2020 , 30, 730-742	6.7	20
148	Early prenatal exposure to suspected endocrine disruptor mixtures is associated with lower IQ at age seven. <i>Environment International</i> , 2020 , 134, 105185	12.9	63

147	Intrauterine exposure to perfluorinated compounds and overweight at age 4: A case-control study 2020 , 15, e0230137		
146	Intrauterine exposure to perfluorinated compounds and overweight at age 4: A case-control study 2020 , 15, e0230137		
145	Intrauterine exposure to perfluorinated compounds and overweight at age 4: A case-control study 2020 , 15, e0230137		
144	Intrauterine exposure to perfluorinated compounds and overweight at age 4: A case-control study 2020 , 15, e0230137		
143	Exposure to polychlorinated compounds and cryptorchidism; A nested case-control study 2020 , 15, e0236394		
142	Exposure to polychlorinated compounds and cryptorchidism; A nested case-control study 2020 , 15, e0236394		
141	Exposure to polychlorinated compounds and cryptorchidism; A nested case-control study 2020 , 15, e0236394		
140	Exposure to polychlorinated compounds and cryptorchidism; A nested case-control study 2020 , 15, e0236394		
139	Association of urinary bisphenols and triclosan with thyroid function during early pregnancy. <i>Environment International</i> , 2019 , 133, 105123	12.9	28
138	Concentrations of perfluoroalkyl substances (PFASs) in human embryonic and fetal organs from first, second, and third trimester pregnancies. <i>Environment International</i> , 2019 , 124, 482-492	12.9	93
137	Cardiovascular disease-related serum proteins in workers occupationally exposed to polycyclic aromatic hydrocarbons. <i>Toxicological Sciences</i> , 2019 ,	4.4	11
136	A Novel Approach to Chemical Mixture Risk Assessment-Linking Data from Population-Based Epidemiology and Experimental Animal Tests. <i>Risk Analysis</i> , 2019 , 39, 2259-2271	3.9	15
135	Caregiving and infants' neurodevelopment in rural Costa Rica: Results from the Infants' Environmental Health Study (ISA). <i>NeuroToxicology</i> , 2019 , 74, 100-107	4.4	4
134	Cancer-related proteins in serum are altered in workers occupationally exposed to polycyclic aromatic hydrocarbons: a cross-sectional study. <i>Carcinogenesis</i> , 2019 , 40, 771-781	4.6	6
133	Bisphenol A in human saliva and urine before and after treatment with dental polymer-based restorative materials. <i>European Journal of Oral Sciences</i> , 2019 , 127, 435-444	2.3	13
132	Filaggrin variations are associated with PAH metabolites in urine and DNA alterations in blood. <i>Environmental Research</i> , 2019 , 177, 108600	7.9	6
131	Early pregnancy serum levels of perfluoroalkyl substances and risk of preeclampsia in Swedish women. <i>Scientific Reports</i> , 2019 , 9, 9179	4.9	34
130	Exposure to Pesticides and Health Effects on Farm Owners and Workers From Conventional and Organic Agricultural Farms in Costa Rica: Protocol for a Cross-Sectional Study. <i>JMIR Research Protocols</i> , 2019 , 8, e10914	2	21

129	Environmental exposures in young adults with declining kidney function in a population at risk of Mesoamerican nephropathy. <i>Occupational and Environmental Medicine</i> , 2019 , 76, 920-926	2.1	14
128	Biomarkers of Exposure to Pyrimethanil After Controlled Human Experiments. <i>Journal of Analytical Toxicology</i> , 2019 , 43, 277-283	2.9	7
127	PVC flooring at home and uptake of phthalates in pregnant women. <i>Indoor Air</i> , 2019 , 29, 43-54	5.4	18
126	Temporal trends of phthalate exposures during 2007-2010 in Swedish pregnant women. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2018 , 28, 437-447	6.7	37
125	Prenatal phthalate exposure was associated with croup in Swedish infants. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2018 , 107, 1011-1019	3.1	13
124	Exposure to perfluoroalkyl substances during pregnancy and child behaviour at 5 to 9years of age. <i>Hormones and Behavior</i> , 2018 , 101, 105-112	3.7	13
123	DNA methylation of the cancer-related genes F2RL3 and AHRR is associated with occupational exposure to polycyclic aromatic hydrocarbons. <i>Carcinogenesis</i> , 2018 , 39, 869-878	4.6	19
122	Brominated Flame Retardants and Organophosphate Esters in Preschool Dust and Children's Hand Wipes. <i>Environmental Science & Technology</i> , 2018 , 52, 4878-4888	10.3	61
121	Prenatal Bisphenol A Exposure is Linked to Epigenetic Changes in Glutamate Receptor Subunit Gene Grin2b in Female Rats and Humans. <i>Scientific Reports</i> , 2018 , 8, 11315	4.9	24
120	Assessment of dermal uptake of diphenylmethane-4,4'-diisocyanate using tape stripping and biological monitoring. <i>European Journal of Dermatology</i> , 2018 , 28, 143-148	0.8	1
119	Amniotic Fluid INSL3 Measured During the Critical Time Window in Human Pregnancy Relates to Cryptorchidism, Hypospadias, and Phthalate Load: A Large Case-Control Study. <i>Frontiers in Physiology</i> , 2018 , 9, 406	4.6	25
118	Occupational exposure to asphalt mixture during road paving is related to increased mitochondria DNA copy number: a cross-sectional study. <i>Environmental Health</i> , 2018 , 17, 29	6	24
117	Comprehensive proteome analysis of nasal lavage samples after controlled exposure to welding nanoparticles shows an induced acute phase and a nuclear receptor, LXR/RXR, activation that influence the status of the extracellular matrix. <i>Clinical Proteomics</i> , 2018 , 15, 20	5	3
116	Half-lives of PFOS, PFHxS and PFOA after end of exposure to contaminated drinking water. <i>Occupational and Environmental Medicine</i> , 2018 , 75, 46-51	2.1	266
115	Excretion of Urinary Metabolites of the Phthalate Esters DEP and DEHP in 16 Volunteers after Inhalation and Dermal Exposure. <i>International Journal of Environmental Research and Public Health</i> , 2018 , 15,	4.6	16
114	Association between paternal smoking at the time of pregnancy and the semen quality in sons. <i>PLoS ONE</i> , 2018 , 13, e0207221	3.7	3
113	Temporal trends and predictors of perfluoroalkyl substances serum levels in Swedish pregnant women in the SELMA study. <i>PLoS ONE</i> , 2018 , 13, e0209255	3.7	32
112	Prenatal Mancozeb Exposure, Excess Manganese, and Neurodevelopment at 1 Year of Age in the Infants' Environmental Health (ISA) Study. <i>Environmental Health Perspectives</i> , 2018 , 126, 057007	8.4	34

111	Inhalation and Dermal Uptake of Particle and Gas-Phase Phthalates-A Human Exposure Study. <i>Environmental Science & Technology</i> , 2018 , 52, 12792-12800	10.3	23
110	Association of Prenatal Phthalate Exposure With Language Development in Early Childhood. <i>JAMA Pediatrics</i> , 2018 , 172, 1169-1176	8.3	34
109	Incorporating regulatory guideline values in analysis of epidemiology data. <i>Environment International</i> , 2018 , 120, 535-543	12.9	4
108	Daily intake of phthalates, MEHP, and DINCH by ingestion and inhalation. <i>Chemosphere</i> , 2018 , 208, 40-49	8.4	39
107	Environmental tobacco smoke exposure during pregnancy has limited effect on infant birthweight and umbilical vein endothelial nitric oxide synthase. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018 , 97, 1309-1316	3.8	5
106	Phthalates, non-phthalate plasticizers and bisphenols in Swedish preschool dust in relation to children's exposure. <i>Environment International</i> , 2017 , 102, 114-124	12.9	123
105	Concentration of perfluorinated compounds and cotinine in human foetal organs, placenta, and maternal plasma. <i>Science of the Total Environment</i> , 2017 , 596-597, 97-105	10.2	63
104	Diverging temporal trends of human exposure to bisphenols and plastizisers, such as phthalates, caused by substitution of legacy EDCs?. <i>Environmental Research</i> , 2017 , 153, 48-54	7.9	66
103	High levels of vitamin D associated with less ischemic heart disease - a nested case-control study among rural men in Sweden. <i>Annals of Agricultural and Environmental Medicine</i> , 2017 , 24, 288-293	1.4	5
102	Early markers of cardiovascular disease are associated with occupational exposure to polycyclic aromatic hydrocarbons. <i>Scientific Reports</i> , 2017 , 7, 9426	4.9	50
101	Female exposure to phthalates and time to pregnancy: a first pregnancy planner study. <i>Human Reproduction</i> , 2017 , 32, 232-238	5.7	20
100	Urinary 8-oxo-7, 8-dihydro-2'-deoxyguanosine concentrations after frying of bacon, a pilot study. <i>Toxicological and Environmental Chemistry</i> , 2017 , 99, 525-534	1.4	2
99	Prenatal smoking exposure, measured as maternal serum cotinine, and children's motor developmental milestones and motor function: A follow-up study. <i>NeuroToxicology</i> , 2016 , 53, 236-245	4.4	10
98	Cotinine Validation of Self-Reported Smoking During Pregnancy in the Swedish Medical Birth Register. <i>Nicotine and Tobacco Research</i> , 2016 , 18, 79-83	4.9	41
97	Prenatal Phthalate, Perfluoroalkyl Acid, and Organochlorine Exposures and Term Birth Weight in Three Birth Cohorts: Multi-Pollutant Models Based on Elastic Net Regression. <i>Environmental Health Perspectives</i> , 2016 , 124, 365-72	8.4	140
96	Perfluorooctane Sulfonate Concentrations in Amniotic Fluid, Biomarkers of Fetal Leydig Cell Function, and Cryptorchidism and Hypospadias in Danish Boys (1980-1996). <i>Environmental Health Perspectives</i> , 2016 , 124, 151-6	8.4	33
95	Manganese concentrations in drinking water from villages near banana plantations with aerial mancozeb spraying in Costa Rica: Results from the Infants' Environmental Health Study (ISA). <i>Environmental Pollution</i> , 2016 , 215, 247-257	9.3	35
94	Analysis of nanoparticle-protein coronas formed in vitro between nanosized welding particles and nasal lavage proteins. <i>Nanotoxicology</i> , 2016 , 10, 226-34	5.3	22

93	Pesticide exposure and neurodevelopment in children aged 6-9 years from Talamanca, Costa Rica. <i>Cortex</i> , 2016 , 85, 137-150	3.8	71
92	Pregnancy serum concentrations of perfluorinated alkyl substances and offspring behaviour and motor development at age 5-9 years--a prospective study. <i>Environmental Health</i> , 2015 , 14, 2	6	47
91	Is the fact of parenting couples cohabitation affecting the serum levels of persistent organohalogen pollutants?. <i>International Journal of Hygiene and Environmental Health</i> , 2015 , 218, 392-400	6.9	6
90	Prenatal phthalate exposures and anogenital distance in Swedish boys. <i>Environmental Health Perspectives</i> , 2015 , 123, 101-7	8.4	179
89	Perfluorinated chemicals in blood serum of inhabitants in central Poland in relation to gender and age. <i>Science of the Total Environment</i> , 2015 , 532, 548-55	10.2	34
88	Phthalates, perfluoroalkyl acids, metals and organochlorines and reproductive function: a multipollutant assessment in Greenlandic, Polish and Ukrainian men. <i>Occupational and Environmental Medicine</i> , 2015 , 72, 385-93	2.1	52
87	Amniotic fluid phthalate levels and male fetal gonad function. <i>Epidemiology</i> , 2015 , 26, 91-9	3.1	75
86	Prenatal phthalate exposure and reproductive function in young men. <i>Environmental Research</i> , 2015 , 138, 264-70	7.9	62
85	Phthalate exposure and reproductive parameters in young men from the general Swedish population. <i>Environment International</i> , 2015 , 85, 54-60	12.9	61
84	Tobacco smoking during pregnancy and risk of adverse behaviour in offspring: A follow-up study. <i>Reproductive Toxicology</i> , 2015 , 58, 65-72	3.4	6
83	Toluene diisocyanate: Induction of the autotaxin-lysophosphatidic acid axis and its association with airways symptoms. <i>Toxicology and Applied Pharmacology</i> , 2015 , 287, 222-31	4.6	12
82	Exposure of hairdressers to ortho- and meta-toluidine in hair dyes. <i>Occupational and Environmental Medicine</i> , 2015 , 72, 57-63	2.1	18
81	Levels of perfluoroalkyl substances and risk of coronary heart disease: Findings from a population-based longitudinal study. <i>Environmental Research</i> , 2015 , 142, 148-54	7.9	26
80	Hazard quotient profiles used as a risk assessment tool for PFOS and PFOA serum levels in three distinctive European populations. <i>Environment International</i> , 2015 , 74, 112-8	12.9	42
79	Serum phthalate levels and time to pregnancy in couples from Greenland, Poland and Ukraine. <i>PLoS ONE</i> , 2015 , 10, e0120070	3.7	28
78	Vitamin D Status at Birth and Future Risk of Attention Deficit/Hyperactivity Disorder (ADHD). <i>PLoS ONE</i> , 2015 , 10, e0140164	3.7	19
77	Anthropometry in 5- to 9-Year-Old Greenlandic and Ukrainian Children in Relation to Prenatal Exposure to Perfluorinated Alkyl Substances. <i>Environmental Health Perspectives</i> , 2015 , 123, 841-6	8.4	65
76	Oxidative stress, telomere shortening, and DNA methylation in relation to low-to-moderate occupational exposure to welding fumes. <i>Environmental and Molecular Mutagenesis</i> , 2015 , 56, 684-93	3.2	47

75	LC-MS-MS Analysis of Urinary Biomarkers of Imazalil Following Experimental Exposures. <i>Journal of Analytical Toxicology</i> , 2015 , 39, 691-7	2.9	8
74	Targeted proteomic analyses of nasal lavage fluid in persulfate-challenged hairdressers with bleaching powder-associated rhinitis. <i>Journal of Proteome Research</i> , 2015 , 14, 860-73	5.6	6
73	Blood and hair manganese concentrations in pregnant women from the infants' environmental health study (ISA) in Costa Rica. <i>Environmental Science & Technology</i> , 2014 , 48, 3467-76	10.3	56
72	Time trends between 1987 and 2007 for perfluoroalkyl acids in plasma from Swedish women. <i>Chemosphere</i> , 2014 , 102, 61-7	8.4	32
71	Fetal exposure to perfluorinated compounds and attention deficit hyperactivity disorder in childhood. <i>PLoS ONE</i> , 2014 , 9, e95891	3.7	58
70	Exposure determinants of phthalates, parabens, bisphenol A and triclosan in Swedish mothers and their children. <i>Environment International</i> , 2014 , 73, 323-33	12.9	198
69	Determination of 5-hydroxythiabendazole in human urine as a biomarker of exposure to thiabendazole using LC/MS/MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014 , 973C, 61-67	3.2	10
68	Aerial application of mancozeb and urinary ethylene thiourea (ETU) concentrations among pregnant women in Costa Rica: the Infants' Environmental Health Study (ISA). <i>Environmental Health Perspectives</i> , 2014 , 122, 1321-8	8.4	47
67	Exposure to perfluoroalkyl substances and sperm DNA global methylation in Arctic and European populations. <i>Environmental and Molecular Mutagenesis</i> , 2014 , 55, 591-600	3.2	33
66	Perfluoroalkyl substances and time to pregnancy in couples from Greenland, Poland and Ukraine. <i>Environmental Health</i> , 2014 , 13, 116	6	25
65	Associations between serum phthalates and biomarkers of reproductive function in 589 adult men. <i>Environment International</i> , 2014 , 66, 146-56	12.9	77
64	Biomarkers of exposure in Monday morning urine samples as a long-term measure of exposure to aromatic diisocyanates. <i>International Archives of Occupational and Environmental Health</i> , 2014 , 87, 365-72	3.2	6
63	Human biological monitoring of suspected endocrine-disrupting compounds. <i>Asian Journal of Andrology</i> , 2014 , 16, 5-16	2.8	34
62	Determinants of maternal and fetal exposure and temporal trends of perfluorinated compounds. <i>Environmental Science and Pollution Research</i> , 2013 , 20, 7970-8	5.1	69
61	High-throughput method for the analysis of ethylenethiourea with direct injection of hydrolysed urine using online on-column extraction liquid chromatography and triple quadrupole mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013 , 934, 53-9	3.2	18
60	Human and methodological sources of variability in the measurement of urinary 8-oxo-7,8-dihydro-2'-deoxyguanosine. <i>Antioxidants and Redox Signaling</i> , 2013 , 18, 2377-91	8.4	107
59	Screening method using selected reaction monitoring for targeted proteomics studies of nasal lavage fluid. <i>Journal of Proteome Research</i> , 2013 , 12, 234-47	5.6	15
58	Serum concentrations of perfluorinated alkyl acids and their associations with diet and personal characteristics among Swedish adults. <i>Molecular Nutrition and Food Research</i> , 2013 , 57, 2206-15	5.9	32

57	Sperm DNA integrity in relation to exposure to environmental perfluoroalkyl substances - a study of spouses of pregnant women in three geographical regions. <i>Reproductive Toxicology</i> , 2012 , 33, 577-583 ^{3,4}		58
56	Serum levels of perfluorinated compounds and sperm Y:X chromosome ratio in two European populations and in Inuit from Greenland. <i>Reproductive Toxicology</i> , 2012 , 34, 644-50	3.4	19
55	Early life low-level cadmium exposure is positively associated with increased oxidative stress. <i>Environmental Research</i> , 2012 , 112, 164-70	7.9	44
54	Indigenous children living nearby plantations with chlorpyrifos-treated bags have elevated 3,5,6-trichloro-2-pyridinol (TCPy) urinary concentrations. <i>Environmental Research</i> , 2012 , 117, 17-26	7.9	22
53	A nested case-control study of intrauterine exposure to persistent organochlorine pollutants and the risk of hypospadias. <i>PLoS ONE</i> , 2012 , 7, e44767	3.7	30
52	Blood serum concentrations of perfluorinated compounds in men from Greenlandic Inuit and European populations. <i>Chemosphere</i> , 2012 , 88, 1269-75	8.4	101
51	Very high correlations between fresh weight and lipid-adjusted PCB-153 serum concentrations: irrespective of fasting status, age, body mass index, gender, or exposure distributions. <i>Chemosphere</i> , 2012 , 88, 828-31	8.4	15
50	Association between pregnancy loss and urinary phthalate levels around the time of conception. <i>Environmental Health Perspectives</i> , 2012 , 120, 458-63	8.4	140
49	Phthalates and perfluorooctanesulfonic acid in human amniotic fluid: temporal trends and timing of amniocentesis in pregnancy. <i>Environmental Health Perspectives</i> , 2012 , 120, 897-903	8.4	88
48	Assessment of long-term and recent pesticide exposure among rural school children in Nicaragua. <i>Occupational and Environmental Medicine</i> , 2012 , 69, 119-25	2.1	15
47	Analysis of chlormequat in human urine as a biomarker of exposure using liquid chromatography triple quadrupole mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2011 , 879, 1551-6	3.2	19
46	Searching for early breast cancer biomarkers by serum protein profiling of pre-diagnostic serum; a nested case-control study. <i>BMC Cancer</i> , 2011 , 11, 381	4.8	66
45	Strategy for identification and detection of multiple oxidative modifications within proteins applied on persulfate-oxidized hemoglobin and human serum albumin. <i>Rapid Communications in Mass Spectrometry</i> , 2011 , 25, 327-40	2.2	6
44	N-nitrosamines are associated with shorter telomere length. <i>Scandinavian Journal of Work, Environment and Health</i> , 2011 , 37, 316-24	4.3	42
43	Time-dependent proteomic iTRAQ analysis of nasal lavage of hairdressers challenged by persulfate. <i>Journal of Proteome Research</i> , 2010 , 9, 5620-8	5.6	9
42	Low 8-oxo-7,8-dihydro-2'-deoxyguanosine levels and influence of genetic background in an Andean population exposed to high levels of arsenic. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2010 , 683, 98-105	3.3	22
41	Chronic exposure to cadmium and arsenic strongly influences concentrations of 8-oxo-7,8-dihydro-2'-deoxyguanosine in urine. <i>Free Radical Biology and Medicine</i> , 2010 , 48, 1211-7	7.8	68
40	Identification of covalent binding sites of ethyl 2-cyanoacrylate, methyl methacrylate and 2-hydroxyethyl methacrylate in human hemoglobin using LC/MS/MS techniques. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2010 , 878, 2474-82	3.2	2

39	A nested case-control study of intrauterine exposure to persistent organochlorine pollutants in relation to risk of type 1 diabetes. <i>PLoS ONE</i> , 2010 , 5, e11281	3.7	26
38	Acute health effects on planters of conifer seedlings treated with insecticides. <i>Annals of Occupational Hygiene</i> , 2009 , 53, 383-90		9
37	Does longstanding nicotine exposure impair bone healing and osseointegration? An experimental study in rabbits. <i>Journal of Biomedical Materials Research - Part B Applied Biomaterials</i> , 2009 , 91, 918-23	3.5	17
36	Methylhexahydrophthalic anhydride adducted albumin tryptic peptides in nasal lavage fluid. <i>Inhalation Toxicology</i> , 2009 , 21, 1013-20	2.7	4
35	N-nitrosamines in the southern Swedish rubber industries - exposure, health effects, and immunologic markers. <i>Scandinavian Journal of Work, Environment and Health</i> , 2009 , 35, 203-11	4.3	7
34	Underestimation of toluene diisocyanate concentration using long-term sampling with 1-(2-methoxyphenyl) piperazine impregnated filters. <i>International Journal of Hygiene and Environmental Health</i> , 2008 , 211, 458-62	6.9	3
33	Identification of covalent binding sites of phthalic anhydride in human hemoglobin. <i>Chemical Research in Toxicology</i> , 2008 , 21, 2156-63	4	4
32	Analysis of phenoxyacetic acid herbicides as biomarkers in human urine using liquid chromatography/triple quadrupole mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2008 , 22, 143-50	2.2	12
31	Analysis of ethylenethiourea as a biomarker in human urine using liquid chromatography/triple quadrupole mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2008 , 22, 2573-9	2.2	14
30	Evaluation of an immunoaffinity extraction column for enrichment of adducts between human serum albumin and hexahydrophthalic anhydride in plasma. <i>Biomedical Chromatography</i> , 2008 , 22, 327-32 ^{1.7}		4
29	Analysis of 3,5-dichloroaniline as a biomarker of vinclozolin and iprodione in human urine using liquid chromatography/triple quadrupole mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2007 , 21, 536-42	2.2	24
28	Semen quality and exposure to persistent organochlorine pollutants. <i>Epidemiology</i> , 2006 , 17, 450-8	3.1	104
27	Reproductive hormone levels in men exposed to persistent organohalogen pollutants: a study of Inuit and three European cohorts. <i>Environmental Health Perspectives</i> , 2006 , 114, 1348-53	8.4	51
26	Xenoestrogenic activity in blood of European and Inuit populations. <i>Environmental Health</i> , 2006 , 5, 12	6	36
25	Dioxin-like activities in serum across European and Inuit populations. <i>Environmental Health</i> , 2006 , 5, 14	6	32
24	Effect of systemic administration of nicotine on healing in osseous defects. An experimental study in rabbits. Part II. <i>Clinical Oral Implants Research</i> , 2006 , 17, 488-94	4.8	10
23	The impact of nicotine on osseointegration. An experimental study in the femur and tibia of rabbits. <i>Clinical Oral Implants Research</i> , 2005 , 16, 389-95	4.8	29
22	Inter-population variations in concentrations, determinants of and correlations between 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) and 1,1-dichloro-2,2-bis (p-chlorophenyl)-ethylene (p,p'-DDE): a cross-sectional study of 3161 men and women from Inuit and European populations. <i>Environmental Health</i> , 2005 , 4, 27	6	81

21	The impact of nicotine on bone healing and osseointegration. <i>Clinical Oral Implants Research</i> , 2005 , 16, 268-76	4.8	37
20	Biological monitoring of carbon disulphide and phthalate exposure in the contemporary rubber industry. <i>International Archives of Occupational and Environmental Health</i> , 2005 , 78, 663-9	3.2	18
19	Characterization of adducts formed between human serum albumin and the butadiene metabolite epoxybutanediol. <i>Rapid Communications in Mass Spectrometry</i> , 2005 , 19, 2488-96	2.2	12
18	Exposure to PCBs and p,p'-DDE and human sperm chromatin integrity. <i>Environmental Health Perspectives</i> , 2005 , 113, 175-9	8.4	79
17	A survey of airborne isocyanate exposure in 13 Swedish polyurethane industries. <i>Annals of Occupational Hygiene</i> , 2004 , 48, 405-14		14
16	Determination of isocyanates in air using 1-(2-methoxyphenyl)piperazine-impregnated filters: long-term sampling performance and field comparison with impingers with dibutylamine. <i>Annals of Occupational Hygiene</i> , 2004 , 48, 415-24		14
15	In vivo conjugation of nasal lavage proteins by hexahydrophthalic anhydride. <i>Toxicology and Applied Pharmacology</i> , 2004 , 194, 69-78	4.6	9
14	Correlations between air levels of hexahydrophthalic anhydride (HHPA) and HHPA-adducted albumin tryptic peptides in nasal lavage fluid from experimentally exposed volunteers. <i>Rapid Communications in Mass Spectrometry</i> , 2004 , 18, 1592-8	2.2	11
13	Biological monitoring of exposure to toluene diisocyanate. <i>Scandinavian Journal of Work, Environment and Health</i> , 2004 , 30, 371-8	4.3	20
12	Development, validation and characterization of an analytical method for the quantification of hydrolysable urinary metabolites and plasma protein adducts of 2,4- and 2,6-toluene diisocyanate, 1,5-naphthalene diisocyanate and 4,4'-methylenediphenyl diisocyanate. <i>Biomarkers</i> , 2003 , 8, 204-17	2.6	40
11	Determination of hexahydrophthalic anhydride adducts to human serum albumin. <i>Biomarkers</i> , 2003 , 8, 343-59	2.6	18
10	Exposure-response relationships for hexahydrophthalic and methylhexahydrophthalic anhydrides with total plasma protein adducts as biomarkers. <i>Scandinavian Journal of Work, Environment and Health</i> , 2003 , 29, 297-303	4.3	10
9	Mass spectrometric characterization of human hemoglobin adducts formed in vitro by hexahydrophthalic anhydride. <i>Chemical Research in Toxicology</i> , 2002 , 15, 562-9	4	11
8	Total plasma protein adducts of allergenic hexahydrophthalic and methylhexahydrophthalic anhydrides as biomarkers of long-term exposure. <i>Scandinavian Journal of Work, Environment and Health</i> , 2001 , 27, 133-9	4.3	6
7	Quantification of protein adducts of hexahydrophthalic anhydride and methylhexahydrophthalic anhydride in human plasma. <i>Journal of Environmental Monitoring</i> , 2000 , 2, 155-60		7
6	Binding of the potent allergen hexahydrophthalic anhydride in the mucosa of the upper respiratory and alimentary tract following single inhalation exposures in guinea pigs and rats. <i>Toxicology</i> , 1999 , 134, 153-68	4.4	8
5	Human hemoglobin adducts following exposure to hexahydrophthalic anhydride and methylhexahydrophthalic anhydride. <i>Toxicology and Applied Pharmacology</i> , 1998 , 153, 152-60	4.6	14
4	Haemoglobin adducts and specific immunoglobulin G in humans as biomarkers of exposure to hexahydrophthalic anhydride. <i>Biomarkers</i> , 1997 , 2, 239-46	2.6	10

- 3 Biological monitoring of methylhexahydrophthalic anhydride by determination of methylhexahydrophthalic acid in urine and plasma from exposed workers. *International Archives of Occupational and Environmental Health*, **1997**, 70, 128-32 3.2 9
- 2 Direct measurement of hexahydrophthalic anhydride in workplace air with a transportable Fourier transform infrared spectrometer. *AIHA Journal*, **1996**, 57, 832-6 8
- 1 Exposure to Pesticides and Health Effects on Farm Owners and Workers From Conventional and Organic Agricultural Farms in Costa Rica: Protocol for a Cross-Sectional Study (Preprint) 1