

Kazuo Komamura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11670766/publications.pdf>

Version: 2024-02-01

20
papers

2,574
citations

471371

17
h-index

752573

20
g-index

20
all docs

20
docs citations

20
times ranked

3193
citing authors

#	ARTICLE	IF	CITATIONS
1	Blockade of sarcolemmal TRPV2 accumulation inhibits progression of dilated cardiomyopathy. <i>Cardiovascular Research</i> , 2013, 99, 760-768.	1.8	70
2	A histamine H2 receptor blocker ameliorates development of heart failure in dogs independently of β -adrenergic receptor blockade. <i>Basic Research in Cardiology</i> , 2010, 105, 787-794.	2.5	38
3	Metformin Prevents Progression of Heart Failure in Dogs. <i>Circulation</i> , 2009, 119, 2568-2577.	1.6	269
4	Activation of Ecto-5'-Nucleotidase in the Blood and Hearts of Patients With Chronic Heart Failure. <i>Journal of Cardiac Failure</i> , 2008, 14, 426-430.	0.7	8
5	Activation of Na ⁺ /H ⁺ Exchanger 1 Is Sufficient to Generate Ca ²⁺ Signals That Induce Cardiac Hypertrophy and Heart Failure. <i>Circulation Research</i> , 2008, 103, 891-899.	2.0	168
6	Erythropoietin Enhances Neovascularization of Ischemic Myocardium and Improves Left Ventricular Dysfunction After Myocardial Infarction in Dogs. <i>Journal of the American College of Cardiology</i> , 2006, 48, 176-184.	1.2	123
7	Long-Term Stimulation of Adenosine A2b Receptors Begun After Myocardial Infarction Prevents Cardiac Remodeling in Rats. <i>Circulation</i> , 2006, 114, 1923-1932.	1.6	92
8	Erythropoietin Just Before Reperfusion Reduces Both Lethal Arrhythmias and Infarct Size via the Phosphatidylinositol-3 Kinase-Dependent Pathway in Canine Hearts. <i>Cardiovascular Drugs and Therapy</i> , 2005, 19, 33-40.	1.3	82
9	Exacerbation of heart failure in adiponectin-deficient mice due to impaired regulation of AMPK and glucose metabolism. <i>Cardiovascular Research</i> , 2005, 67, 705-713.	1.8	207
10	UGT1A1 haplotypes associated with reduced glucuronidation and increased serum bilirubin in irinotecan-administered Japanese patients with cancer*1. <i>Clinical Pharmacology and Therapeutics</i> , 2004, 75, 501-515.	2.3	243
11	A novel mechanism of myocyte degeneration involving the Ca ²⁺ -permeable growth factor-regulated channel. <i>Journal of Cell Biology</i> , 2003, 161, 957-967.	2.3	250
12	Comprehensive UGT1A1 Genotyping in a Japanese Population by Pyrosequencing. <i>Clinical Chemistry</i> , 2003, 49, 1182-1185.	1.5	55
13	Salutary effect of adjunctive intracoronary nicorandil administration on restoration of myocardial blood flow and functional improvement in patients with acute myocardial infarction. <i>American Heart Journal</i> , 1997, 133, 616-621.	1.2	94
14	Ecto-5'-Nucleotidase Mediates Infarct Size-Limiting Effect by Ischemic Preconditioning in the Rabbit Heart. <i>Journal of Cardiovascular Pharmacology</i> , 1997, 30, 775-783.	0.8	7
15	Plasma Adenosine Levels Increase in Patients With Chronic Heart Failure. <i>Circulation</i> , 1997, 95, 1363-1365.	1.6	187
16	Intracoronary morphology of culprit lesions after reperfusion in acute myocardial infarction: Serial angiographic observations. <i>Journal of the American College of Cardiology</i> , 1996, 27, 606-610.	1.2	106
17	Role of Activation of Protein Kinase C in the Infarct Size-Limiting Effect of Ischemic Preconditioning Through Activation of Ecto-5'-nucleotidase. <i>Circulation</i> , 1996, 93, 781-791.	1.6	100
18	752-4 Transient Intracoronary Infusion of ATP After Reperfusion Reduces the Extent of No-reflow and Infarct Size in Dogs. <i>Journal of the American College of Cardiology</i> , 1995, 25, 227A-228A.	1.2	6

#	ARTICLE	IF	CITATIONS
19	Progressive decreases in coronary vein flow during reperfusion in acute myocardial infarction: Clinical documentation of the no reflow phenomenon after successful thrombolysis. Journal of the American College of Cardiology, 1994, 24, 370-377.	1.2	70
20	Neointimal coverage of stents in human coronary arteries observed by angioscopy. Journal of the American College of Cardiology, 1994, 23, 341-346.	1.2	399