

Song Gao

List of Publications by Citations

Source: <https://exaly.com/author-pdf/11683162/song-gao-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

496
papers

32,036
citations

86
h-index

155
g-index

527
ext. papers

33,966
ext. citations

6.4
avg, IF

7.26
L-index

#	Paper	IF	Citations
496	Synthesis, functionalization, and biomedical applications of multifunctional magnetic nanoparticles. <i>Advanced Materials</i> , 2010 , 22, 2729-42	24	1129
495	A Stable Pentagonal Bipyramidal Dy(III) Single-Ion Magnet with a Record Magnetization Reversal Barrier over 1000 K. <i>Journal of the American Chemical Society</i> , 2016 , 138, 5441-50	16.4	738
494	An organometallic single-ion magnet. <i>Journal of the American Chemical Society</i> , 2011 , 133, 4730-3	16.4	645
493	Chemically diverse and multifunctional hybrid organic/inorganic perovskites. <i>Nature Reviews Materials</i> , 2017 , 2,	73.3	608
492	A mononuclear dysprosium complex featuring single-molecule-magnet behavior. <i>Angewandte Chemie - International Edition</i> , 2010 , 49, 7448-51	16.4	555
491	An organic-inorganic perovskite ferroelectric with large piezoelectric response. <i>Science</i> , 2017 , 357, 306-309	39.3	506
490	Strategies towards single-chain magnets. <i>Coordination Chemistry Reviews</i> , 2010 , 254, 1081-1100	23.2	488
489	3d-4f combined chemistry: synthetic strategies and magnetic properties. <i>Inorganic Chemistry</i> , 2009 , 48, 3342-59	5.1	472
488	Constructing magnetic molecular solids by employing three-atom ligands as bridges. <i>Chemical Communications</i> , 2008 , 281-94	5.8	467
487	An azide-bridged homospin single-chain magnet: [Co(2,2'-bithiazoline)(N ₃) ₂] _n . <i>Journal of the American Chemical Society</i> , 2003 , 125, 13976-7	16.4	466
486	Coexistence of magnetic and electric orderings in the metal-formate frameworks of [NH ₄][M(HCOO) ₃]. <i>Journal of the American Chemical Society</i> , 2011 , 133, 14948-51	16.4	413
485	Single-crystal dendritic micro-pines of magnetic alpha-Fe ₂ O ₃ : large-scale synthesis, formation mechanism, and properties. <i>Angewandte Chemie - International Edition</i> , 2005 , 44, 4197-201	16.4	407
484	Framework-structured weak ferromagnets. <i>Chemical Society Reviews</i> , 2011 , 40, 3157-81	58.5	361
483	From Cubane to Supercubane: The Design, Synthesis, and Structure of a Three-Dimensional Open Framework Based on a Ln(4)O(4) Cluster This work was supported by the National Natural Science Foundation of China (No. 29771001, 29831010), National Key Project for Fundamental Research (G1998061306), and the Excellent Young Teachers Fund of MOE, P.R. China. <i>Angewandte Chemie - International Edition</i> , 2004 , 43, 4615-25	16.4	347
482	Perovskite-like Metal Formates with Weak Ferromagnetism and as Precursors to Amorphous Materials. <i>Inorganic Chemistry</i> , 2004 , 43, 4615-25	5.1	305
481	High-dimensional architectures from the self-assembly of lanthanide ions with benzenedicarboxylates and 1,10-phenanthroline. <i>Inorganic Chemistry</i> , 2003 , 42, 4985-94	5.1	294
480	Disorder-order ferroelectric transition in the metal formate framework of [NH ₄][Zn(HCOO) ₃]. <i>Journal of the American Chemical Society</i> , 2010 , 132, 9588-90	16.4	291

479	Solvent-tuned azido-bridged Co ²⁺ layers: square, honeycomb, and Kagomí. <i>Journal of the American Chemical Society</i> , 2006 , 128, 674-5	16.4	273
478	Understanding the Magnetic Anisotropy toward Single-Ion Magnets. <i>Accounts of Chemical Research</i> , 2016 , 49, 2381-2389	24.3	266
477	Cyano-Bridged 4f-3d Coordination Polymers with a Unique Two-Dimensional Topological Architecture and Unusual Magnetic Behavior. <i>Angewandte Chemie - International Edition</i> , 2001 , 40, 434-437	16.4	262
476	Two-Coordinate Co(II) Imido Complexes as Outstanding Single-Molecule Magnets. <i>Journal of the American Chemical Society</i> , 2017 , 139, 373-380	16.4	261
475	Chiral magnetic metal-organic frameworks of dimetal subunits: magnetism tuning by mixed-metal compositions of the solid solutions. <i>Inorganic Chemistry</i> , 2006 , 45, 7069-76	5.1	252
474	Zero-field slow magnetic relaxation from single Co(II) ion: a transition metal single-molecule magnet with high anisotropy barrier. <i>Chemical Science</i> , 2013 , 4, 1802	9.4	251
473	The silica-like extended polymorphism of cobalt(II) imidazolate three-dimensional frameworks: X-ray single-crystal structures and magnetic properties. <i>Chemistry - A European Journal</i> , 2003 , 9, 5673-85	4.8	247
472	The observation of superparamagnetic behavior in molecular nanowires. <i>Journal of the American Chemical Society</i> , 2004 , 126, 8900-1	16.4	241
471	Influence of Guest Exchange on the Magnetization Dynamics of Dilanthanide Single-Molecule-Magnet Nodes within a Metal-Organic Framework. <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 9861-5	16.4	230
470	Capping ligand perturbed slow magnetic relaxation in dysprosium single-ion magnets. <i>Chemistry - A European Journal</i> , 2011 , 17, 12476-81	4.8	225
469	A {Co ₃₂ } nanosphere supported by p-tert-butylthiacalix[4]arene. <i>Journal of the American Chemical Society</i> , 2009 , 131, 11650-1	16.4	219
468	Formate-based magnetic metal-organic frameworks templated by protonated amines. <i>Advanced Materials</i> , 2010 , 22, 1526-33	24	217
467	Two three-dimensional metal-organic frameworks containing one-dimensional hydroxyl/carboxylate mixed bridged metal chains: syntheses, crystal structures, and magnetic properties. <i>Inorganic Chemistry</i> , 2006 , 45, 1508-16	5.1	217
466	Formation of two-dimensional supramolecular icelike layer containing (H ₂ O) ₁₂ rings. <i>Angewandte Chemie - International Edition</i> , 2004 , 43, 1374-6	16.4	217
465	Series of lanthanide organometallic single-ion magnets. <i>Inorganic Chemistry</i> , 2012 , 51, 3079-87	5.1	210
464	Metal-organic perovskites: synthesis, structures, and magnetic properties of [C(NH ₂) ₃][M(II)(HCOO) ₃] (M = Mn, Fe, Co, Ni, Cu, and Zn; C(NH ₂) ₃ = guanidinium). <i>Chemistry - A European Journal</i> , 2009 , 15, 12050-64	4.8	206
463	Stringing oxo-centered trinuclear [MnIII ₃ O] units into single-chain magnets with formate or azide linkers. <i>Angewandte Chemie - International Edition</i> , 2007 , 46, 7388-92	16.4	206
462	An azido-bridged disc-like heptanuclear cobalt(II) cluster: towards a single-molecule magnet. <i>Chemical Communications</i> , 2006 , 3302-4	5.8	196

- 461 High symmetry or low symmetry, that is the question - high performance Dy(III) single-ion magnets by electrostatic potential design. *Chemical Science*, **2016**, 7, 684-691 9.4 194
- 460 Synthesis and characterization of two novel lanthanide coordination polymers with an open framework based on an unprecedented [Ln(7)(micro(3)-OH)(8)](13+) cluster. *Inorganic Chemistry*, **2004**, 43, 1600-2 5.1 188
- 459 A 64-nuclear cubic cage incorporating propeller-like Fe(III)₈ apices and HCOO⁻ edges. *Journal of the American Chemical Society*, **2008**, 130, 10500-1 16.4 179
- 458 Solvothermal reduction synthesis and characterization of superparamagnetic magnetite nanoparticles. *Journal of Materials Chemistry*, **2003**, 13, 1983 176
- 457 Syntheses, crystal structures, and magnetic characterization of five new dimeric manganese(III) tetradentate Schiff base complexes exhibiting single-molecule-magnet behavior. *Inorganic Chemistry*, **2006**, 45, 3538-48 5.1 174
- 456 Magnetic Nanomaterials: Chemical Design, Synthesis, and Potential Applications. *Accounts of Chemical Research*, **2018**, 51, 404-413 24.3 172
- 455 A 2D cyano- and oxamidato-bridged heterotrimetallic Cr(III)-Cu(II)-Gd(III) complex. *Angewandte Chemie - International Edition*, **2003**, 42, 3288-91 16.4 166
- 454 Two 3D porous lanthanide-fumarate-oxalate frameworks exhibiting framework dynamics and luminescent change upon reversible de- and rehydration. *Inorganic Chemistry*, **2007**, 46, 1337-42 5.1 165
- 453 Dehydrogenative coupling of phenanthroline under hydrothermal conditions: crystal structure of a novel layered vanadate complex constructed of 4,8,10-net sheets: [(2,2'-biphen)Co]V₃O₈.5. *Chemical Communications*, **2001**, 1670-1 5.8 159
- 452 Coexistence of Spin-Canting, Metamagnetism, and Spin-Flop in a (4,4) Layered Manganese Azide Polymer. *Chemistry of Materials*, **2005**, 17, 6369-6380 9.6 158
- 451 [NH₂NH₃][M(HCOO)₃] (M = Mn²⁺, Zn²⁺, Co²⁺ and Mg²⁺): structural phase transitions, prominent dielectric anomalies and negative thermal expansion, and magnetic ordering. *Inorganic Chemistry Frontiers*, **2014**, 1, 83-98 6.8 156
- 450 Coordination polymers with end-on azido and pyridine carboxylate N-oxide bridges displaying long-range magnetic ordering with low dimensional character. *Inorganic Chemistry*, **2006**, 45, 6694-705 5.1 156
- 449 Formate--the analogue of azide: structural and magnetic properties of M(HCOO)₂(4,4'-bpy).nH₂O (M = Mn, Co, Ni; n = 0, 5). *Inorganic Chemistry*, **2005**, 44, 572-83 5.1 156
- 448 Monodisperse nickel nanoparticles prepared from a monosurfactant system and their magnetic properties. *Journal of Materials Chemistry*, **2003**, 13, 1510 150
- 447 Single-molecule magnetism of tetrapyrrole lanthanide compounds with sandwich multiple-decker structures. *Coordination Chemistry Reviews*, **2016**, 306, 195-216 23.2 142
- 446 Slow magnetic relaxation in a novel carboxylate/oxalate/hydroxyl bridged dysprosium layer. *Chemical Science*, **2015**, 6, 3095-3101 9.4 138
- 445 Hydroxide-bridged five-coordinate Dy single-molecule magnet exhibiting the record thermal relaxation barrier of magnetization among lanthanide-only dimers. *Chemical Science*, **2017**, 8, 1288-1294^{9.4} 137
- 444 Thiocalix[4]arene-supported planar Ln(4) (Ln = Tb(III), Dy(III)) clusters: toward luminescent and magnetic bifunctional materials. *Inorganic Chemistry*, **2009**, 48, 11743-7 5.1 137

443	Octahedral Fe ₃ O ₄ nanoparticles and their assembled structures. <i>Chemical Communications</i> , 2009 , 4378-808	58.5	133
442	Exchange-coupled nanocomposites: chemical synthesis, characterization and applications. <i>Chemical Society Reviews</i> , 2014 , 43, 8098-113	58.5	132
441	Cobalt-lanthanide coordination polymers constructed with metalloligands: a ferromagnetic coupled quasi-1D Dy ³⁺ chain showing slow relaxation. <i>Chemistry - A European Journal</i> , 2008 , 14, 10340-74.8	7.8	130
440	A pillared layer MOF with anion-tunable magnetic properties and photochemical [2 + 2] cycloaddition. <i>Chemical Communications</i> , 2007 , 1127-9	5.8	129
439	A family of 3d-4f octa-nuclear [Mn(III)(4)Ln(III)(4)] wheels (Ln = Sm, Gd, Tb, Dy, Ho, Er, and Y): synthesis, structure, and magnetism. <i>Inorganic Chemistry</i> , 2010 , 49, 11587-94	5.1	124
438	Phase transitions, prominent dielectric anomalies, and negative thermal expansion in three high thermally stable ammonium magnesium-formate frameworks. <i>Chemistry - A European Journal</i> , 2014 , 20, 1146-58	4.8	123
437	Extended Networks of Co ²⁺ and Mn ²⁺ Bridged by NCS ⁻ /N ₃ ⁻ Anions and Flexible Long Spacers: Syntheses, Structures, and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2005 , 2005, 3277-3286	2.3	122
436	Monodisperse lanthanide oxysulfide nanocrystals. <i>Journal of the American Chemical Society</i> , 2006 , 128, 11758-9	16.4	120
435	Ferrimagnetic-like ordering in a unique three-dimensional coordination polymer featuring mixed azide/carboxylate-bridged trinuclear manganese (II) clusters as subunits. <i>Chemical Communications</i> , 2001 , 2320-1	5.8	120
434	Cyclic water pentamer in a tape-like structure. <i>Chemical Communications</i> , 2004 , 2220-1	5.8	118
433	Unexpected assembly of a unique cyano-bridged three-dimensional Cu ₃ Cr ₂ ferromagnet. <i>Journal of the American Chemical Society</i> , 2001 , 123, 11809-10	16.4	113
432	Copper(II) Complexes of Schiff-Base and Reduced Schiff-Base Ligands: Influence of Weakly Coordinating Sulfonate Groups on the Structure and Oxidation of 3,5-DTBC. <i>European Journal of Inorganic Chemistry</i> , 2005 , 2005, 4635-4645	2.3	110
431	Second ligand-directed self-assembly of lanthanide(III) coordination polymers with 1,4-naphthalenedicarboxylate. <i>New Journal of Chemistry</i> , 2005 , 29, 798	3.6	109
430	Azide-bridged one-dimensional Mn(III) polymers: effects of side group of Schiff base ligands on structure and magnetism. <i>Inorganic Chemistry</i> , 2007 , 46, 11235-42	5.1	108
429	Synthesis, crystal structures, and magnetism of cobalt coordination polymers based on dicyanamide and pyrazine-dioxide derivatives. <i>Inorganic Chemistry</i> , 2005 , 44, 2169-76	5.1	108
428	New Porous Lanthanide-Organic Frameworks: Synthesis, Characterization, and Properties of Lanthanide 2,6-Naphthalenedicarboxylates. <i>European Journal of Inorganic Chemistry</i> , 2004 , 2004, 3262-3268	2.3	106
427	Two molecular tapes consisting of serial or parallel azido-bridged eight-membered copper rings. <i>Angewandte Chemie - International Edition</i> , 2005 , 44, 5841-6	16.4	106
426	Ferromagnetic Ordering in a Two-Dimensional Copper Complex with Dual End-to-End and End-On Azide Bridges This work was supported by the National Natural Science Foundation of China (NSF 29631040 and 29929001), The Major State Basic Research Development Program (Grant No. G2000077500 and No. G1998061206), and the Malaysian Government research grant R&D No. 305/pfizik/622004. <i>Angewandte Chemie - International Edition</i> 2000 , 39, 3633-3635	16.4	105

425	Highly near-IR emissive ytterbium(iii) complexes with unprecedented quantum yields. <i>Chemical Science</i> , 2017 , 8, 2702-2709	9.4	102
424	Spin crossover in a series of iron(II) complexes of 2-(2-Alkyl-2H-tetrazol-5-yl)-1,10-phenanthroline: effects of alkyl side chain, solvent, and anion. <i>Inorganic Chemistry</i> , 2007 , 46, 2541-55	5.1	102
423	Diversity of azido magnetic chains constructed with flexible ligand 2,2'-dipyridylamine. <i>Inorganic Chemistry</i> , 2009 , 48, 1301-8	5.1	99
422	Two chain compounds of $[M(N_3)_2(HCOO)][(CH_3)_2NH_2]$ (M = Fe and Co) with a mixed azido/formate bridge displaying metamagnetic behavior. <i>Inorganic Chemistry</i> , 2006 , 45, 2782-4	5.1	99
421	Four 2D metal-organic networks incorporating Cd-cluster SUBs: hydrothermal synthesis, structures and photoluminescent properties. <i>CrystEngComm</i> , 2009 , 11, 122-129	3.3	95
420	A Mononuclear Dysprosium Complex Featuring Single-Molecule-Magnet Behavior. <i>Angewandte Chemie</i> , 2010 , 122, 7610-7613	3.6	95
419	An interesting magnetic behavior in molecular solid containing one-dimensional Ni(III) chain. <i>Chemical Physics Letters</i> , 2004 , 396, 353-358	2.5	95
418	Synthesis and size-dependent magnetic properties of monodisperse EuS nanocrystals. <i>Small</i> , 2006 , 2, 244-8	11	94
417	Observation of the single-ion magnet behavior of d ions on two-coordinate Co(i)-NHC complexes. <i>Chemical Science</i> , 2015 , 6, 7156-7162	9.4	93
416	Pure trinuclear 4f single-molecule magnets: synthesis, structures, magnetism and ab initio investigation. <i>Chemistry - A European Journal</i> , 2011 , 17, 2458-66	4.8	91
415	An A-site mixed-ammonium solid solution perovskite series of $[(NH_2 NH_3)_x (CH_3 NH_3)_{1-x}][Mn(HCOO)_3]$ (x=1.00-0.67). <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 11093-6	16.4	89
414	$[M(N_3)_2(H_2O)_2] \times (bpeado)$: unusual antiferromagnetic Heisenberg chain (M = Mn) and ferromagnetic Ising chain (M = Co) with large coercivity and magnetic relaxation (bpeado = 1,2-bis(4-pyridyl)ethane-N,N'-dioxide). <i>Chemistry - A European Journal</i> , 2009 , 15, 1757-64	4.8	89
413	Rational synthesis and magnetic properties of a family of low-dimensional heterometallic Cr-Mn complexes based on the versatile building block $[Cr(2,2'-bipyridine)(CN)_4]^-$. <i>Inorganic Chemistry</i> , 2005 , 44, 4534-45	5.1	88
412	Synthesis, crystal structures, and magnetic properties of cyano-bridged heterobimetallic chains based on $[(Tp)Fe(CN)_3]^-$. <i>Inorganic Chemistry</i> , 2006 , 45, 8942-9	5.1	87
411	Coexistence of long-range ferromagnetic ordering and glassy behavior in one-dimensional bimetallic cyano-bridged polymers. <i>Inorganic Chemistry</i> , 2003 , 42, 6123-9	5.1	87
410	A unique 3D alternating ferro- and antiferromagnetic manganese azide system with threefold interpenetrating (10,3) nets. <i>Angewandte Chemie - International Edition</i> , 2004 , 43, 990-4	16.4	86
409	A three-dimensional manganeseII complex exhibiting ferrimagnetic and metamagnetic behaviors. <i>Inorganic Chemistry</i> , 2003 , 42, 5486-8	5.1	86
408	A 3D porous lanthanide-fumarate framework with water hexamer occupied cavities, exhibiting a reversible dehydration and rehydration procedure. <i>Dalton Transactions</i> , 2006 , 765-8	4.3	85

407	[CH ₃ NH ₂ (CH ₂) ₂ NH ₂ CH ₃][M ₂ (HCOO) ₆] (M = MnII and CoII): weak ferromagnetic metal formate frameworks of unique binodal 6-connected (4(12).6(3))(4(9).6(6)) topology, templated by a diammonium cation. <i>Inorganic Chemistry</i> , 2007 , 46, 8439-41	5.1	84
406	Cu ₄ (CH ₃ C(OH)(PO ₃) ₂) ₂ (C ₄ H ₄ N ₂)(H ₂ O) ₄ : a novel, three-dimensional copper diphosphonate with metamagnetism. <i>Chemical Communications</i> , 2001 , 2346-7	5.8	84
405	Single-molecule-magnet behavior in a Fe(12)Sm(4) cluster. <i>Inorganic Chemistry</i> , 2010 , 49, 9734-6	5.1	83
404	Thermostability and photoluminescence of Dy(III) single-molecule magnets under a magnetic field. <i>Chemical Science</i> , 2016 , 7, 5020-5031	9.4	83
403	Rational enhancement of the energy barrier of bis(tetrapyrrole) dysprosium SMMs replacing atom of porphyrin core. <i>Chemical Science</i> , 2015 , 6, 5947-5954	9.4	82
402	Sandwich-type tetrakis(phthalocyaninato) dysprosium-cadmium quadruple-decker SMM. <i>Chemical Communications</i> , 2011 , 47, 9624-6	5.8	82
401	Observation of an octameric water cluster containing a book-shaped hexamer in a 4f-3d complex. <i>Chemical Communications</i> , 2005 , 2336-8	5.8	82
400	Detailed magnetic studies on Co(N ₃) ₂ (4-acetylpyridine) ₂ : a weak-ferromagnet with a very big canting angle. <i>Inorganic Chemistry</i> , 2008 , 47, 5720-6	5.1	81
399	Multiple Regulated Assembly, Crystal Structures and Magnetic Properties of Porous Coordination Polymers with Flexible Ligands. <i>European Journal of Inorganic Chemistry</i> , 2005 , 2005, 4150-4159	2.3	81
398	Ferromagnetic Ordering in a Diamond-Like Cyano-Bridged Mn(II) Ru(III) Bimetallic Coordination Polymer. <i>Angewandte Chemie - International Edition</i> , 2001 , 40, 3031-3	16.4	81
397	Synthesis, crystal structures and magnetic properties of 1D polymeric [MnIII(salen)N ₃] and [MnIII(salen)Ag(CN) ₂] complexes. <i>New Journal of Chemistry</i> , 2002 , 26, 1025-1028	3.6	81
396	Modulating the phases of iron carbide nanoparticles: from a perspective of interfering with the carbon penetration of Fe@FeO by selectively adsorbed halide ions. <i>Chemical Science</i> , 2017 , 8, 473-481	9.4	80
395	Magnetic molecular materials with paramagnetic lanthanide ions. <i>Science in China Series B: Chemistry</i> , 2009 , 52, 1739-1758		80
394	Linking cyano-bridged ladders by azide to form a layered metamagnet. <i>Chemical Communications</i> , 2004 , 1906-7	5.8	80
393	A 36-fold multiple unit cell and switchable anisotropic dielectric responses in an ammonium magnesium formate framework. <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 2534-7	16.4	79
392	Chemical synthesis and magnetic study of nanocrystalline thin films of cobalt spinel ferrites. <i>Solid State Communications</i> , 1998 , 107, 471-476	1.6	79
391	Synthesis, Structures and Catecholase Activity of a New Series of Dicopper(II) Complexes of Reduced Schiff Base Ligands. <i>European Journal of Inorganic Chemistry</i> , 2006 , 2006, 2656-2670	2.3	79
390	Synthesis, structure, and magnetism of a ferric 24-azametallacrown-8 complex. <i>Inorganic Chemistry</i> , 2004 , 43, 2222-4	5.1	78

- 389 A Family of Co(II)Co(III)₃ Single-Ion Magnets with Zero-Field Slow Magnetic Relaxation: Fine Tuning of Energy Barrier by Remote Substituent and Counter Cation. *Inorganic Chemistry*, **2015**, 54, 5475-86 5.1 77
- 388 Shape-dependent magnetic properties of low-dimensional nanoscale Prussian blue (PB) analogue SmFe(CN)₆·4H₂O. *Chemical Communications*, **2005**, 4339-41 5.8 77
- 387 A two-step field-induced magnetic transition in a novel layered cobalt diphosphonate. *Chemical Communications*, **2003**, 1076-7 5.8 76
- 386 Characterization of a soluble molecular magnet: unusual magnetic behavior of cyano-bridged Gd(III)-Cr(III) complexes with one-dimensional and nanoscaled square structures. *Inorganic Chemistry*, **2002**, 41, 4756-62 5.1 76
- 385 One-dimensional cobalt diphosphonates exhibiting weak ferromagnetism and field-induced magnetic transitions. *Inorganic Chemistry*, **2004**, 43, 2151-6 5.1 75
- 384 Ferromagnetic ordering and metamagnetism in malonate bridged 3D diamond-like and honeycomb-like networks: [Cu(mal)(DMF)]_n and [[Cu(mal)(0.5pyz)]·H₂O]_n (mal = Malonate Dianion, DMF = N,N-dimethylformamide, pyz = Pyrazine). *Inorganic Chemistry*, **2003**, 42, 4792-4 5.1 75
- 383 Controlled synthesis of heterotrimetallic single-chain magnets from anisotropic high-spin 3 d-4 f nodes and paramagnetic spacers. *Chemistry - A European Journal*, **2013**, 19, 294-303 4.8 74
- 382 A family of porous magnets, [M₃(HCOO)₆] (M=Mn, Fe, Co and Ni). *Polyhedron*, **2007**, 26, 2207-2215 2.7 73
- 381 Low-Coordinate Single-Ion Magnets by Intercalation of Lanthanides into a Phenol Matrix. *Angewandte Chemie - International Edition*, **2018**, 57, 4673-4676 16.4 71
- 380 Novel antitumor agent, trilacunary Keggin-type tungstobismuthate, inhibits proliferation and induces apoptosis in human gastric cancer SGC-7901 cells. *Inorganic Chemistry*, **2013**, 52, 5119-27 5.1 71
- 379 Heterometallic MIIIRuIII₂ compounds constructed from trans-[Ru(Salen)(CN)₂]- and trans-[Ru(Acac)₂(CN)₂]-. Synthesis, structures, magnetic properties, and density functional theoretical study. *Inorganic Chemistry*, **2005**, 44, 6579-90 5.1 71
- 378 A Novel Azido and Pyrazine-Dioxide Bridged Three-Dimensional Manganese(II) Network with Antiferromagnetic Ordering (TN = 62 K) and a Spin Flop State. *Chemistry of Materials*, **2001**, 13, 1946-1948 8.6 71
- 377 Endohedral Metallofullerene as Molecular High Spin Qubit: Diverse Rabi Cycles in Gd@CN. *Journal of the American Chemical Society*, **2018**, 140, 1123-1130 16.4 70
- 376 A new series of chiral metal formate frameworks of [HONH₃][M(II)(HCOO)₃] (M = Mn, Co, Ni, Zn, and Mg): synthesis, structures, and properties. *Inorganic Chemistry*, **2012**, 51, 13363-72 5.1 70
- 375 Making a [Co₂₄] metallamacrocycle from the shuttlecock-like tetranuclear cobalt-calixarene building blocks. *Chemical Communications*, **2010**, 46, 6362-4 5.8 70
- 374 An enantiopure Fe(III)₄ single-molecule magnet. *Chemical Communications*, **2011**, 47, 8049-51 5.8 70
- 373 One-dimensional ferromagnetically coupled bimetallic chains constructed with trans-[Ru(acac)₂(CN)₂]-: syntheses, structures, magnetic properties, and density functional theoretical study. *Chemistry - A European Journal*, **2010**, 16, 3524-35 4.8 70
- 372 Hydrothermal synthesis and crystal structure of a novel two-dimensional vanadium oxide complex with a 6,14-net sinusoidal ruffling anionic layer: [Ni(phen)₂V(4)O(11)] (phen = 1,10-phenanthroline). *Inorganic Chemistry*, **2002**, 41, 140-3 5.1 70

371	Structures and magnetism of two novel heptanuclear lanthanide-centered trigonal prismatic clusters: $[\text{LnCu}_6(\mu_3\text{-OH})_3(\text{HL})_2(\text{L})_4](\text{ClO}_4)_2 \cdot 2.25\text{H}_2\text{O}$ (Ln = La, Tb; H ₂ L = iminodiacetic acid). <i>Inorganic Chemistry</i> , 2000 , 39, 2488-92	5.1	70
370	Magnetic properties of EuS nanoparticles synthesized by thermal decomposition of molecular precursors. <i>Journal of Materials Chemistry</i> , 2005 , 15, 4209		69
369	Metamagnetism of the First Cyano-Bridged Two-Dimensional Brick-Wall-like 4fBd Array. <i>Chemistry of Materials</i> , 2001 , 13, 1431-1433	9.6	69
368	Three-Dimensional Eight- or Four-Connected MetalOrganic Frameworks Tuned by Hydrothermal Temperatures. <i>Crystal Growth and Design</i> , 2007 , 7, 1312-1317	3.5	68
367	Synthesis, crystal structure, and magnetic properties of two cyano-bridged bimetallic 4f-3d arrays with one-dimensional chain and two-dimensional brick wall molecular structures. <i>Inorganic Chemistry</i> , 2001 , 40, 6295-300	5.1	68
366	Hydrothermal Syntheses, Architectures and Magnetic Properties of Six Novel MnII Coordination Polymers with Mixed Ligands. <i>European Journal of Inorganic Chemistry</i> , 2006 , 2006, 2411-2421	2.3	67
365	Long-range ferromagnetic ordering in two-dimensional coordination polymers $\text{Co}[\text{N}(\text{CN})_2]_2(\text{L})$ [L = pyrazine dioxide (pzdo) and 2-methyl pyrazine dioxide (mpdo)] with dual μ - and μ_3 -[N(CN) ₂] bridges. <i>Inorganic Chemistry</i> , 2003 , 42, 5399-404	5.1	67
364	Soft chemical synthesis and transport properties of $\text{La}_{0.7}\text{Sr}_{0.3}\text{MnO}_3$ granular perovskites. <i>Solid State Communications</i> , 2000 , 114, 43-47	1.6	67
363	Synthesis, structures, and magnetic properties of a series of cyano-bridged Fe-Mn bimetallic complexes. <i>Inorganic Chemistry</i> , 2006 , 45, 5018-26	5.1	66
362	Synthesis, structure, and magnetism of three azido-bridged Co^{2+} compounds with a flexible coligand 1,2-(tetrazole-1-yl)ethane. <i>Inorganic Chemistry</i> , 2008 , 47, 8134-42	5.1	65
361	A cyano-bridged molecular magnet with a novel two-dimensional brick wall structure. <i>Chemical Communications</i> , 2000 , 1309-1310	5.8	65
360	A Prussian-blue type ferrimagnet $\text{Na}[\text{MnCr}(\text{CN})_6]$: single crystal structure and magnetic properties. <i>Inorganic Chemistry</i> , 2004 , 43, 2465-7	5.1	64
359	Elucidation of slow magnetic relaxation in a ferromagnetic 1D dysprosium chain through magnetic dilution. <i>Chemical Communications</i> , 2014 , 50, 6052-5	5.8	63
358	Magnetic Properties of Metal Diphosphonate Compounds with One-Dimensional Chain Structures. <i>Chemistry of Materials</i> , 2003 , 15, 3233-3236	9.6	63
357	Syntheses, structures and magnetic properties of three-dimensional co-ordination polymers constructed by dimer subunits. <i>Dalton Transactions RSC</i> , 2000 , 4187-4191		63
356	Cobalt(II) coordination polymer exhibiting single-ion-magnet-type field-induced slow relaxation behavior. <i>Inorganic Chemistry</i> , 2015 , 54, 3716-8	5.1	62
355	Six-coordinate lanthanide complexes: slow relaxation of magnetization in the dysprosium(III) complex. <i>Chemistry - A European Journal</i> , 2014 , 20, 15975-80	4.8	62
354	Constructing single-chain magnets by supramolecular π -stacking and spin canting: a case study on manganese(III) corroles. <i>Chemistry - A European Journal</i> , 2012 , 18, 915-24	4.8	62

- 353 Angular-resolved magnetometry beyond triclinic crystals: out-of-equilibrium studies of Cp*ErCOT single-molecule magnet. *Chemistry - A European Journal*, **2013**, 19, 13726-31 4.8 62
- 352 Tuning the magnetic behavior via dehydration/hydration treatment of a new ferrimagnet with the composition of K_{0.2}Mn_{1.4}Cr(CN)₆ × 6H₂O. *Inorganic Chemistry*, **2006**, 45, 999-1004 5.1 61
- 351 The First One-Dimensional Molecular Material Featuring Maleonitriledithiolate as a Subunit Exhibiting Metamagnetic-Like Behavior. *European Journal of Inorganic Chemistry*, **2003**, 2003, 2393-2396 2.3 61
- 350 Pendant macrocyclic metallic building blocks for the design of cyano-bridged heterometallic complexes with 1D chain and 2D layer structures. *Inorganic Chemistry*, **2003**, 42, 5604-11 5.1 61
- 349 A 1D dysprosium chain with slow magnetic relaxation constructed from a pyridine-N-oxide ligand. *Chemical Communications*, **2014**, 50, 10434-7 5.8 60
- 348 Metamagnetic Copper(II) Diphosphonates with Layered Structures. *Chemistry of Materials*, **2002**, 14, 3143-3147 9.6 60
- 347 (Boratabenzene)(cyclooctatetraenyl) lanthanide complexes: a new type of organometallic single-ion magnet. *Inorganic Chemistry Frontiers*, **2016**, 3, 828-835 6.8 59
- 346 Variable dimensionality from mononuclear and trinuclear to one and two dimensions: a series of copper(II) compounds with 4,4'-dipyridine dioxide. *Inorganic Chemistry*, **2001**, 40, 6247-53 5.1 59
- 345 Synthesis, Structure, and Magnetic Property of a New Open-Framework Manganese Borophosphate, [NH₄]₄[Mn₉B₂(OH)₂(HPO₄)₄(PO₄)₆]. *Chemistry of Materials*, **2006**, 18, 476-481 9.6 58
- 344 Crystal structure and magnetic properties of a new two-dimensional cyano-bridged bimetallic assembly [Ni(L[?])₃[Cr(CN)₅(NO)]₂ · 10H₂O (L[?] = 3,10-dimethyl-1,3,5,8,10,12-hexaazacyclotetradecane). *Chemical Communications*, **2000**, 713-714 5.8 58
- 343 Does the thermal evolution of molecular structures critically affect the magnetic anisotropy?. *Chemical Science*, **2015**, 6, 4587-4593 9.4 57
- 342 Transition Metal (Fe, Co and Ni) Carbide and Nitride Nanomaterials: Structure, Chemical Synthesis and Applications. *ChemNanoMat*, **2015**, 1, 376-398 3.5 57
- 341 Azide and oxo bridged ferromagnetic clusters: Three face-shared tetracubane Ni(II)/Co(II) hexamers and a wheel-shaped SMM-like Co(II) heptamer. *Inorganica Chimica Acta*, **2008**, 361, 3895-3902 2.7 57
- 340 Hydrothermal Syntheses, Structures, and Properties of Three 3-D Lanthanide Coordination Polymers that Form 1-D Channels. *European Journal of Inorganic Chemistry*, **2004**, 2004, 2968-2973 2.3 57
- 339 New route to the mixed valence semiquinone-catecholate based mononuclear Fe(III) and catecholate based dinuclear Mn(III) complexes: first experimental evidence of valence tautomerism in an iron complex. *Inorganic Chemistry*, **2004**, 43, 5908-18 5.1 57
- 338 Weak Ligand-Field Effect from Ancillary Ligands on Enhancing Single-Ion Magnet Performance. *Chemistry - A European Journal*, **2016**, 22, 12724-31 4.8 56
- 337 Reversible de-/resolution and accompanied magnetism modulation in a framework of topologically ferrimagnetic [Co₃(μ₃-OH)₂]_n chains linked by a V-shaped ligand 4,4'-dicarboxybiphenyl sulfone. *Chemical Communications*, **2010**, 46, 4339-41 5.8 56
- 336 Field-induced magnetic transitions in metal phosphonates with ladderlike chain structures: (NH₃C₆H₄NH₃)₂M₂(hedpH)₂ · H₂O [M = Fe, Co, Mn, Zn; hedp = C(CH₃)(OH)(PO₃)₂]. *Inorganic Chemistry*, **2005**, 44, 2761-5 5.1 56

- 335 Abrupt spin transition around room temperature and light induced properties in Fe(II) complexes with N4O2 coordination sphere. *Chemical Communications*, **2010**, 46, 2554-6 5.8 55
- 334 Synthesis and characterization of copper(II) azide complexes containing polyamines as co-ligands. *Polyhedron*, **2004**, 23, 1243-1248 2.7 55
- 333 A two-fold interpenetrated three-dimensional cobalt(II) complex with dual dicyanamide and 4,4'-bipyridine bridges. *New Journal of Chemistry*, **2000**, 24, 953-954 3.6 55
- 332 Synthesis, crystal structure and magnetic property of a new 1D molecular material [1-(4'-chlorobenzyl)2-aminopyridinium](+) bis(maleonitriledithiolato)nickel(II) *Journal of Solid State Chemistry*, **2005**, 178, 100-105 3.3 54
- 331 Oxo-centered regular octahedral lanthanide clusters. *New Journal of Chemistry*, **2000**, 24, 61-62 3.6 54
- 330 Three p-tert-butylthiacalix[4]arene-supported cobalt compounds obtained in one pot involving in situ formation of N6H2 ligand. *Inorganic Chemistry*, **2010**, 49, 7735-40 5.1 53
- 329 Structures and magnetism of cyano-bridged grid-like two-dimensional 4f3d arrays. *Journal of Materials Chemistry*, **2006**, 16, 2625-2634 5.3
- 328 Single domain SmCo5@Co exchange-coupled magnets prepared from core/shell Sm[Co(CN)6]4H2O@GO particles: a novel chemical approach. *Scientific Reports*, **2013**, 3, 3542 4.9 52
- 327 Assembly of hydrogen bonded diamondoid networks based on synthetic metal-organic tetrahedral nodes. *Inorganic Chemistry*, **2005**, 44, 837-9 5.1 52
- 326 [Cu(tn)]3[W(CN)8]2·2H2O and [Cu(pn)]3[W(CN)8]2·2H2O: Two Novel Cu(II)/W(V) Cyano-Bridged Two-Dimensional Coordination Polymers with Metamagnetism. *Chemistry of Materials*, **2003**, 15, 2094-2098 8.6 52
- 325 Solvothermal reduction synthesis and magnetic properties of polymer protected iron and nickel nanocrystals. *Journal of Alloys and Compounds*, **2004**, 365, 112-116 5.7 52
- 324 Magneto-structural correlation in a series of bimetallic alternating chain complexes of [CrIII(LCN)4]n[MnIII(salpn)n]n·solvents (L=2,2'-bipy or 9,10-phen, salpn=substituted salicylaldehyde, solvents=water and methanol). *Inorganic Chemistry*, **2007**, 46, 10221-8 5.1 51
- 323 Syntheses, structures, and magnetic properties of mixed-valent diruthenium(II,III) diphosphonates with discrete and one-dimensional structures. *Inorganic Chemistry*, **2005**, 44, 4309-14 5.1 51
- 322 Metal-organic niccolite: synthesis, structures, phase transition, and magnetic properties of [CH3NH2(CH2)2NH2CH3][M2(HCOO)6] (M=divalent Mn, Fe, Co, Ni, Cu and Zn). *Chemistry - an Asian Journal*, **2011**, 6, 3084-96 4.5 50
- 321 New heterometallic [Mn(III)(4)Ln(III)(4)] wheels incorporating formate ligands. *Dalton Transactions*, **2010**, 39, 3375-7 4.3 50
- 320 Hydrogencyanamide-bridged one-dimensional polymers built on Mn(III)-Schiff base fragments: synthesis, structure, and magnetism. *Chemistry - A European Journal*, **2007**, 13, 2937-52 4.8 50
- 319 One-dimensional coordination polymers [Co(acac)2pz]n and [Co(acac)2(4,4'-bipy)]n (acac=acetylacetonate, pz=pyrazine, bipy=4,4'-bipyridine): synthesis, structures and magnetic properties. *Polyhedron*, **2001**, 20, 1255-1261 2.7 50
- 318 Observation of an unusual field-dependent slow magnetic relaxation and two distinct transitions in a family of rare-earth/transition-metal complexes. *Physical Review B*, **2001**, 63, 3.3 50

- 317 A novel bimetallic cage complex constructed from six V4Co pentatomic rings: hydrothermal synthesis and crystal structure of [(2,2LPy2NH)2Co]3V8O23. *Chemical Communications*, **2001**, 1636-7 5.8 50
- 316 2D LnIII RuIII 2 compounds constructed from trans-[Ru(acac)2(CN)2]-. Syntheses, structures, and magnetic properties. *Inorganic Chemistry*, **2006**, 45, 6756-60 5.1 49
- 315 Novel three-dimensional networks cross-linked by two sets of hydrogen bonded frameworks based on 1-D {[M(H2O)4(4,4?-dpdo)][ClO4]2[2(4,4?-dpdo)]n} (4,4?-dpdo = 4,4?-bipyridine dioxide; M = Co, Ni, Cu or Zn). *Dalton Transactions RSC*, **2001**, 130-133 49
- 314 Erbium-formate frameworks templated by diammonium cations: syntheses, structures, structural transition and magnetic properties. *Dalton Transactions*, **2011**, 40, 6038-46 4.3 48
- 313 Two Mn(II)(2)Ln(III)(4) (Ln = Gd, Eu) hexanuclear compounds of p-tert-butylsulfanylcalix[4]arene. *Dalton Transactions*, **2009**, 2250-4 4.3 48
- 312 Edta-linked 4f-3d heterometallic two dimensional sheet in Ln2M3(edta)3(H2O)11·2H2O (Ln?Nd, Gd; M?Mn, Co). *Polyhedron*, **1998**, 17, 2243-2248 2.7 48
- 311 Eight-connected three-dimensional lanthanide coordination networks constructed by pyrazine-dioxide (pzdo). *Microporous and Mesoporous Materials*, **2004**, 73, 89-95 5.3 48
- 310 Lanthanide contraction and anions-controlled dimensional diversity in Ln?Cu?NTA (Ln=lanthanide, H3NTA=nitrilotriacetic acid) coordination polymers: crystal structure and magnetic properties. *Polyhedron*, **2002**, 21, 1097-1103 2.7 48
- 309 Antiferromagnetic ordering in a novel five-connected 3D polymer {Cu2(2,5-Me2pyz)[N(CN)2]4}n(2,5-Me2pyz, 2,5-dimethylpyrazine). *New Journal of Chemistry*, **2002**, 26, 523-525 3.6 48
- 308 Anion controlled 2D assembly of a La?Cu cation array and its unusual magnetic properties. *Chemical Communications*, **2000**, 1685-1686 5.8 48
- 307 A soft phosphorus atom to "harden" an erbium(iii) single-ion magnet. *Chemical Science*, **2018**, 9, 7540-7545 4.7
- 306 The slow magnetic relaxation regulated by ligand conformation of a lanthanide single-ion magnet [Hex4N][Dy(DBM)4]. *Inorganic Chemistry Frontiers*, **2014**, 1, 503-509 6.8 47
- 305 Thermal cleavage of cyclobutane rings in photodimerized coordination-polymeric sheets. *Chemistry - A European Journal*, **2012**, 18, 7869-77 4.8 47
- 304 A Novel Three-Dimensional Network Containing Mn(II) Ions and Tricyanomethanide with Rare 4f-64 Topology. *Crystal Growth and Design*, **2005**, 5, 1331-1333 3.5 47
- 303 Synthesis, crystal structure and metamagnetic properties of a two-dimensional honeycomb network based on ferricyanide and (3,10-dimethyl-1,3,5,8,10,12-hexaazacyclotetradecane)nickel(II) building blocks. *Journal of the Chemical Society Dalton Transactions*, **1999**, 2477-2480 47
- 302 The inhibitory effects of a new cobalt-based polyoxometalate on the growth of human cancer cells. *Dalton Transactions*, **2014**, 43, 6070-8 4.3 46
- 301 Syntheses, structures and properties of seven isomorphous 1D Ln3+ complexes Ln(BTA)(HCOO)(H2O)3 (H2BTA = bis(tetrazoly)amine, Ln = Pr, Gd, Eu, Tb, Dy, Er, Yb) and two 3D Ln3+ complexes Ln(HCOO)3 (Ln = Pr, Nd). *Dalton Transactions*, **2008**, 6165-9 4.3 46
- 300 Structures and Magnetism of a Series Mn(II) Coordination Polymers Containing Pyrazine-Dioxide Derivatives and Different Anions. *Crystal Growth and Design*, **2005**, 5, 269-277 3.5 45

299	Syntheses, crystal structures and magnetic properties of manganese(II)-hedp compounds involving alkylenediamine templates (hedp = 1-hydroxyethylidene-diphosphonate). <i>Dalton Transactions RSC</i> , 2002 , 2752-2759		45
298	A novel Cu(II)-W(V) bimetallic assembly magnet $[\{Cu(en)_2\}_3\{W(CN)_8\}_2\{H_2O\}_4]_n$ (en = ethylenediamine) with cube-like W_8Cu_{12} units from a coordinated anion template self-assembly reaction. <i>New Journal of Chemistry</i> , 2002 , 26, 485-489	3.6	45
297	Coupling Influences SMM Properties for Pure 4 f Systems. <i>Chemistry - A European Journal</i> , 2018 , 24, 6079-6086	4.6	44
296	A half-sandwich organometallic single-ion magnet with hexamethylbenzene coordinated to the Dy(III) ion. <i>Chemical Communications</i> , 2014 , 50, 11418-20	5.8	44
295	Controlled synthesis and multifunctional properties of FePt-Au heterostructures. <i>Nano Research</i> , 2011 , 4, 836-848	10	44
294	Magnetization relaxation in a three-dimensional ligated cobalt phosphonate containing ferrimagnetic chains. <i>Chemistry - A European Journal</i> , 2011 , 17, 3579-83	4.8	44
293	A cyano-bridged MnII/MoV bimetallic ferrimagnet with a novel moniliform structure. <i>Dalton Transactions RSC</i> , 2002 , 2805		44
292	Reactions between macrocyclic nickel(II) complexes and ferricyanide. Metamagnetic properties of a two-dimensional honeycomb assembly $[NiL_3]_3[Fe(CN)_6]_2\{8H_2O\}$ (L3 = 3,10-dihydroxyethyl-1,3,6,8,10,12-hexaazacyclotetradecane). <i>Dalton Transactions RSC</i> , 2000 , 2996-3000		44
291	Effects of Organic Templates on Directing the Structures of Nickel(II)-Hydroxyethylidenediphosphonate Compounds: A Structural and Magnetic Study. <i>Chemistry of Materials</i> , 1999 , 11, 2382-2388	9.6	44
290	Bimetallic sheet and 3D threefold interpenetrating diamond-like network constructed by chelate Cu cations and Mn dicyanamide polymeric chains. Synthesis, crystal structure, and magnetism of $[Cu(L)_2][Mn(dca)_4]$ (L=ethylenediamine or 1,3-diaminopropane; dca=dicyanamide $N(CN)_2$)	2.7	43
289	Temperature-Induced Irreversible Phase Transition From Perovskite to Diamond But Pressure-Driven Back-Transition in an Ammonium Copper Formate. <i>Angewandte Chemie - International Edition</i> , 2016 , 55, 2097-100	16.4	43
288	Honeycomb layer of cobalt(II) azide hydrazine showing weak ferromagnetism. <i>Inorganic Chemistry</i> , 2007 , 46, 10452-4	5.1	42
287	3D Self-penetrating coordination network constructed by dicyanamide and 1,2-bis(4-pyridyl)ethane-N,N'-dioxide (bpeado). <i>CrystEngComm</i> , 2004 , 6, 579-583	3.3	42
286	Syntheses, structures and magnetic properties of 1-D coordination polymers containing both dicyanamide and 2-pyrrolidone. <i>Inorganic Chemistry Communication</i> , 2001 , 4, 72-75	3.1	42
285	Dinuclear dysprosium SMMs bridged by a neutral bipyrimidine ligand: two crystal systems that depend on different lattice solvents lead to a distinct slow relaxation behaviour. <i>Dalton Transactions</i> , 2016 , 45, 8790-4	4.3	42
284	Synthesis, magnetic and photomagnetic study of new iron(II) spin-crossover complexes with N_4O_4 coordination sphere. <i>Dalton Transactions</i> , 2010 , 39, 4856-68	4.3	41
283	Heterometallic Cr-Mn complexes containing cyanide and oxalate bridges. <i>Inorganic Chemistry</i> , 2006 , 45, 5447-54	5.1	41
282	$[Zn_3(HCOO)_6]$: A Porous Diamond Framework Conformable to Guest Inclusion. <i>Australian Journal of Chemistry</i> , 2006 , 59, 617	1.2	41

281	Template- and pH-directed assembly of diruthenium diphosphonates with different topologies and oxidation states. <i>Inorganic Chemistry</i> , 2006 , 45, 4205-13	5.1	41
280	Copper phosphonates with dinuclear and layer structures: a structural and magnetic study. <i>Journal of Solid State Chemistry</i> , 2004 , 177, 4557-4563	3.3	41
279	Can Non-Kramers Tm(III) Mononuclear Molecules be Single-Molecule Magnets (SMMs)?. <i>Chemistry - A European Journal</i> , 2016 , 22, 4704-8	4.8	40
278	A New Bis(phthalocyaninato) Terbium Single-Ion Magnet with an Overall Excellent Magnetic Performance. <i>Inorganic Chemistry</i> , 2017 , 56, 13889-13896	5.1	40
277	Hydrothermal synthesis, structures and magnetic studies of transition metal sulfates containing hydrazine. <i>Inorganic Chemistry</i> , 2011 , 50, 144-54	5.1	40
276	Pyrolysis of single molecular precursor for monodisperse lanthanide sulfide/oxysulfide nanocrystals. <i>Journal of Materials Chemistry</i> , 2008 , 18, 949-953		40
275	Structure and magnetic properties of a novel copper diphosphonate with pillared layered structure:: Cu ₂ (H ₂ O) ₂ {O ₃ PCH ₂ N(C ₂ H ₄) ₂ NCH ₂ PO ₃ }. <i>Journal of Solid State Chemistry</i> , 2004 , 177, 1297-1307	3.3	40
274	Magnetic characteristics of new cyano-bridged two-dimensional honeycomb-like bimetallic assemblies containing Ni(II)-N[triple bond]C-Cr(III) or Ni(II)-N[triple bond]C-Cr(I) linkages. <i>Inorganic Chemistry</i> , 2001 , 40, 6287-94	5.1	40
273	Hydrothermal syntheses and crystal structures of two-dimensional (2D) layered vanadium oxide complexes: M(bipy)(H ₂ O)V ₂ O ₆ (M = Ni, Co, bipy = bipyridine) and [Ni(bipy) ₂ V ₆ O ₁₇]. <i>Dalton Transactions RSC</i> , 2002 , 598		39
272	Pyrazine dioxide bridged two-dimensional antiferromagnets [M(NCS) ₂ (pzdo) ₂] (M = Mn, Co; pzdo = pyrazine dioxide). <i>Chemical Communications</i> , 2001 , 2586-2587	5.8	39
271	Co-crystallized fullerene and a mixed (phthalocyaninato)(porphyrinato) dysprosium double-decker SMM. <i>Chemical Science</i> , 2014 , 5, 3214-3220	9.4	38
270	Synthesis, crystal structure and magnetic behavior of two cobalt coordination polymers with 1,2-bis(1,2,4-triazol-1-yl)ethane and dicyanamide. <i>Polyhedron</i> , 2007 , 26, 5219-5224	2.7	38
269	Spontaneous magnetization below 7.7 K based on an extended 3-D H-bonding network material: synthesis, crystal structure and magnetic properties. <i>Dalton Transactions RSC</i> , 2002 , 3915-3918		38
268	Crystal structures and magnetic properties of new cyano-bridged two-dimensional grid-like bimetallic assemblies [Ni(tn) ₂] ₂ [Cr(CN) ₅ ((NO))OH*H ₂ O and [Ni(tn) ₂] ₂ [Co(CN) ₆]NO ₃ *2H ₂ O (tn=1,3-propanediamine). <i>Inorganic Chemistry</i> , 2001 , 40, 4839-44	5.1	38
267	Enhancing the reactivity of nickel(ii) in hydrogen evolution reactions (HERs) by hydrogenation of porphyrinoid ligands. <i>Chemical Science</i> , 2017 , 8, 5953-5961	9.4	37
266	Syntheses, Structures, and Characterization of Two Manganese(II)-Aminobenzoic Complexes. <i>European Journal of Inorganic Chemistry</i> , 2006 , 2006, 1649-1656	2.3	37
265	Towards 3-D Spherical Self-Assembly by Ternary Surfactant Combinations: The Case of Magnetite Nanoparticles. <i>European Journal of Inorganic Chemistry</i> , 2004 , 2004, 1169-1173	2.3	37
264	Hydrothermal syntheses, structures and magnetic properties of two transition metal coordination polymers with a square grid framework. <i>Polyhedron</i> , 2004 , 23, 1257-1262	2.7	37

263	Slow magnetic relaxation in a mixed-valence Mn(II/III) Complex: $[Mn^{II}_2(bispicen)_2(\mu-3-Cl)_2Mn^{III}(Cl_4Cat)_2Mn^{III}(Cl_4Cat)_2(H_2O)_2]$ infinity. <i>Inorganic Chemistry</i> , 2004 , 43, 849-51	5.1	37
262	Synthesis and characterization of substituted Schiff-base ligands and their d(10) metal complexes: structure-induced luminescence tuning behaviors and applications in co-sensitized solar cells. <i>Dalton Transactions</i> , 2015 , 44, 5306-22	4.3	36
261	Cooperative Crystallization of Chiral Heterometallic Indium(III) Potassium(I) Metal Organic Frameworks as Photosensitizers in Luminescence Sensors and Dye-Sensitized Solar Cells. <i>Crystal Growth and Design</i> , 2016 , 16, 1737-1745	3.5	36
260	Half-Sandwich Complexes of Dy(III): A Janus-Motif with Facile Tunability of Magnetism. <i>Inorganic Chemistry</i> , 2015 , 54, 5162-8	5.1	36
259	Lanthanide contraction and pH value controlled structural change in a series of rare earth complexes with p-aminobenzoic acid. <i>Journal of Molecular Structure</i> , 2004 , 702, 77-83	3.4	36
258	The Formation of Gd ₄ O ₄ cubane cluster controlled by L-valine. <i>New Journal of Chemistry</i> , 2000 , 24, 251-258		36
257	Assembling Dysprosium Dimer Units into a Novel Chain Featuring Slow Magnetic Relaxation via Formate Linker. <i>Inorganic Chemistry</i> , 2016 , 55, 12904-12911	5.1	36
256	Dramatic impact of the lattice solvent on the dynamic magnetic relaxation of dinuclear dysprosium single-molecule magnets. <i>Inorganic Chemistry Frontiers</i> , 2018 , 5, 1575-1586	6.8	35
255	NaCl-type frameworks of $[M(\text{pyrazine})_2\text{NO}_2]\text{ClO}_4$ (M = Co, Cu), the first examples containing $\mu(1,3)$ -nitrito bridges showing antiferromagnetism. <i>Inorganic Chemistry</i> , 2006 , 45, 9148-50	5.1	35
254	A novel 3D coordination polymer containing pentacoordinate cobalt(II) dimers with antiferromagnetic ordering. <i>Inorganic Chemistry Communication</i> , 2004 , 7, 864-867	3.1	35
253	An antiferromagnetic Mn(III) chain bridged by hydrogencyanamide: $[Mn^{III}(5\text{-Brsalen})(\mu 1,3\text{-NCNH})]_n$. <i>Inorganic Chemistry</i> , 2004 , 43, 8221-3	5.1	35
252	Low temperature synthesis and magnetism of La _{0.75} Ca _{0.25} MnO ₃ nanoparticles. <i>Journal of Physics and Chemistry of Solids</i> , 2000 , 61, 1407-1413	3.9	35
251	The first ladder-like lanthanide double chain complex $[Tm(\text{NTA})(\text{H}_2\text{O})_2]_2 \cdot 2\text{H}_2\text{O}$ (H ₃ NTA=nitritotriacetic acid). <i>Inorganic Chemistry Communication</i> , 2000 , 3, 319-321	3.1	35
250	A Variety of Phase-Transition Behaviors in a Niccolite Series of $[\text{NH}_3(\text{CH}_2)_4\text{NH}_3][\text{M}(\text{HCOO})_3]_2$. <i>Chemistry - A European Journal</i> , 2016 , 22, 6199-203	4.8	35
249	Observation of magnetic bistability in polymorphs of the $[\text{Ni}(\text{dmit})(2)](-)$ complexes. <i>Inorganic Chemistry</i> , 2009 , 48, 9623-30	5.1	34
248	Syntheses, crystal structures, and magnetic properties of two cyclic dimer M ₂ L ₂ complexes constructed from a new nitronyl nitroxide ligand and $M(\text{hfac})_2$ (M = Cu ²⁺ , Mn ²⁺). <i>Inorganica Chimica Acta</i> , 2007 , 360, 3553-3559	2.7	34
247	$\{[\text{Cu}_2(\text{bpdado})_2(\text{H}_2\text{O})_2] \cdot \text{H}_2\text{O}\}_n$: a 1D nanotubular coordination polymer with wall made of edge-sharing hexagons, where bpdado = 2,2'-bipyridine-3,3'-dicarboxylate-1,1'-dioxide. <i>Dalton Transactions</i> , 2005 , 2976-8	4.3	34
246	Synthesis, Crystal Structures and Magnetism of a Series of Heptanuclear Rare-Earth-Centered Trigonal Prism Clusters. <i>European Journal of Inorganic Chemistry</i> , 2003 , 2003, 731-736	2.3	34

- 245 Structural change of supramolecular coordination polymers of itaconic acid and 1,10-phenanthroline along lanthanide series. *Inorganica Chimica Acta*, **2005**, 358, 2687-2696 2.7 34
- 244 A copper-formate framework showing a simple to helical antiferroelectric transition with prominent dielectric anomalies and anisotropic thermal expansion, and antiferromagnetism. *Chemistry - A European Journal*, **2014**, 20, 15872-83 4.8 33
- 243 Copper(I)-iodide based coordination polymers: bifunctional properties related to thermochromism and PMMA-doped polymer film materials. *Journal of Materials Chemistry C*, **2015**, 3, 6249-6259 7.1 33
- 242 Chiral crystalline solids of ammonium-templated Er(III)formate frameworks assembled from three achiral acentric components. *CrystEngComm*, **2011**, 13, 5285 3.3 33
- 241 Synthesis, structure, and magnetic properties of (A)[Fe(III)(oxalate)Cl₂] (A = alkyl ammonium cations) with anionic 1D [Fe(III)(oxalate)Cl₂]- chains. *Inorganic Chemistry*, **2007**, 46, 3089-96 5.1 33
- 240 Three-dimensional heterometallic chiral Cr-Mn compound constructed by cyanide and dicyanamide bridges. *Inorganic Chemistry*, **2006**, 45, 10404-6 5.1 33
- 239 Syntheses, structure and magnetic properties of pillared layered diphosphonates: M₂(O₃PC₆H₄PO₃)(H₂O)₂ (M=Co(II), Ni(II)). *Journal of Solid State Chemistry*, **2004**, 177, 2311-2315 3.3 33
- 238 Peculiar Magnetic Transition Behaviour Around 208 K in One-dimensional Nickel(III) Chain Complex [1-(4'-Vinylbenzyl)pyridinium][Ni(mnt)₂] (mnt₂ = Maleonitriledithiolate). *Chemistry Letters*, **2005**, 34, 680-681 1.7 33
- 237 Crystal structure and luminescence property of ternary terbium p-aminobenzoic acid complexes with different second ligands. *Journal of Molecular Structure*, **2005**, 740, 175-180 3.4 33
- 236 New ternary lanthanide coordination polymers of 1,4-naphthalenedicarboxylate with 1,10-phenanthroline. *Inorganic Chemistry Communication*, **2005**, 8, 72-75 3.1 33
- 235 Highly stable open frameworks of Ln₂U (Ln=La, Ce) 3D coordination polymers containing spacious hexagon channels and unusual μ₂-H₂O bridges. *Inorganic Chemistry Communication*, **2001**, 4, 301-304 3.1 33
- 234 Tuning Slow Magnetic Relaxation in a Two-Dimensional Dysprosium Layer Compound through Guest Molecules. *Inorganic Chemistry*, **2016**, 55, 7980-7 5.1 33
- 233 Angular-resolved magnetometry beyond triclinic crystals part II: torque magnetometry of Cp*ErCOT single-molecule magnets. *Chemistry - A European Journal*, **2014**, 20, 14051-6 4.8 32
- 232 Cobalt diphosphonate with a new double chain structure exhibiting field-induced magnetic transition. *Dalton Transactions*, **2007**, 4681-4 4.3 32
- 231 Thermodynamics of spin-1/2 antiferromagnet-antiferromagnet-ferromagnet and ferromagnet-ferromagnet-antiferromagnet trimerized quantum Heisenberg chains. *Physical Review B*, **2006**, 73, 3.3 32
- 230 Three-Dimensional Lanthanoid-Containing Coordination Frameworks: Structure, Magnetic and Fluorescent Properties. *European Journal of Inorganic Chemistry*, **2005**, 2005, 766-772 2.3 32
- 229 Two mononuclear single molecule magnets derived from dysprosium(III) and tmphen (tmphen = 3,4,7,8-tetramethyl-1,10-phenanthroline). *Dalton Transactions*, **2015**, 44, 9020-6 4.3 31
- 228 Hierarchical cobalt-formate framework series with (412?63)(49?66)_n (n = 1B) topologies exhibiting slow dielectric relaxation and weak ferromagnetism. *APL Materials*, **2014**, 2, 124104 5.7 31

227	Syntheses, structures and magnetic properties of five iron(II) coordination polymers with flexible bis(imidazole) and bis(triazole) ligands. <i>Polyhedron</i> , 2012 , 31, 77-81	2.7	30
226	Magnetic nanoparticle-based cancer therapy. <i>Chinese Physics B</i> , 2013 , 22, 027506	1.2	30
225	Synthesis, crystal structure and magnetic properties of a CuII/IV bimetallic complex with a novel open framework structure. <i>Dalton Transactions</i> , 2003 , 3283-3287	4.3	30
224	The First Three-Leg Ladder Coordination Polymer: [Zn ₃ (Oac) ₄ (4,4'-bpy) ₃ {N(CN) ₂ }] ₂ . <i>Chemistry Letters</i> , 2001 , 30, 2-3	1.7	30
223	Novel bis(phthalocyaninato) rare earth complexes with the bulky and strong electron-donating dibutylamino groups: synthesis, spectroscopy, and SMM properties. <i>Inorganic Chemistry Frontiers</i> , 2017 , 4, 1465-1471	6.8	29
222	Irreversible transformation of chiral to achiral polymorph of K[Co(HCOO) ₃]: synthesis, structures, and magnetic properties. <i>Dalton Transactions</i> , 2011 , 40, 4465-73	4.3	29
221	Transition metal coordination frameworks with bridges of 1,2-bis(4-pyridyl)ethane-N,N'-dioxide incorporating anions of different size. <i>CrystEngComm</i> , 2008 , 10, 1796	3.3	29
220	Unexpected assembly of a cyanide-bridged one-dimensional nickel(II) complex with alternating [Ni(TIM)] ²⁺ and [Ni(CN) ₄] ²⁻ ions (TIM=2,3,9,10-tetramethyl-1,4,8,11-tetraazacyclotetradeca-1,3,8,10-tetraene). <i>Inorganic Chemistry Frontiers</i> , 2017 , 4, 1465-1471	3.1	29
219	Combined Effect of Hydrogen Bonding and π -Stacking Interactions in the Assembly of Indium(III) Metal-Organic Materials: Structure-Directing and Aggregation-Induced Emission Behavior. <i>Crystal Growth and Design</i> , 2015 , 15, 2402-2412	3.5	28
218	Linear trinuclear cobalt(II) single molecule magnet. <i>Dalton Transactions</i> , 2015 , 44, 2865-70	4.3	28
217	Effects of solvents and temperature on the luminescence properties of Cd-isonicotinic acid frameworks based on mono-, bi-, and trinuclear cluster units. <i>CrystEngComm</i> , 2014 , 16, 1113-1125	3.3	28
216	A novel tricyanoruthenium(III) building block for the construction of bimetallic coordination polymers. <i>Chemical Communications</i> , 2010 , 46, 6102-4	5.8	28
215	Stringing Oxo-Centered Trinuclear [MnIII ₃ O] Units into Single-Chain Magnets with Formate or Azide Linkers. <i>Angewandte Chemie</i> , 2007 , 119, 7532-7536	3.6	28
214	Crystal structure and magnetic behavior of a three-dimensional cyano-bridged assembly [CuL ₁] ₂ [Cr(CN) ₆]ClO ₄ ·0.5H ₂ O (L ₁ = 3,10-dipropyl-1,3,5,8,10,12-hexaazacyclotetradecane). <i>New Journal of Chemistry</i> , 2004 , 28, 996-999	3.6	28
213	A convenient solvothermal route to ruthenium nanoparticles. <i>New Journal of Chemistry</i> , 2000 , 24, 739-740	3.0	28
212	Enhanced magnetic anisotropy in a tellurium-coordinated cobalt single-ion magnet. <i>Inorganic Chemistry Frontiers</i> , 2017 , 4, 701-705	6.8	27
211	Slow magnetic relaxation in a mononuclear 8-coordinate Fe(II) complex. <i>Chemical Communications</i> , 2017 , 53, 1474-1477	5.8	27
210	Qubit crossover in the endohedral fullerene ScC@C. <i>Chemical Science</i> , 2018 , 9, 457-462	9.4	27

- 209 FePt concave nanocubes with enhanced methanol oxidation activity. *CrystEngComm*, **2012**, 14, 7572 3.3 27
- 208 Synthesis, structures, and magnetism of three 1D Mn(III) chains with oxazoline-based ligands. *Inorganic Chemistry*, **2010**, 49, 5868-75 5.1 27
- 207 Syntheses, crystal structures, and spectroscopic and magnetic properties of [Mn²⁺(H₂L¹)(Cl₄Cat)·4.2H₂O] and [Mn²⁺(H₂L²)(Cl₄Cat)·4.2CH₃CN·2H₂O]_n: temperature-dependent valence tautomerism in solution. *Inorganic Chemistry*, **2005**, 44, 9714-22 5.1 27
- 206 Synthesis, Crystal Structure, and Magnetic Properties of a Three-Dimensional Cyano-Bridged Bimetallic Coordination Polymer with an Aromatic Amine Capping Ligand: [Cu(2,2′-bpa)]₃[Cr(CN)₆]₂·2H₂O (2,2′-bpa = 2,2′-Dipicolylamine). *Crystal Growth and Design*, **2006**, 6, 94-98 3.5 27
- 205 Syntheses, structures and magnetic properties of two novel 3D helical dicyanamide complexes containing polyamine ligand. *Inorganic Chemistry Communication*, **2004**, 7, 669-672 3.1 27
- 204 A Novel One-Dimensional Chain Cyano-Bridged Complex [Sm(DMF)₄(H₂O)₂Mn(CN)₆·nH₂O]_n: Long-Range Magnetic Ordering, T_c = 18 K and Coercive Force H_c = 600 Oe. *Chemistry Letters*, **2001**, 30, 350-351 1.7 27
- 203 Crystal structures and magnetic properties of two octacyanometalate-based tungstate(V)–copper(II) bimetallic assemblies. *New Journal of Chemistry*, **2002**, 26, 1190-1195 3.6 27
- 202 Zinc-diluted magnetic metal formate perovskites: synthesis, structures, and magnetism of [CH₃NH₃][Mn(x)Zn(1-x)(HCOO)₃] (x = 0-1). *Chemistry - an Asian Journal*, **2012**, 7, 1697-707 4.5 26
- 201 [NiL¹]₃[Cr(CN)₆]₂·18H₂O [L¹ = 3,10-bis(2-hydroxyethyl)-1,3,5,8,10,12-hexaazacyclotetradecane]: a two-dimensional bimetallic assembly exhibiting antiferromagnetic ordering and metamagnetic behavior. *New Journal of Chemistry*, **2001**, 25, 875-878 3.6 26
- 200 Lanthanide dinuclear complexes constructed from mixed oxygen-donor ligands: the effect of substituent positions of the neutral ligand on the magnetic dynamics in Dy analogues. *Dalton Transactions*, **2016**, 45, 4614-21 4.3 25
- 199 A family of one-dimensional lanthanide complexes bridged by two distinct carboxylate ligands with the Dy analogue displaying magnetic relaxation behaviour. *Dalton Transactions*, **2017**, 46, 14114-14121 4.3 25
- 198 Two EE-azido-bridged nickel(II) layered compounds: vigorous twisted torsion angle Ni-N₃-Ni results in ferromagnetic behavior. *Inorganic Chemistry*, **2009**, 48, 7174-80 5.1 25
- 197 A Novel Acentric Three-Dimensional Network with Complicated Topology. *Crystal Growth and Design*, **2006**, 6, 1261-1263 3.5 25
- 196 Preparation of three terbium complexes with p-aminobenzoic acid and investigation of crystal structure influence on luminescence property. *Journal of Solid State Chemistry*, **2004**, 177, 3735-3742 3.3 25
- 195 A Series of Bimetallic Ammonium AlNa Formates. *Chemistry - A European Journal*, **2017**, 23, 9857-9871 4.8 24
- 194 Advances in Lanthanide Single-Ion Magnets. *Structure and Bonding*, **2014**, 111-141 0.9 24
- 193 Understanding the magnetic anisotropy in a family of N₂(3-) radical-bridged lanthanide complexes: density functional theory and ab initio calculations. *Journal of Physical Chemistry A*, **2013**, 117, 10873-80 2.8 24
- 192 Two Ferromagnetic One-Dimensional Nickel(II) Complexes Containing the Diamagnetic [Ni(CN)₄]²⁻ Group. *European Journal of Inorganic Chemistry*, **2002**, 2002, 699-702 2.3 24

191	A novel molecular ladder structure of Cu(II)-Ba(II) coordination polymer exhibiting ferromagnetic coupling. <i>Chemical Communications</i> , 2003 , 2946-7	5.8	24
190	Co ₃ {CH ₃ C(NH ₃)(PO ₃ H)(PO ₃) ₂ } ₂ {CH ₃ C(NH ₃)(PO ₃ H) ₂ } ₂ (H ₂ O) ₄ ·2H ₂ O: A cobalt 1-aminoethylidenediphosphonate with a layer structure. <i>Journal of Solid State Chemistry</i> , 2005 , 178, 1049-1053 ²⁴	3.3	24
189	Observation of the asphericity of 4f-electron density and its relation to the magnetic anisotropy axis in single-molecule magnets. <i>Nature Chemistry</i> , 2020 , 12, 213-219	17.6	24
188	A Six-Coordinate Dysprosium Single-Ion Magnet with Trigonal-Prismatic Geometry. <i>Inorganic Chemistry</i> , 2017 , 56, 7320-7323	5.1	23
187	Slow Magnetic Relaxation in Weak Easy-Plane Anisotropy: the Case of a Combined Magnetic and HFEP Study. <i>Inorganic Chemistry</i> , 2017 , 56, 697-700	5.1	23
186	Constructing a Series of Azide-Bridged CuII Magnetic Low-Dimensional Coordination Polymers by using Pybox Ligands. <i>European Journal of Inorganic Chemistry</i> , 2013 , 2013, 3101-3111	2.3	23
185	trans-[Os(III)(salen)(CN) ₂] ⁻ : a new paramagnetic building block for the construction of molecule-based magnetic materials. <i>Inorganic Chemistry</i> , 2010 , 49, 1607-14	5.1	23
184	Thermodynamics of spin-12 tetrameric Heisenberg antiferromagnetic chain. <i>Physical Review B</i> , 2009 , 80,	3.3	23
183	Synthesis, crystal structure, EPR and magnetic properties of a cyano-bridged CuII/NiII heterobimetallic complex: an unusual structure with long-range ferromagnetic exchange through hydrogen bonding. <i>New Journal of Chemistry</i> , 2004 , 28, 412-417	3.6	23
182	Structures and magnetism of bimetallic octacyanometallates: honeycomb-like [Mo(CN) ₈ Ni ₂ (pn) ₄] _n ·4nH ₂ O and binuclear [Mo(CN) ₈ Ni(en) ₂ (H ₂ O)] ₂ [Ni(en) ₃] ₂ ·2H ₂ O (pn=1,3-diaminopropane, en=1,2-diaminoethane). <i>Inorganic Chemistry Communication</i> , 2002 , 5, 660-663	3.1	23
181	Probing the influence of molecular symmetry on the magnetic anisotropy of octahedral cobalt(II) complexes. <i>Inorganic Chemistry Frontiers</i> , 2017 , 4, 1909-1916	6.8	22
180	Spin transitions in a series of [Fe(pybox) ₂] ²⁺ complexes modulated by ligand structures, counter anions, and solvents. <i>Inorganic Chemistry Frontiers</i> , 2016 , 3, 1624-1636	6.8	22
179	Evaporable lanthanide single-ion magnet. <i>CrystEngComm</i> , 2016 , 18, 4165-4171	3.3	22
178	Syntheses and Structures of Layered Copper(II) Diphosphonates with Mixed Ligands. <i>European Journal of Inorganic Chemistry</i> , 2003 , 2003, 726-730	2.3	22
177	Preparation of nanosized La ₂ CuO ₄ perovskite oxide using an amorphous heteronuclear complex as a precursor at low-temperature. <i>Journal of Alloys and Compounds</i> , 2000 , 311, 16-21	5.7	22
176	Synthesis and structure of an unprecedented heterometallic tetranuclear cluster [Gd ₂ Co ₂ (β-O)(H ₂ O)(CDTA) ₂ (H ₂ O) ₆] ₂ ·10H ₂ O (H ₄ CDTA = trans-1,2-cyclohexanedinitrilotetraacetic acid). <i>Dalton Transactions RSC</i> , 2000 , 1003-1004		22
175	A 36-Fold Multiple Unit Cell and Switchable Anisotropic Dielectric Responses in an Ammonium Magnesium Formate Framework. <i>Angewandte Chemie</i> , 2015 , 127, 2564-2567	3.6	21
174	Tunable Luminescence and Application in Dye-Sensitized Solar Cells of Zn(II)/Hg(II) Complexes: Methyl Substitution-Induced Supramolecular Structures Based on (E)-N-(6-Methoxypyridin-2-ylmethylene)arylamine Derivatives. <i>Inorganic Chemistry</i> , 2015 , 54, 7742-52	5.1	21

- 173 Thermal and light induced spin crossover behavior of a dinuclear Fe(II) compound. *Dalton Transactions*, **2015**, 44, 11282-5 4.3 21
- 172 Synthesis, crystal structure and effect of deuterated solvents and temperature on visible and near infrared luminescence of N4-donor Schiff base lanthanide complexes. *Journal of Luminescence*, **2014**, 149, 75-85 3.8 21
- 171 A family of enantiopure Fe(III)₄ single molecule magnets: fine tuning of energy barrier by remote substituent. *Dalton Transactions*, **2014**, 43, 11897-907 4.3 21
- 170 Polymorphism of (H₂mela)₂[CuCl₅]Cl (mela = melamine): structures, transformation and magnetic properties. *CrystEngComm*, **2011**, 13, 4683 3.3 21
- 169 Spin-canting in a 1D chain Mn(II) complex with alternating double end-on and double end-to-end azido bridging ligands. *Inorganic Chemistry Communication*, **2007**, 10, 897-901 3.1 21
- 168 Structures and Magnetic Properties of Iron Diphosphonates: [NH₃(CH₂)₂NH₃][FeII{HO₃PC(CH₃)(OH)(PO₃H)}₂]**2**H₂O and [NH₃(CH₂)₂NH₂(CH₂)₂NH₃][FeIII{O₃PC(CH₃)(OH)(PO₃H)}₂]**1**H₂O. *Journal of Solid State Chemistry*, **2008**, 174, 347-353 3.3 21
- 167 Solvothermal synthesis, crystal structure and magnetic property of a new dinuclear manganese(II) azido complex: [Mn(2,2'-dpa)(N₃)₂]₂ (2,2'-dpa = 2,2'-dipicolylamine). *Inorganica Chimica Acta*, **2005**, 358, 834-838 2.7 21
- 166 Magnetic properties of J₁J₂ quantum Heisenberg chains with spin S = 1/2, 1, 3/2 and 2 in a magnetic field. *Journal of Physics Condensed Matter*, **2005**, 17, 6081-6094 1.8 21
- 165 Structural phase transition in a multi-induced mononuclear Fe(II) spin-crossover complex. *Dalton Transactions*, **2015**, 44, 20906-12 4.3 20
- 164 Slow Magnetic Relaxation in a Series of Mononuclear 8-Coordinate Fe(II) and Co(II) Complexes. *Inorganic Chemistry*, **2018**, 57, 3761-3774 5.1 20
- 163 Controllable synthesis of Zn/Cd(ii) coordination polymers: dual-emissive luminescent properties, and tailoring emission tendency under varying excitation energies. *Dalton Transactions*, **2016**, 45, 4863-4873 4.3 20
- 162 The synthesis, structures and magnetic properties of polynuclear Ru(III)-3d (3d = Mn(II/III), Ni(II), Cu(II)) compounds based on [Ru(III)(Q)₂(CN)₂]⁻. *Dalton Transactions*, **2013**, 42, 3876-87 4.3 20
- 161 A oximato-bridged linear trinuclear [Mn(IV)Mn(III)Mn(IV)] single-molecule magnet. *Dalton Transactions*, **2012**, 41, 13620-5 4.3 20
- 160 Static field induced magnetic relaxations in dinuclear lanthanide compounds of [phen₂Ln₂(HCOO)₄(HCOO)₂]**x** (NO₃)₂**x**] (1, Ln = Gd and x = 0.52; 2, Ln = Er and x = 0.90; phen = 1,10-phenanthroline). *Science China Chemistry*, **2012**, 55, 926-933 7.9 20
- 159 Cyano-bridged molecular squares: Synthesis and structures of [Ni(cyclen)]₂[Pt(CN)₄]₂ **6**H₂O, [Ni(cyclen)]₂[Ni(CN)₄]₂ **6**H₂O and [Mn(cyclen)]₂[Ni(CN)₄]₂ **6**H₂O. *Polyhedron*, **2006**, 25, 1256-1262 2.7 20
- 158 Perovskite-Like Polar Lanthanide Formate Frameworks of [NHNH][Ln(HCOO)] (Ln = Tb-Lu and Y): Synthesis, Structures, Magnetism, and Anisotropic Thermal Expansion. *Inorganic Chemistry*, **2016**, 55, 10075-10082 5.1 20
- 157 A distinct magnetic anisotropy enhancement in mononuclear dysprosium-sulfur complexes by controlling the Dy-ligand bond length. *Dalton Transactions*, **2016**, 45, 8149-53 4.3 20
- 156 Modulating Slow Magnetic Relaxation of Dysprosium Compounds through the Position of Coordinating Nitrate Group. *Inorganic Chemistry*, **2017**, 56, 13430-13436 5.1 19

155	Four Dinuclear and One-Dimensional-Chain Dysprosium and Terbium Complexes Based on 2-Hydroxy-3-methoxybenzoic Acid: Structures, Fluorescence, Single-Molecule-Magnet, and Ab Initio Investigation. <i>Inorganic Chemistry</i> , 2020 , 59, 4414-4423	5.1	19
154	Two unusual organic/inorganic hybrid 3-D frameworks based on Keggin-type heteropoly blue anion-chains, 40-membered macrocycles, and sodium linker units. <i>CrystEngComm</i> , 2014 , 16, 8449	3.3	19
153	An A-Site Mixed-Ammonium Solid Solution Perovskite Series of [(NH ₂ NH ₃) _x (CH ₃ NH ₃) _{1-x}][Mn(HCOO) ₃] (x=1.000-0.67). <i>Angewandte Chemie</i> , 2015 , 127, 11245-11248	3.6	19
152	Influence of Guest Exchange on the Magnetization Dynamics of Dilanthanide Single-Molecule-Magnet Nodes within a Metal/Organic Framework. <i>Angewandte Chemie</i> , 2015 , 127, 9999-10003	3.6	19
151	Field-induced slow magnetic relaxation in a hydrogen-bonding linked Co(II) 1D supramolecular coordination polymer. <i>Supramolecular Chemistry</i> , 2015 , 27, 401-406	1.8	19
150	Cobalt(II) Coordination Polymers Containing trans-1,2-Bis(4-pyridyl)ethene and Their Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2010 , 2010, 3856-3865	2.3	19
149	Two diammonium copper azides with similar layerlike magnetic substructures made of chains of serially connected Cu ₆ rings show cation-modulated magnetism. <i>Chemistry - an Asian Journal</i> , 2008 , 3, 950-7	4.5	19
148	Bis(ethylenedithio)tetrathiafulvalene Radical Salts with Anderson Type Heteropolymolybdates Containing Tris(alkoxo) Ligands. <i>Crystal Growth and Design</i> , 2005 , 5, 1531-1538	3.5	19
147	Construction and theoretical study of a new Dy(III)-diketone chain featuring slow magnetic relaxation. <i>CrystEngComm</i> , 2015 , 17, 5620-5624	3.3	18
146	Structure variations of a series of lanthanide complexes constructed from quinoline carboxylate ligands: photoluminescent properties and PMMA matrix doping. <i>RSC Advances</i> , 2015 , 5, 38254-38263	3.7	18
145	Getting Insights into the Temperature-Specific Active Sites on Platinum Nanoparticles for CO Oxidation: A Combined in Situ Spectroscopic and ab Initio Density Functional Theory Study. <i>ACS Catalysis</i> , 2019 , 9, 7759-7768	13.1	18
144	M ₂ (N ₃) ₄ (hmt)(H ₂ O) (M = Co ²⁺ and Ni ²⁺ , hmt = hexamethylenetetramine): mixed azide-hmt bridged 3D metal frameworks with long-range magnetic ordering. <i>CrystEngComm</i> , 2009 , 11, 2096	3.3	18
143	Two-dimensional rare earth coordination polymers involving different coordination modes of thiodiglycolic acid. <i>Inorganic Chemistry Communication</i> , 2002 , 5, 28-31	3.1	18
142	Cyano-Bridged 4f/3d Coordination Polymers with a Unique Two-Dimensional Topological Architecture and Unusual Magnetic Behavior. <i>Angewandte Chemie</i> , 2001 , 113, 448-451	3.6	18
141	The solvent effect in an axially symmetric Fe(III) ₄ single-molecule magnet. <i>Chemical Communications</i> , 2014 , 50, 15090-3	5.8	17
140	Metal monophosphonates M{(2-C ₅ H ₄ NO)CH ₂ PO ₃ }(H ₂ O) ₂ (M = Co, Ni, Mn, Cd): synthesis, structure, and magnetism. <i>Inorganic Chemistry</i> , 2010 , 49, 8474-80	5.1	17
139	Bimetallic sandwiches assembled with chelated Cu/Zn cations and manganese dicyanamide polymeric ladders. <i>Polyhedron</i> , 2003 , 22, 433-439	2.7	17
138	Structure and Magnetism of Cyano-Bridged Hetero-Binuclear Complexes RE(dpdo) ₂ (H ₂ O) ₃ Fe(CN) ₆ ·4H ₂ O (RE=Gd, Y). <i>Molecular Crystals and Liquid Crystals</i> , 1999 , 335, 201-210		17

137	Experimental Determination of Magnetic Anisotropy in Exchange-Bias Dysprosium Metallocene Single-Molecule Magnets. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 13037-13043	16.4	16
136	An insight into the controllable synthesis of Cd(II) complexes with a new multifunctional ligand and its application in dye-sensitized solar cells and luminescence properties. <i>Journal of Materials Chemistry A</i> , 2015 , 3, 6053-6063	13	16
135	Novel heterobimetallic ruthenium(III)-cobalt(II) compounds constructed from trans-[Ru(III)(Q) ₂ (CN) ₂] ⁻ (Q = 8-quinolinolato): synthesis, structures and magnetic properties. <i>Chemical Communications</i> , 2011 , 47, 8694-6	5.8	16
134	Synthesis, structural characterization, and magnetism of a butterfly-shaped hexanuclear Ni(II) complex. <i>Inorganic Chemistry Communication</i> , 2008 , 11, 769-771	3.1	16
133	Synthesis, crystal structure and magnetic properties of μ -oxo-centered tetranuclear lanthanide clusters. <i>New Journal of Chemistry</i> , 2001 , 25, 531-533	3.6	16
132	Rational design of an "all-in-one" phototheranostic. <i>Chemical Science</i> , 2020 , 11, 8204-8213	9.4	16
131	Magnetic and HFEPR Studies of Exchange Coupling in a Series of ECL Dicobalt Complexes. <i>Inorganic Chemistry</i> , 2017 , 56, 2417-2425	5.1	15
130	[(C ₂ H ₅) ₄ N][U ₂ O ₄ (HCOO) ₅], an ammonium uranyl formate framework showing para- to ferro-electric transition: synthesis, structures, and properties. <i>Inorganic Chemistry</i> , 2014 , 53, 8708-16	5.1	15
129	Phase transitions and thermodynamics of the two-dimensional Ising model on a distorted kagome lattice. <i>Physical Review B</i> , 2010 , 82,	3.3	15
128	Magnetic properties tuned by oxamido bridging ligand derivatives in two new hybrid organic inorganic nitronyl nitroxide copper(II) complexes. <i>CrystEngComm</i> , 2007 , 9, 799	3.3	15
127	Synthesis, structure and magnetic properties of one chiral heptanuclear Fe cluster: {Fe[(FeL) ₂ (μ -OH)] ₃ (μ -OH) ₆ }] \cdot 3H ₂ O. <i>Inorganic Chemistry Communication</i> , 2008 , 11, 256-259	3.1	15
126	Synthesis and characterization of new polynuclear lanthanide coordination polymers with 4,4'-oxybis(benzoic acid). <i>Polyhedron</i> , 2005 , 24, 823-830	2.7	15
125	Synthesis, structure and magnetic studies of two-dimensional network of Cu(II) polymer using malonate and 4,4'-azobispyridine ligands. <i>Inorganica Chimica Acta</i> , 2005 , 358, 3519-3524	2.7	15
124	Advanced Cd(II) complexes as high efficiency co-sensitizers for enhanced dye-sensitized solar cell performance. <i>Dalton Transactions</i> , 2015 , 44, 18187-95	4.3	14
123	In situ tetrazole templated chair-like decanuclear azido-cobalt(II) SMM containing both tetra- and octa-hedral Co(II) ions. <i>Dalton Transactions</i> , 2015 , 44, 480-3	4.3	14
122	Tri-Manganese(III) Salen-Based Cryptands: A Metal Cooperative Antioxidant Strategy that Overcomes Ischemic Stroke Damage. <i>Journal of the American Chemical Society</i> , 2020 , 142, 10219-10227	16.4	14
121	Low-Coordinate Single-Ion Magnets by Intercalation of Lanthanides into a Phenol Matrix. <i>Angewandte Chemie</i> , 2018 , 130, 4763-4766	3.6	14
120	Alkylamine-Templated Niccolite Frameworks of [GaM(HCOO)] (M = Fe, Ni): Structure, Magnetism, and Dielectricity. <i>Inorganic Chemistry</i> , 2018 , 57, 3941-3947	5.1	14

119	Elucidation of the two-step relaxation processes of a tetranuclear dysprosium molecular nanomagnet through magnetic dilution. <i>Dalton Transactions</i> , 2018 , 47, 11636-11644	4.3	14
118	Two Mn ₆ single-molecule magnets with sulfur-contained capping ligand. <i>Polyhedron</i> , 2011 , 30, 3279-3283	3.7	14
117	Synthesis, structure and magnetic properties of carboxylate-bridged polynuclear copper(II)lanthanide(III) rings: [Ln(H ₂ O) ₅ (CuL) ₂][CuL]·8H ₂ O (Ln = La, Ce). <i>Polyhedron</i> , 2004 , 23, 2253-2258	2.7	14
116	Two novel types of Ln ₂ M ₂ clusters bridged by cyanide and carboxylate simultaneously. <i>Journal of Solid State Chemistry</i> , 2003 , 171, 201-207	3.3	14
115	Unusual molecular ladder structure containing zinc(II) and natural T-shape ligand nitrilotriacetate. <i>Inorganic Chemistry Communication</i> , 2001 , 4, 79-81	3.1	14
114	Spin Frustrated Double Chain {[Fe(PZDA)H ₂ O] ₂ ·2H ₂ O} _n (H ₂ PZDA=2,3-Pyrazinedicarboxylic Acid). <i>Chemistry Letters</i> , 1999 , 28, 773-774	1.7	14
113	Rational construction of a porous lanthanide coordination polymer featuring reversible guest-dependent magnetic relaxation behavior. <i>Inorganic Chemistry Frontiers</i> , 2018 , 5, 2875-2884	6.8	14
112	Determination of zero-field splitting in Co halide complexes with magnetic and far-IR measurements. <i>Dalton Transactions</i> , 2017 , 46, 7408-7411	4.3	13
111	Dysprosium complexes bearing unsupported Dy-Ge/Sn metal-metal bonds as single-ion magnets. <i>Chemical Communications</i> , 2019 , 55, 8250-8253	5.8	13
110	Reactivity of a Two-Coordinate Cobalt(0) Cyclic (Alkyl)(amino)carbene Complex. <i>Organometallics</i> , 2020 , 39, 729-739	3.8	13
109	Unprecedented one-dimensional chain and two-dimensional network dysprosium(III) single-molecule toroids with white-light emission. <i>Chemical Communications</i> , 2020 , 56, 2590-2593	5.8	13
108	Nanomagnetism: Principles, nanostructures, and biomedical applications. <i>Chinese Physics B</i> , 2014 , 23, 057505	1.2	13
107	(H ₂ mela) ₂ [FeCl ₅]Cl (mela=melamine): A Cl-bridged single-chain magnet based on weak ferromagnetism. <i>Coordination Chemistry Reviews</i> , 2013 , 257, 2484-2490	23.2	13
106	Coordination Polymers Constructed from [Mn(N)(CN) ₄] ²⁻ Synthesis, Structures, and Magnetic Properties. <i>European Journal of Inorganic Chemistry</i> , 2008 , 2008, 158-163	2.3	13
105	Hydrothermal syntheses, crystal structures and magnetic properties of two inorganic-organic hybrid materials: [{Cu(phen)} ₂ (VVO ₂) ₂ (VVO ₂ (H ₂ O)(PO ₄) ₂)] and [V ₄ O ₇ (2,2'-bpy) ₂ (HPO ₄) ₂] (phen=1,10-phenanthroline, bpy=bipyridine). <i>Journal of Solid State Chemistry</i> , 2003 , 176, 69-75	3.3	13
104	A rare chloride-bridged dysprosium chain with slow magnetic relaxation: a thermally activated mechanism via a second-excited state promoted by magnetic interactions. <i>Inorganic Chemistry Frontiers</i> , 2019 , 6, 786-790	6.8	12
103	Multifunctional Zn(II)/Cd(II) metal complexes for tunable luminescence properties and highly efficient dye-sensitized solar cells. <i>RSC Advances</i> , 2015 , 5, 43705-43716	3.7	12
102	Magnetic anisotropy investigation on light lanthanide complexes. <i>Dalton Transactions</i> , 2018 , 47, 1966-1971	4.1	12

- 101 Structural Distortion Controlled Spin-Crossover Behavior. *Crystal Growth and Design*, **2015**, 15, 2565-2567. 3.5 12
- 100 Chemical synthesis of magnetic nanocrystals: Recent progress. *Chinese Physics B*, **2013**, 22, 107503 1.2 11
- 99 Weak ferromagnetism in chiral diamond-like neutral networks: Mn(2-pymS)₂ and Co(2-pymS)₂ (2-pymSH = 2-mercaptopyrimidine). *Dalton Transactions*, **2012**, 41, 2626-31 4.3 11
- 98 A novel triazidoruthenium(III) building block for the construction of polynuclear compounds. *Dalton Transactions*, **2012**, 41, 5794-8 4.3 11
- 97 Chlorido-Bridged MnII Schiff-Base Complex with Ferromagnetic Exchange Interactions. *European Journal of Inorganic Chemistry*, **2007**, 2007, 3663-3668 2.3 11
- 96 Formation of Two-Dimensional Supramolecular Icelike Layer Containing (H₂O)₁₂ Rings. *Angewandte Chemie*, **2004**, 116, 1398-1400 3.6 11
- 95 Synthesis, Structure and Magnetic Properties of a μ -Carboxylato-Bridged Ferromagnetic Trinuclear Copper(II) Complex with Lariat Tetraazacrown Ether as Ligand. *European Journal of Inorganic Chemistry*, **2004**, 2004, 1311-1317 2.3 11
- 94 The highly stable open frameworks and the adsorption performance of a 3D ErIII coordination polymer containing spacious channels. *Inorganic Chemistry Communication*, **2003**, 6, 841-844 3.1 11
- 93 Vertex-Sharing Water Tape Consisting of Cyclic Hexamers. *European Journal of Inorganic Chemistry*, **2005**, 2005, 3902-3906 2.3 11
- 92 Two Molecular Tapes Consisting of Serial or Parallel Azido-Bridged Eight-Membered Copper Rings. *Angewandte Chemie*, **2005**, 117, 5991-5996 3.6 11
- 91 Regulating the structural dimensionality and dynamic properties of a porous dysprosium coordination polymer through solvent molecules. *Inorganic Chemistry Frontiers*, **2020**, 7, 930-938 6.8 11
- 90 Three New Niccolites: High-Temperature Phase Transitions, Prominent Anisotropic Thermal Expansions, Dielectric Anomalies, and Magnetism. *Chemistry - A European Journal*, **2019**, 25, 9303-9314 4.8 10
- 89 Magnetic layered perovskites of [CHC(NH)][M(HCOO)] (M = Co and Ni): synthesis, structures and properties. *Dalton Transactions*, **2018**, 47, 11925-11933 4.3 10
- 88 Crystal structure and magnetic property of a metal-organic framework (MOF) containing double-stranded chain with metallomacrocycles and dinuclear Mn(II) subunits. *Journal of Molecular Structure*, **2008**, 891, 357-363 3.4 10
- 87 Construction of a Novel 2D Polymer [Co(dmpzm)(dca)₂] [From Reaction of a Mononuclear Complex [Co(dmpzm)Cl₂] with Sodium Dicyanamide (dca) [dmpzm=bis(3,5-dimethylpyrazolyl)methane]. *Chinese Journal of Chemistry*, **2006**, 24, 1716-1720 4.9 10
- 86 Enhanced magnetoresistance in La_{0.7}Sr_{0.3}MnO₃ nanoscaledgranular composites. *Science in China Series B: Chemistry*, **2000**, 43, 561-566 10
- 85 Field-Dependent Magnetic Relaxation and Magnetocaloric Effect in Mononuclear Gd Complexes. *Acta Chimica Sinica*, **2013**, 71, 1022 3.3 10
- 84 A Gallium(III) Complex that Engages Protein Disulfide Isomerase A3 (PDIA3) as an Anticancer Target. *Angewandte Chemie - International Edition*, **2020**, 59, 20147-20153 16.4 10

83	Dy@CN: a new member of dimetalloazafullerenes with strong single molecular magnetism. <i>Nanoscale</i> , 2020 , 12, 11130-11135	7.7	9
82	Electric field manipulation enhanced by strong spin-orbit coupling: promoting rare-earth ions as qubits. <i>National Science Review</i> , 2020 , 7, 1557-1563	10.8	9
81	Thermally-induced single-crystal-to-single-crystal transformations from a 2D two-fold interpenetrating square lattice layer to a 3D four-fold interpenetrating diamond framework and its application in dye-sensitized solar cells. <i>Physical Chemistry Chemical Physics</i> , 2016 , 18, 19001-10	3.6	9
80	A new Fe(6) ferric cluster: synthesis, crystal structure and magnetic properties. <i>Dalton Transactions</i> , 2008 , 6948-52	4.3	9
79	pH-Directed assembly and magnetic properties of two polynuclear MnII complexes: $[Mn_3(phen)_2(OOCCH_3)_6]$ and 1-D $[Mn(phen)Cl_2]_n$. <i>Polyhedron</i> , 2008 , 27, 3669-3673	2.7	9
78	Syntheses, spectra, crystal structures and magnetic properties of molecular solids based on the ion (M=Ni, Pt). <i>Polyhedron</i> , 2005 , 24, 1669-1676	2.7	9
77	1-D polymer containing the [Ru-N-Ru] mu-nitrido moiety: crystal structure and magnetic properties of $([Cu(en)_2]_3[Ru_2N(CN)_{10}].ClO_4)_n$ (en = 1,2-diaminoethane). <i>Chemical Communications</i> , 2002 , 2090-1	5.8	9
76	Weak exchange coupling effects leading to fast magnetic relaxations in a trinuclear dysprosium single-molecule magnet. <i>Inorganic Chemistry Frontiers</i> , 2020 , 7, 447-454	6.8	9
75	Temperature-Induced Irreversible Phase Transition From Perovskite to Diamond But Pressure-Driven Back-Transition in an Ammonium Copper Formate. <i>Angewandte Chemie</i> , 2016 , 128, 2137-2140	2.6	8
74	Two trihydrazine-bridged cobalt(II) chain compounds exhibiting antiferromagnetic ordering and single-chain magnetic behavior. <i>Chemistry - an Asian Journal</i> , 2014 , 9, 2463-6	4.5	8
73	Evolution of molecular nanomagnets in China. <i>Philosophical Transactions Series A, Mathematical, Physical, and Engineering Sciences</i> , 2013 , 371, 20120316	3	8
72	Supramolecular self-assembly of 1D and 3D heterometallic coordination polymers with triruthenium building blocks. <i>Chemistry - A European Journal</i> , 2012 , 18, 11228-37	4.8	8
71	Detailed magnetic study on the formate-bridged MOFs with anion-tunable magnetic properties. <i>Science China Chemistry</i> , 2012 , 55, 1055-1063	7.9	8
70	Lanthanide Based Magnetic Molecular Materials 2010 , 355-405		8
69	Synthesis and characterization of a new iron(III) oxide cluster containing two different discrete $[Fe_3O]$ units: $[Fe_3O(O_2CCH_2CH_2)_6(H_2O)_3]$ and $[Fe_3O(O_2CCH_2CH_2)_7(H_2O)_2(OH_2CCH_2CH_2)_4(OH)(H_2O)]$. <i>Inorganic Chemistry Communication</i> , 2004 , 7, 584-587	3.1	8
68	Ferromagnetic Ordering in a Diamond-Like Cyano-Bridged MnII/RuIII Bimetallic Coordination Polymer. <i>Angewandte Chemie</i> , 2001 , 113, 3121-3123	3.6	8
67	A One-dimensional Nickel(III) Chain Complex Showing Ferromagnetic Ordering: Crystal Structure and Magnetic Property. <i>Chemistry Letters</i> , 2002 , 31, 576-577	1.7	8
66	Ferromagnetic Ordering in a Two-Dimensional Copper Complex with Dual End-to-End and End-On Azide Bridges. <i>Angewandte Chemie</i> , 2000 , 112, 3779-3781	3.6	8

65	Understanding the near-infrared fluorescence and field-induced single-molecule-magnetic properties of dinuclear and one-dimensional-chain ytterbium complexes based on 2-hydroxy-3-methoxybenzoic acid. <i>Inorganic Chemistry Frontiers</i> , 2020 , 7, 3136-3145	6.8	8
64	Two Magnetic Switching Complexes Based on the Fe(II) Ion. <i>Inorganic Chemistry</i> , 2016 , 55, 7805-7	5.1	8
63	Adducts of Tris(alkyl) Holmium(III) Showing Magnetic Relaxation. <i>Inorganic Chemistry</i> , 2020 , 59, 5835-5844	4.1	8
62	Two-step magnetic switching in a mononuclear iron(II) complex around room temperature. <i>Dalton Transactions</i> , 2015 , 44, 8938-41	4.3	7
61	A Pair of Enantiopure Cubane-Type CuII4O4 Clusters: Synthesis, Structure, Chirality and Magnetism. <i>European Journal of Inorganic Chemistry</i> , 2014 , 2014, 5385-5390	2.3	7
60	NiIIrIII and CuIIrIII Coordination Polymers Constructed from [Ru(CN)6]4- <i>European Journal of Inorganic Chemistry</i> , 2005 , 2005, 364-370	2.3	7
59	Structural, magnetic, and transport properties in La0.7Ca0.3Mn1-xScxO3. <i>Journal of Applied Physics</i> , 2001 , 90, 4609-4613	2.5	7
58	A novel one-dimensional Ni(II)Be(II) polymer containing β -cyanides: [Ni(cyclen)]2[Fe(CN)6]·8H2O. <i>New Journal of Chemistry</i> , 2002 , 26, 1099-1101	3.6	7
57	Single-Crystal Study of a Low Spin Co(II) Molecular Qubit: Observation of Anisotropic Rabi Cycles. <i>Inorganic Chemistry</i> , 2019 , 58, 2330-2335	5.1	7
56	Coherent manipulation and quantum phase interference in a fullerene-based electron triplet molecular qubit. <i>Npj Quantum Information</i> , 2021 , 7,	8.6	7
55	Assembly of two-fold interpenetrated silver supramolecular coordination polymer using Keggin phosphotungstate template. <i>Inorganic Chemistry Communication</i> , 2014 , 44, 91-95	3.1	6
54	Synthesis, Crystal Structures, and Magnetic Properties of Heterodimetallic RuIIIBd Coordination Compounds Based on a Meridional Tricyanoruthenium(III) Building Block. <i>European Journal of Inorganic Chemistry</i> , 2015 , 2015, 1065-1073	2.3	6
53	Synthesis, crystal structures, and magnetic properties of two three-dimensional octacyanotungstate(IV)-based bimetallic frameworks with 4,4'-bipyridine dioxide (4,4'-dpdo). <i>Science in China Series B: Chemistry</i> , 2009 , 52, 266-275		6
52	Orbital-dependent magnetic properties of molecular cluster containing high-spin Co(II) ions. <i>International Journal of Quantum Chemistry</i> , 2009 , 109, 3368-3378	2.1	6
51	Hydrogen bonding-directed supramolecular structural motifs based on 4,4'-bipyridine dioxide. <i>CrystEngComm</i> , 2001 , 3, 147-151	3.3	6
50	The differential magnetic relaxation behaviours of slightly distorted triangular dodecahedral dysprosium analogues in a type of cyano-bridged 3d-4f zig-zag chain compounds. <i>Dalton Transactions</i> , 2020 , 49, 6867-6875	4.3	6
49	Rare earth permanent magnetic nanostructures: chemical design and microstructure control to optimize magnetic properties. <i>Inorganic Chemistry Frontiers</i> , 2021 , 8, 383-395	6.8	6
48	Electric and magnetic transitions with 90° turning of polarizations in a layered perovskite of [NH4Cl]2[Ni(HCOO)2(NH3)2]. <i>APL Materials</i> , 2018 , 6, 114205	5.7	6

47	Coexistence of magnetic and electric orderings in a divalent Cr-based multiaxial molecular ferroelectric. <i>Chemical Science</i> , 2021 , 12, 9742-9747	9.4	6
46	Two half-sandwich organometallic single-ion magnets with toluene coordinated to the Dy(III) ion: The [(C ₇ H ₈)Dy(AlCl ₄) ₃] and [(C ₇ H ₈)Dy(AlBr ₄) ₃] complexes. <i>Inorganic Chemistry Communication</i> , 2017 , 86, 312-314	3.1	5
45	Structural complexity in the rare earth metallocene hydride complexes, [(CMe) ₂ InH] <i>Dalton Transactions</i> , 2014 , 43, 15526-31	4.3	5
44	Bitriazole bridged manganese(II) framework with weak ferromagnet behavior. <i>Polyhedron</i> , 2011 , 30, 3112-3115	2.7	5
43	Syntheses, structures and magnetic properties of a family of one-dimensional M(II)-lanthanide(III) (M = Ni(II) and Zn(II)) coordination polymers. <i>Science in China Series B: Chemistry</i> , 2007 , 50, 308-317		5
42	Magnetic Nanostructures: Rational Design and Fabrication Strategies toward Diverse Applications.. <i>Chemical Reviews</i> , 2022 ,	68.1	5
41	Triple Helical Molecular Braid and Parallel Packed Wavy Chain-Based Supramolecular Organic Frameworks with Conformation- and Packing-Dependent Luminescent Properties. <i>Crystal Growth and Design</i> , 2016 , 16, 4727-4735	3.5	5
40	Design principle of half-sandwich type erbium single-ion magnets through crystal field engineering: a combined magnetic and electronic structure study. <i>Dalton Transactions</i> , 2019 , 48, 10407-10411	4.3	4
39	Amination of the Gd@C endohedral fullerene: tunable substitution effect on quantum coherence behaviors. <i>Chemical Science</i> , 2020 , 11, 10737-10743	9.4	4
38	Crystal Structures and Magnetic Properties of 2D Supramolecular Architectures Assembled from Benzimidazolecarboxylato-Bridged 1D Double-Stranded Coordinating Chains Featuring Metallomacrocycles as Subunits. <i>European Journal of Inorganic Chemistry</i> , 2008 , 2008, 3776-3785	2.3	4
37	Salts of ammonium-crown ether supramolecular cation with bis(maleonitriledithiolato)copperate(II): Crystal structures, magnetic susceptibilities, single crystal EPR spectra and DFT calculation. <i>Inorganica Chimica Acta</i> , 2005 , 358, 3841-3848	2.7	4
36	Ferromagnetic interactions in EO-azido-bridged binuclear transition metal(II) systems: Syntheses, crystal structures and magnetostructural correlations. <i>Science China Chemistry</i> , 2012 , 55, 942-950	7.9	3
35	Unusual magnetic transition in a new 1D molecular solid [1-(4?-iodobenzyl)pyridinium][Ni(mnt) ₂](⁰). <i>Solid State Sciences</i> , 2006 , 8, 520-525	3.4	3
34	A cyano-bridged dinuclear 4f-3d array. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2002 , 58, m325-6		3
33	Free-standing 2D non-van der Waals antiferromagnetic hexagonal FeSe semiconductor: halide-assisted chemical synthesis and Fe related magnetic transitions.. <i>Chemical Science</i> , 2021 , 13, 203-209	9.4	3
32	Recent progress in metal-based molecular probes for optical bioimaging and biosensing. <i>Current Opinion in Chemical Biology</i> , 2021 , 66, 102097	9.7	3
31	Effect of the Transition Metal Ions on the Single-Molecule Magnet Properties in a Family of Air-Stable 3d-4f Ion-Pair Compounds with Pentagonal Bipyramidal Ln(III) Ions. <i>Inorganic Chemistry</i> , 2021 ,	5.1	3
30	High temperature Fe(III) spin crossover behaviours in three unprecedented FeIII ^{II} BeIII (M = Fe, Cd) linear trinuclear complexes. <i>Inorganic Chemistry Frontiers</i> , 2020 , 7, 1526-1531	6.8	3

29	Orientation mapping of Rabi frequencies in a rare-earth molecular qudit. <i>Inorganic Chemistry Frontiers</i> , 2020 , 7, 3875-3881	6.8	3
28	The comparative studies on the magnetic relaxation behaviour of the axially-elongated pentagonal-bipyramidal dysprosium and erbium ions in similar one-dimensional chain structures. <i>Dalton Transactions</i> , 2021 , 50, 8736-8745	4.3	3
27	Multiple magnetic relaxation pathways in T-shaped N-heterocyclic carbene-supported Fe(I) single-ion magnets. <i>Inorganic Chemistry Frontiers</i> , 2019 , 6, 1050-1057	6.8	2
26	Experimental Determination of Magnetic Anisotropy in Exchange-Bias Dysprosium Metallocene Single-Molecule Magnets. <i>Angewandte Chemie</i> , 2020 , 132, 13137-13143	3.6	2
25	Direct observation of a fast single-crystal-to-single-crystal transformation from a CuII-framework to a CuI-chain mediated by ascorbic acid. <i>CrystEngComm</i> , 2016 , 18, 1878-1882	3.3	2
24	Hydrothermal synthesis, crystal structure and magnetic properties of a new 2D layered vanadium oxide complex: [Ni(phen)(H ₂ O)] ₂ V ₄ O ₁₂ . <i>Journal of Coordination Chemistry</i> , 2005 , 58, 327-334	1.6	2
23	Synthesis, crystal structure and magnetic properties of a nickel(III) maleonitriledithiolate complex using 1-methyl-3-benzylbenzimidazolium cation. <i>Journal of Molecular Structure</i> , 2005 , 743, 197-200	3.4	2
22	Paramagnetic properties adjustment for Gd@C(9)-C82 by regioselective multi-amination. <i>Carbon</i> , 2020 , 158, 320-326	10.4	2
21	Slow magnetic relaxation in structurally similar mononuclear 8-coordinate Fe(II) and Fe(III) compounds. <i>Chemical Communications</i> , 2021 , 57, 781-784	5.8	2
20	Enhancing the magnetic performance of pyrazine-N-oxide bridged dysprosium chains through controlled variation of ligand coordination modes. <i>Dalton Transactions</i> , 2021 , 50, 7048-7055	4.3	2
19	Neutral polar hybrid perovskites of [(CH ₃) ₂ SO][RE(HCOO) ₃] (RE = Lu and Y): Phase transitions driven by transformation of weak C-H...O interactions. <i>APL Materials</i> , 2021 , 9, 021112	5.7	2
18	High temperature anionic Fe(III) spin crossover behavior in a mixed-valence Fe(II)/Fe(III) complex. <i>Dalton Transactions</i> , 2021 , 50, 5960-5967	4.3	2
17	The construction of dynamic dysprosium-carboxylate ribbons by utilizing the hybrid-ligand conception. <i>Dalton Transactions</i> , 2021 , 50, 1246-1252	4.3	2
16	A neutral auxiliary ligand enhanced dysprosium(III) single molecule magnet. <i>Dalton Transactions</i> , 2018 , 47, 7395-7398	4.3	1
15	Metal-Organic Frameworks: Functional Magnetic Materials with Formate 2014 , 1-23		1
14	Synthesis and crystal structure of 1D Cd-amine coordination polymer and its luminescent properties. <i>Chemical Research in Chinese Universities</i> , 2014 , 30, 720-725	2.2	1
13	New tricyanoiron(III) building blocks for the construction of molecule-based magnets. <i>Science China Chemistry</i> , 2010 , 53, 2106-2111	7.9	1
12	Single-Crystal Dendritic Micro-Pines of Magnetic Fe ₂ O ₃ : Large-Scale Synthesis, Formation Mechanism, and Properties.. <i>ChemInform</i> , 2005 , 36, no		1

- 11 Gallium (III) Complexes in Cancer Chemotherapy. *European Journal of Inorganic Chemistry*, 2.3 1
- 10 Slow magnetic relaxation in high-coordinate Co(II) and Fe(II) compounds bearing neutral tetradentate ligands. *Dalton Transactions*, **2021**, 50, 15327-15335 4.3 1
- 9 First-order-reversal-curve analysis of rare earth permanent magnet nanostructures: insight into the coercivity enhancement mechanism through regulating the Nd-rich phase. *Inorganic Chemistry Frontiers*, **2021**, 8, 1975-1982 6.8 1
- 8 A series of counter cation-dependent tetra-β-diketonate mononuclear lanthanide(III) single-molecule magnets and immobilization on pre-functionalised GaN substrates by anion exchange reaction. *Journal of Materials Chemistry C*, 7.1 1
- 7 Single-Ion Anisotropy: An Insight to Complicated Magnetic Molecules. *Topics in Organometallic Chemistry*, **2018**, 227-252 0.6 1
- 6 Tuning the dynamic magnetic behaviour and proton conductivity via water-induced reversible single-crystal to single-crystal structural transformation. *Journal of Materials Chemistry C*, **2021**, 9, 15858-15867 7.1 1
- 5 A Unique Layered Cu-formate Hydrate of Cu(HCOO)₂·1/3H₂O: Structures, Dehydration, and Thermal and Magnetic Properties. *Chemical Research in Chinese Universities*, **2022**, 38, 107-116 2.2
- 4 Constructing Magnetic Molecular Solids by Employing Three-Atom Ligands as Bridges 195-228
- 3 Hybrid Dicyanamide Perovskites **2020**, 181-197
- 2 Hybrid Azide Perovskites **2020**, 151-179
- 1 Hybrid Formate Perovskites **2020**, 79-149