Takashi Fujimura

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1169288/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Increase of regulatory T cells and the ratio of specific IgE to total IgE are candidates for response monitoring or prognostic biomarkers in 2-year sublingual immunotherapy (SLIT) for Japanese cedar pollinosis. Clinical Immunology, 2011, 139, 65-74.	1.4	80
2	A Randomized Controlled Trial of Sublingual Immunotherapy for Japanese Cedar Pollinosis. International Archives of Allergy and Immunology, 2008, 146, 76-84.	0.9	65
3	Isolation and characterization of native Cry j 3 from Japanese cedar (Cryptomeria japonica) pollen. Allergy: European Journal of Allergy and Clinical Immunology, 2007, 62, 547-553.	2.7	44
4	The Induced Regulatory T Cell Level, Defined as the Proportion of IL-10 ⁺ Foxp3 ⁺ Cells among CD25 ⁺ CD4 ⁺ Leukocytes, Is a Potential Therapeutic Biomarker for Sublingual Immunotherapy: A Preliminary Report. International Archives of Allergy and Immunology, 2010, 153, 378-387.	0.9	43
5	Anticancer effect of nor-wogonin (5, 7, 8-trihydroxyflavone) on human triple-negative breast cancer cells via downregulation of TAK1, NF-κB, and STAT3. Pharmacological Reports, 2019, 71, 289-298.	1.5	34
6	Two-Dimensional IgE-Binding Spectrum of Japanese Cedar <i>(Cryptomeria japonica)</i> Pollen Allergens. International Archives of Allergy and Immunology, 2004, 133, 125-135.	0.9	33
7	Spectrum of allergens for Japanese cedar pollinosis and impact of component-resolved diagnosis on allergen-specific immunotherapy. Allergology International, 2015, 64, 312-320.	1.4	33
8	A flavanone derivative from the Asian medicinal herb (Perilla frutescens) potently suppresses IgE-mediated immediate hypersensitivity reactions. Biochemical and Biophysical Research Communications, 2017, 483, 674-679.	1.0	31
9	Influences of Maternal Factors Over Offspring Allergies and the Application for Food Allergy. Frontiers in Immunology, 2019, 10, 1933.	2.2	31
10	Molecular cloning and characterization of a new Japanese cedar pollen allergen homologous to plant isoflavone reductase family. Clinical and Experimental Allergy, 2002, 32, 1064-1070.	1.4	28
11	Toward elucidating the full spectrum of mite allergens — state of the art. Journal of Bioscience and Bioengineering, 2002, 94, 285-298.	1.1	26
12	Molecular cloning of a class IV chitinase allergen from Japanese cedar (Cryptomeria japonica) pollen and competitive inhibition of its immunoglobulin E-binding capacity by latex C-serum. Clinical and Experimental Allergy, 2005, 35, 234-243.	1.4	26
13	A novel moonlight function of glyceraldehydeâ€3â€phosphate dehydrogenase (GAPDH) for immunomodulation. BioFactors, 2018, 44, 597-608.	2.6	22
14	Der f 35: An MD-2-like house dust mite allergen that cross-reacts with Der f 2 and Pso o 2. Allergy: European Journal of Allergy and Clinical Immunology, 2017, 72, 1728-1736.	2.7	21
15	Toward elucidating the full spectrum of mite allergensstate of the art. Journal of Bioscience and Bioengineering, 2002, 94, 285-98.	1.1	20
16	Antigen-Specific Immunotherapy against Allergic Rhinitis: The State of the Art. Allergology International, 2010, 59, 21-31.	1.4	18
17	Der f 34, a Novel Major House Dust Mite Allergen Belonging to a Highly Conserved Rid/YjgF/YER057c/UK114 Family of Imine Deaminases. Journal of Biological Chemistry, 2016, 291, 21607-21615.	1.6	17
18	Induction of Th1 immune responses to Japanese cedar pollen allergen (Cry j 1) in mice immunized with Cry j 1 conjugated with CpG oligodeoxynucleotide. Comparative Immunology, Microbiology and Infectious Diseases, 2011, 34, 157-161.	0.7	13

Takashi Fujimura

#	Article	IF	CITATIONS
19	lgE reactivity to a Cry j 3, an allergen of Japanese cedar (Cryptomeria japonica) pollen in dogs with canine atopic dermatitis. Veterinary Immunology and Immunopathology, 2012, 149, 132-135.	0.5	13
20	Recombinant Fusion Allergens, Cry j 1 and Cry j 2 from Japanese Cedar Pollen, Conjugated with Polyethylene Glycol Potentiate the Attenuation of Cry j 1-Specific IgE Production in Cry j 1-Sensitized Mice and Japanese Cedar Pollen Allergen-Sensitized Monkeys. International Archives of Allergy and Immunology, 2015, 168, 32-43.	0.9	13
21	Design, Synthesis and Cytotoxicity Evaluation of New 3, 5-Disubstituted-2-Thioxoimidazolidinones. Anti-Cancer Agents in Medicinal Chemistry, 2018, 18, 573-582.	0.9	13
22	Approaches to immunotherapies for Japanese cedar pollinosis. Auris Nasus Larynx, 2011, 38, 431-438.	0.5	11
23	Synergistic tumor suppression by a Perilla frutescens-derived methoxyflavanone and anti-cancer tyrosine kinase inhibitors in A549 human lung adenocarcinoma. Cytotechnology, 2018, 70, 913-919.	0.7	11
24	A methoxyflavanone derivative from the Asian medicinal herb (Perilla frutescens) induces p53-mediated G2/M cell cycle arrest and apoptosis in A549 human lung adenocarcinoma. Cytotechnology, 2018, 70, 899-912.	0.7	10
25	Prominent IgE-binding and cytokine-inducing capacities of a newly cloned N-terminal region of Der f 14, an apolipophorin-like house dust mite allergen. Journal of Biochemistry, 2018, 163, 51-60.	0.9	9
26	Impact of Histone H1 on the Progression of Allergic Rhinitis and Its Suppression by Neutralizing Antibody in Mice. PLoS ONE, 2016, 11, e0153630.	1.1	9
27	Anti-inflammatory intravenous immunoglobulin (IVIg) suppresses homeostatic proliferation of B cells. Cytotechnology, 2018, 70, 921-927.	0.7	8
28	Immunological Parameters Associated with the Development of Allergic Rhinitis: A Preliminary Prospective Study. American Journal of Rhinology and Allergy, 2012, 26, 92-96.	1.0	7
29	Intravenous immunoglobulin (IVIg) acts directly on conventional T cells to suppress T cell receptor signaling. Biochemical and Biophysical Research Communications, 2020, 522, 792-798.	1.0	5
30	Therapeutic Effects and Biomarkers in Sublingual Immunotherapy: A Review. Journal of Allergy, 2012, 2012, 1-9.	0.7	4
31	Exposure to positively- and negatively-charged plasma cluster ions impairs IgE-binding capacity of indoor cat and fungal allergens. World Allergy Organization Journal, 2016, 9, 27.	1.6	3
32	Intake of a fermented plant product attenuates allergic symptoms without changing systemic immune responses in a mouse model of Japanese cedar pollinosis. World Allergy Organization Journal, 2018, 11, 31.	1.6	2
33	New treatment strategies for allergic rhinitis utilizing the oral mucosa. Clinical and Experimental Allergy Reviews, 2012, 12, 7-13.	0.3	0
34	Plasma Cluster Ions Reduce the IgE-Binding Capacity of House Dust Mite Allergens under a Simulated Indoor Environmental Condition. International Archives of Allergy and Immunology, 2017, 173, 199-203.	0.9	0
35	Abstract 3040: A novel methoxyflavanone from a Chinese medicinal herb (Perilla frutescens) induces G2/M cell cycle arrest and apoptosis in A549 human lung adenocarcinoma cells. , 2016, , .		0