

Francesca Budillon

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1173596/publications.pdf>

Version: 2024-02-01

26
papers

515
citations

623734

14
h-index

677142

22
g-index

28
all docs

28
docs citations

28
times ranked

851
citing authors

#	ARTICLE	IF	CITATIONS
1	Coralligenous banks along the western and northern continental shelf of Sardinia Island (Mediterranean Sea). <i>Journal of Maps</i> , 2022, 18, 200-209.	2.0	9
2	Large-Scale Mercury Dispersion at Sea: Modelling a Multi-Hazard Case Study from Augusta Bay (Central Mediterranean Sea). <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 3956.	2.6	3
3	Present-day infralittoral prograding wedges (IPWs) in Central-Eastern Tyrrhenian Sea: Critical issues and challenges to their use as geomorphological indicators of sea level. <i>Marine Geology</i> , 2022, 450, 106821.	2.1	9
4	Evaluation of disturbance induced on soft offshore sediments by two types of gravity piston coring techniques. <i>Marine Geology</i> , 2019, 417, 106005.	2.1	6
5	A Quantitative Evaluation of Hyperpycnal Flow Occurrence in a Temperate Coastal Zone: The Example of the Salerno Gulf (Southern Italy). <i>Geosciences (Switzerland)</i> , 2019, 9, 501.	2.2	1
6	Open-slope, translational submarine landslide in a tectonically active volcanic continental margin (Licosa submarine landslide, southern Tyrrhenian Sea). <i>Geological Society Special Publication</i> , 2019, 477, 133-150.	1.3	9
7	Seabed mapping in the Pelagie Islands marine protected area (Sicily Channel, southern Mediterranean) using Remote Sensing Object Based Image Analysis (RSOBI). <i>Marine Geophysical Researches</i> , 2019, 40, 333-355.	1.2	39
8	A critical review of potential tsunamigenic sources as first step towards the tsunami hazard assessment for the Napoli Gulf (Southern Italy) highly populated area. <i>Natural Hazards</i> , 2018, 92, 43-76.	3.4	13
9	Coastal Evolution, Hydrothermal Migration Pathways and Soft Deformation along the Campania Continental Shelf (Southern Tyrrhenian Sea): Insights from High-Resolution Seismic Profiles. <i>Geosciences (Switzerland)</i> , 2018, 8, 121.	2.2	6
10	Pockmark morphology and turbulent buoyant plumes at a submarine spring. <i>Continental Shelf Research</i> , 2017, 148, 19-36.	1.8	7
11	Distribution of <i>Posidonia oceanica</i> (L.) Delile meadows around Lampedusa Island (Strait of Tj ETQq1 1 0.784314 rgBT /Overlock 2.0 17	2.0	17
12	A newly discovered Pliocene volcanic field on the western Sardinia continental margin (western Tj ETQq0 0 0 rgBT /Overlock 1.1 14 Tf 50 30	1.1	14
13	Mediterranean coccolith ecobiostratigraphy since the penultimate Glacial (the last 145,000 years) and ecobioevent traceability. <i>Marine Micropaleontology</i> , 2015, 115, 24-38.	1.2	17
14	Sorted bedforms over transgressive deposits along the continental shelf of western Sardinia (Mediterranean Sea). <i>Marine Geology</i> , 2015, 359, 75-88.	2.1	27
15	Combined palaeomagnetic secular variation and petrophysical records to time-constrain geological and hazardous events: An example from the eastern Tyrrhenian Sea over the last 120ka. <i>Global and Planetary Change</i> , 2014, 113, 91-109.	3.5	34
16	Seismic facies and seabed morphology in a tectonically controlled continental shelf: The Augusta Bay (offshore eastern Sicily, Ionian Sea). <i>Marine Geology</i> , 2013, 335, 35-51.	2.1	13
17	Late Holocene sedimentary changes in shallow water settings: the case of the Sele river offshore in the Salerno Gulf (south-eastern Tyrrhenian Sea, Italy). <i>Rendiconti Lincei</i> , 2012, 23, 25-43.	2.2	17
18	Evolution of the Sele River coastal plain (southern Italy) during the Late Quaternary by inland and offshore stratigraphical analyses. <i>Rendiconti Lincei</i> , 2012, 23, 81-102.	2.2	24

#	ARTICLE	IF	CITATIONS
19	The key role played by the Augusta basin (southern Italy) in the mercury contamination of the Mediterranean Sea. <i>Journal of Environmental Monitoring</i> , 2011, 13, 1753.	2.1	59
20	The Bulgheria canyon-fan: a small-scale proximal system in the eastern Tyrrhenian Sea (Italy). <i>Marine Geophysical Researches</i> , 2011, 32, 83-97.	1.2	14
21	Integrated stratigraphic reconstruction for the last 80kyr in a deep sector of the Sardinia Channel (Western Mediterranean). <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2009, 56, 725-737.	1.4	18
22	Sediment transport and deposition during extreme sea storm events at the Salerno Bay (Tyrrhenian) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 Sciences</i> , 2006, 6, 839-852.	3.6	23
23	Event beds in the recent prodelta stratigraphic record of the small flood-prone Bonea Stream (Amalfi) <i>Tj ETQq1 1 0,784314 rgBT /Overlock 10 Tf 5</i>	2.1	96
24	Evidence for a variable paleomagnetic lock-in depth in the Holocene sequence from the Salerno Gulf (Italy): Implications for "high-resolution" paleomagnetic dating. <i>Geochemistry, Geophysics, Geosystems</i> , 2005, 6, n/a-n/a.	2.5	58
25	High-resolution petrophysical and palaeomagnetic study of late-Holocene shelf sediments, Salerno Gulf, Tyrrhenian Sea. <i>Holocene</i> , 2004, 14, 426-435.	1.7	23
26	The geological record of storm events over the last 1000 years in the Salerno Bay (Southern) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 462 17</i>	12.0	17