

Carla Calagua

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11744468/publications.pdf>

Version: 2024-02-01

14
papers

475
citations

1040056

9
h-index

1058476

14
g-index

16
all docs

16
docs citations

16
times ranked

1088
citing authors

#	ARTICLE	IF	CITATIONS
1	EZH2 inhibition activates a dsRNAâ€“STINGâ€“interferon stress axis that potentiates response to PD-1 checkpoint blockade in prostate cancer. <i>Nature Cancer</i> , 2021, 2, 444-456.	13.2	118
2	Evaluation of Intense Androgen Deprivation Before Prostatectomy: A Randomized Phase II Trial of Enzalutamide and Leuprolide With or Without Abiraterone. <i>Journal of Clinical Oncology</i> , 2019, 37, 923-931.	1.6	78
3	Expression of PD-L1 in Hormone-naïve and Treated Prostate Cancer Patients Receiving Neoadjuvant Abiraterone Acetate plus Prednisone and Leuprolide. <i>Clinical Cancer Research</i> , 2017, 23, 6812-6822.	7.0	77
4	Neoadjuvant-Intensive Androgen Deprivation Therapy Selects for Prostate Tumor Foci with Diverse Subclonal Oncogenic Alterations. <i>Cancer Research</i> , 2018, 78, 4716-4730.	0.9	56
5	Downregulation of <i>Dipeptidyl Peptidase 4</i> Accelerates Progression to Castration-Resistant Prostate Cancer. <i>Cancer Research</i> , 2018, 78, 6354-6362.	0.9	42
6	A Subset of Localized Prostate Cancer Displays an Immunogenic Phenotype Associated with Losses of Key Tumor Suppressor Genes. <i>Clinical Cancer Research</i> , 2021, 27, 4836-4847.	7.0	20
7	Phosphorylation of androgen receptor serine 81 is associated with its reactivation in castration-resistant prostate cancer. <i>Cancer Letters</i> , 2018, 438, 97-104.	7.2	19
8	Phosphorylation of the androgen receptor at Ser81 is coâ€“sustained by CDK1 and CDK9 and leads to ARâ€“mediated transactivation in prostate cancer. <i>Molecular Oncology</i> , 2021, 15, 1901-1920.	4.6	19
9	Autocrine Canonical Wnt Signaling Primes Noncanonical Signaling through ROR1 in Metastatic Castration-Resistant Prostate Cancer. <i>Cancer Research</i> , 2022, 82, 1518-1533.	0.9	15
10	Prostate cancer susceptibility gene <i>HIST1H1A</i> is a modulator of androgen receptor signaling and epithelial to mesenchymal transition. <i>Oncotarget</i> , 2018, 9, 28532-28546.	1.8	7
11	Primary adenocarcinoma of the bladder lacks mismatch repair deficiency and demonstrates PD-L1 expression in tumor-infiltrating immune cells, with implications in both diagnosis and therapeutics. <i>Human Pathology</i> , 2019, 94, 58-63.	2.0	6
12	Metastatic Castration-Resistant Prostate Cancer Remains Dependent on Oncogenic Drivers Found in Primary Tumors. <i>JCO Precision Oncology</i> , 2021, 5, 1514-1522.	3.0	6
13	Collision tumors revealed by prospectively assessing subtype-defining molecular alterations in 904 individual prostate cancer foci. <i>JCI Insight</i> , 2022, 7, .	5.0	6
14	A phase II study of nivolumab in patients with high-risk biochemically recurrent (BCR) prostate cancer (PCa).. <i>Journal of Clinical Oncology</i> , 2019, 37, TPS341-TPS341.	1.6	1