

Mona Badawy

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11757464/publications.pdf>

Version: 2024-02-01

9
papers

296
citations

1307594

7
h-index

1474206

9
g-index

9
all docs

9
docs citations

9
times ranked

346
citing authors

#	ARTICLE	IF	CITATIONS
1	Antibiotic-Loaded Bone Cement in Prevention of Periprosthetic Joint Infections in Primary Total Knee Arthroplasty: A Register-based Multicentre Randomised Controlled Non-inferiority Trial (ALBA trial). <i>BMJ Open</i> , 2021, 11, e041096.	1.9	15
2	Development of an internet-delivered cognitive behavioral therapy program for use in combination with exercise therapy and education by patients at increased risk of chronic pain following total knee arthroplasty. <i>BMC Health Services Research</i> , 2021, 21, 1151.	2.2	9
3	The effectiveness of exercise therapy and education plus cognitive behavioral therapy, alone or in combination with total knee arthroplasty in patients with knee osteoarthritis – study protocol for the MultiKnee trial. <i>BMC Musculoskeletal Disorders</i> , 2021, 22, 1054.	1.9	7
4	Primary constrained and hinged total knee arthroplasty: 2- and 5-year revision risk compared with unconstrained total knee arthroplasty: a report on 401 cases from the Norwegian Arthroplasty Register 1994–2017. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2019, 90, 467-472.	3.3	22
5	Have the Causes of Revision for Total and Unicompartmental Knee Arthroplasties Changed During the Past Two Decades?. <i>Clinical Orthopaedics and Related Research</i> , 2017, 475, 1874-1886.	1.5	98
6	Hospital volume and the risk of revision in Oxford unicompartmental knee arthroplasty in the Nordic countries -an observational study of 14,496 cases. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 388.	1.9	35
7	Patient and surgical factors affecting procedure duration and revision risk due to deep infection in primary total knee arthroplasty. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 544.	1.9	35
8	A randomized study on migration of the Spectron EF and the Charnley flanged 40 cemented femoral components using radiostereometric analysis at 2 years. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2011, 82, 538-544.	3.3	21
9	Wear and migration of highly cross-linked and conventional cemented polyethylene cups with cobalt chrome or Oxinium femoral heads: A randomized radiostereometric study of 150 patients. <i>Journal of Orthopaedic Research</i> , 2011, 29, 1222-1229.	2.3	54