

List of Publications by Year in  
Descending Order

**Source:** <https://exaly.com/author-pdf/11774447/p-siddhuraju-publications-by-year.pdf>  
**Version:** 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19 papers	1,643 citations	16 h-index	19 g-index
19 ext. papers	1,757 ext. citations	5.2 avg, IF	4.51 L-index

#	Paper	IF	Citations
19	Antioxidant and free radical scavenging capacity of the underutilized legume, <i>Vigna vexillata</i> (L.) A. Rich. <i>Journal of Food Composition and Analysis</i> , <b>2011</b> , 24, 160-165	4.1	28
18	A comparative study on in vitro antioxidant activity of the legumes <i>Acacia auriculiformis</i> and <i>Acacia ferruginea</i> with a conventional legume <i>Cajanus cajan</i> Estudio comparativo de la actividad antioxidante in vitro de las legumbres <i>Acacia auriculiformis</i> y <i>Acacia ferruginea</i> con la legumbre convencional <i>Cajanus cajan</i> . <i>CYTA - Journal of Food</i> , <b>2011</b> , 9, 8-16	2.3	8
17	The antioxidant activity and free radical scavenging potential of two different solvent extracts of <i>Camellia sinensis</i> (L.) O. Kuntz, <i>Ficus bengalensis</i> L. and <i>Ficus racemosa</i> L.. <i>Food Chemistry</i> , <b>2008</b> , 107, 1000-1007	8.5	190
16	The antioxidant activity and free radical-scavenging capacity of dietary phenolic extracts from horse gram ( <i>Macrotyloma uniflorum</i> (Lam.) Verdc.) seeds. <i>Food Chemistry</i> , <b>2007</b> , 105, 950-958	8.5	143
15	Antioxidant activity of polyphenolic compounds extracted from defatted raw and dry heated <i>Tamarindus indica</i> seed coat. <i>LWT - Food Science and Technology</i> , <b>2007</b> , 40, 982-990	5.4	85
14	L-DOPA (L-3,4-Dihydroxyphenylalanine). <i>Methods in Molecular Biology</i> , <b>2007</b> , 51-53	1.4	
13	Chemical composition, toxic/antimetabolic constituents, and effects of different treatments on their levels, in four provenances of L. from Mexico. <i>Food Chemistry</i> , <b>2006</b> , 96, 80-89	8.5	274
12	Alternative food/feed perspectives of an underutilized legume <i>Mucuna pruriens</i> var. utilis--a review. <i>Plant Foods for Human Nutrition</i> , <b>2005</b> , 60, 201-18	3.9	69
11	Comparative nutritional evaluation of differentially processed mucuna seeds [ <i>Mucuna pruriens</i> (L.) DC. var. utilis (Wall ex Wight) Baker ex Burck] on growth performance, feed utilization and body composition in Nile tilapia ( <i>Oreochromis niloticus</i> L.). <i>Aquaculture Research</i> , <b>2003</b> , 34, 487-500	1.9	36
10	Comparative nutritional evaluation of raw, methanol extracted residues and methanol extracts of moringa ( <i>Moringa oleifera</i> Lam.) leaves on growth performance and feed utilization in Nile tilapia ( <i>Oreochromis niloticus</i> L.). <i>Aquaculture Research</i> , <b>2003</b> , 34, 1147-1159	1.9	51
9	Effect of phenolic nonprotein amino acid L-dopa (L-3,4-dihydroxyphenylalanine) on growth performance, metabolic rates and feed nutrient utilization of common carp ( <i>Cyprinus carpio</i> L.). <i>Aquaculture Nutrition</i> , <b>2002</b> , 8, 69-77	3.2	15
8	The effect of ionising radiation on antinutritional factors and the nutritional value of plant materials with reference to human and animal food. <i>Food Chemistry</i> , <b>2002</b> , 78, 187-205	8.5	99
7	Effect of soaking and ionising radiation on various antinutritional factors of seeds from different species of an unconventional legume, <i>Sesbania</i> and a common legume, green gram ( <i>Vigna radiata</i> ). <i>Food Chemistry</i> , <b>2002</b> , 79, 273-281	8.5	47
6	Studies on the antioxidant activity of Indian Laburnum ( <i>Cassia fistula</i> L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. <i>Food Chemistry</i> , <b>2002</b> , 79, 61-67	8.5	311
5	Effect of various domestic processing methods on antinutrients and in vitro protein and starch digestibility of two indigenous varieties of Indian tribal pulse, <i>Mucuna pruriens</i> Var. utilis. <i>Journal of Agricultural and Food Chemistry</i> , <b>2001</b> , 49, 3058-67	5.7	78
4	Studies on the nutritional composition and antinutritional factors of three different germplasm seed materials of an under-utilized tropical legume, <i>Mucuna pruriens</i> var. utilis. <i>Journal of Agricultural and Food Chemistry</i> , <b>2000</b> , 48, 6048-60	5.7	87
3	Effect of soaking and heat processing on the levels of antinutrients and digestible proteins in seeds of <i>Vigna aconitifolia</i> and <i>Vigna sinensis</i> . <i>Food Chemistry</i> , <b>1998</b> , 63, 259-264	8.5	57

2	Effect of different post-harvest treatments on antinutritional factors in seeds of the tribal pulse, <i>Mucuna pruriens</i> (L.) DC. <i>International Journal of Food Sciences and Nutrition</i> , <b>1996</b> , 47, 263-72	3.7	30
1	Effects of various water or hydrothermal treatments on certain antinutritional compounds in the seeds of the tribal pulse, <i>Dolichos lablab</i> var. <i>vulgaris</i> L. <i>Plant Foods for Human Nutrition</i> , <b>1995</b> , 48, 17-29 <sup>3.9</sup>		35