
Oya YÃ-nal

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11776627/publications.pdf Version: 2024-02-01



Ονα ΥΔ-ΝΑΙ

#	Article	IF	CITATIONS
1	Serum levels of omentin, chemerin and adipsin in patients with biopsy-proven nonalcoholic fatty liver disease. Scandinavian Journal of Gastroenterology, 2011, 46, 91-97.	1.5	107
2	Circulating vaspin levels and epicardial adipose tissue thickness are associated with impaired coronary flow reserve in patients with nonalcoholic fatty liver disease. Atherosclerosis, 2011, 217, 125-129.	0.8	53
3	Serum levels of vaspin, obestatin, and apelin-36 in patients with nonalcoholic fatty liver disease. Metabolism: Clinical and Experimental, 2011, 60, 544-549.	3.4	61
4	Serum Progranulin as an Independent Marker of Liver Fibrosis in Patients with Biopsy-Proven Nonalcoholic Fatty Liver Disease. Disease Markers, 2011, 31, 205-210.	1.3	24
5	Decreased Prohepcidin Levels in Patients with HBV-Related Liver Disease: Relation with Ferritin Levels. Digestive Diseases and Sciences, 2010, 55, 3548-3551.	2.3	11
6	Serum levels of osteoprotegerin in the spectrum of nonalcoholic fatty liver disease. Scandinavian Journal of Clinical and Laboratory Investigation, 2010, 70, 541-546.	1.2	38
7	Coronary flow reserve is impaired in patients with nonalcoholic fatty liver disease: Association with liver fibrosis. Atherosclerosis, 2010, 211, 182-186.	0.8	84
8	What is the most accurate method for the diagnosis of intestinal tuberculosis?. Turkish Journal of Gastroenterology, 2010, 21, 91-96.	1.1	20
9	How should we follow the Crohn's disease patients in remission?. Turkish Journal of Gastroenterology, 2010, 21, 97-99.	1.1	Ο
10	HFE Gene Mutation, Chronic Liver Disease, and Iron Overload In Turkey. Digestive Diseases and Sciences, 2007, 52, 3298-3302.	2.3	14