

# Lindsay C Hewitt

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/117864/publications.pdf>

Version: 2024-02-01

10  
papers

112  
citations

1306789

7  
h-index

1473754

9  
g-index

10  
all docs

10  
docs citations

10  
times ranked

336  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prognostic value of pathological lymph node status and primary tumour regression grading following neoadjuvant chemotherapy – results from the MRC OE02 oesophageal cancer trial. <i>Histopathology</i> , 2018, 72, 1180-1188.	1.6	31
2	KRAS status is related to histological phenotype in gastric cancer: results from a large multicentre study. <i>Gastric Cancer</i> , 2019, 22, 1193-1203.	2.7	16
3	Frequent Coamplification of Receptor Tyrosine Kinase and Downstream Signaling Genes in Japanese Primary Gastric Cancer and Conversion in Matched Lymph Node Metastasis. <i>Annals of Surgery</i> , 2018, 267, 114-121.	2.1	15
4	Computational Image Analysis of T-Cell Infiltrates in Resectable Gastric Cancer: Association with Survival and Molecular Subtypes. <i>Journal of the National Cancer Institute</i> , 2021, 113, 88-98.	3.0	15
5	Biopsy proportion of tumour predicts pathological tumour response and benefit from chemotherapy in resectable oesophageal carcinoma: results from the UK MRC OE02 trial. <i>Oncotarget</i> , 2016, 7, 77565-77575.	0.8	12
6	Reduced genomic tumor heterogeneity after neoadjuvant chemotherapy is related to favorable outcome in patients with esophageal adenocarcinoma. <i>Oncotarget</i> , 2016, 7, 44084-44095.	0.8	10
7	Technical Reproducibility of Single-Nucleotide and Size-Based DNA Biomarker Assessment Using DNA Extracted from Formalin-Fixed, Paraffin-Embedded Tissues. <i>Journal of Molecular Diagnostics</i> , 2015, 17, 242-250.	1.2	8
8	Increasing frequency of gene copy number aberrations is associated with immunosuppression and predicts poor prognosis in gastric adenocarcinoma. <i>British Journal of Surgery</i> , 2022, 109, 291-297.	0.1	4
9	Comprehensive biomarker analyses identifies HER2, EGFR, MET RNA expression and thymidylate synthase 5'UTR SNP as predictors of benefit from S-1 adjuvant chemotherapy in Japanese patients with stage II/III gastric cancer. <i>Journal of Cancer</i> , 2019, 10, 5130-5138.	1.2	1
10	Author response to: Increasing frequency of gene copy number aberrations is associated with immunosuppression and predicts poor prognosis in gastric adenocarcinoma. <i>British Journal of Surgery</i> , 2022, , .	0.1	0