

Lindsay C Hewitt

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/117864/publications.pdf>

Version: 2024-02-01

10
papers

112
citations

1306789

7
h-index

1473754

9
g-index

10
all docs

10
docs citations

10
times ranked

336
citing authors

#	ARTICLE	IF	CITATIONS
1	Increasing frequency of gene copy number aberrations is associated with immunosuppression and predicts poor prognosis in gastric adenocarcinoma. <i>British Journal of Surgery</i> , 2022, 109, 291-297.	0.1	4
2	Author response to: Increasing frequency of gene copy number aberrations is associated with immunosuppression and predicts poor prognosis in gastric adenocarcinoma. <i>British Journal of Surgery</i> , 2022, , .	0.1	0
3	Computational Image Analysis of T-Cell Infiltrates in Resectable Gastric Cancer: Association with Survival and Molecular Subtypes. <i>Journal of the National Cancer Institute</i> , 2021, 113, 88-98.	3.0	15
4	KRAS status is related to histological phenotype in gastric cancer: results from a large multicentre study. <i>Gastric Cancer</i> , 2019, 22, 1193-1203.	2.7	16
5	Comprehensive biomarker analyses identifies <i>HER2</i> , <i>EGFR</i> , <i>MET</i> RNA expression and thymidylate synthase 5'UTR SNP as predictors of benefit from S-1 adjuvant chemotherapy in Japanese patients with stage II/III gastric cancer. <i>Journal of Cancer</i> , 2019, 10, 5130-5138.	1.2	1
6	Prognostic value of pathological lymph node status and primary tumour regression grading following neoadjuvant chemotherapy – results from the MRC OE02 oesophageal cancer trial. <i>Histopathology</i> , 2018, 72, 1180-1188.	1.6	31
7	Frequent Coamplification of Receptor Tyrosine Kinase and Downstream Signaling Genes in Japanese Primary Gastric Cancer and Conversion in Matched Lymph Node Metastasis. <i>Annals of Surgery</i> , 2018, 267, 114-121.	2.1	15
8	Biopsy proportion of tumour predicts pathological tumour response and benefit from chemotherapy in resectable oesophageal carcinoma: results from the UK MRC OE02 trial. <i>Oncotarget</i> , 2016, 7, 77565-77575.	0.8	12
9	Reduced genomic tumor heterogeneity after neoadjuvant chemotherapy is related to favorable outcome in patients with esophageal adenocarcinoma. <i>Oncotarget</i> , 2016, 7, 44084-44095.	0.8	10
10	Technical Reproducibility of Single-Nucleotide and Size-Based DNA Biomarker Assessment Using DNA Extracted from Formalin-Fixed, Paraffin-Embedded Tissues. <i>Journal of Molecular Diagnostics</i> , 2015, 17, 242-250.	1.2	8