

# Abdullah H Baqui

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/11816462/abdullah-h-baqui-publications-by-citations.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

113  
papers

6,063  
citations

37  
h-index

76  
g-index

115  
ext. papers

6,900  
ext. citations

6.7  
avg, IF

4.9  
L-index

| #   | Paper  | IF   | Citations |
|-----|--|------|-----------|
| 113 | Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: a pooled country analysis. <i>Lancet, The</i> , <b>2013</b> , 382, 417-425  | 4.0  | 452       |
| 112 | Effect of community-based newborn-care intervention package implemented through two service-delivery strategies in Sylhet district, Bangladesh: a cluster-randomised controlled trial. <i>Lancet, The</i> , <b>2008</b> , 371, 1936-44 | 4.0  | 443       |
| 111 | National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. <i>The Lancet Global Health</i> , <b>2013</b> , 1, e26-36                             | 13.6 | 404       |
| 110 | Effect of community-based behaviour change management on neonatal mortality in Shivgarh, Uttar Pradesh, India: a cluster-randomised controlled trial. <i>Lancet, The</i> , <b>2008</b> , 372, 1151-62                                  | 4.0  | 357       |
| 109 | Effect of zinc supplementation started during diarrhoea on morbidity and mortality in Bangladeshi children: community randomised trial. <i>BMJ, The</i> , <b>2002</b> , 325, 1059  | 5.9  | 211       |
| 108 | Iron and zinc supplementation promote motor development and exploratory behavior among Bangladeshi infants. <i>American Journal of Clinical Nutrition</i> , <b>2004</b> , 80, 903-10   | 7    | 189       |
| 107 | Effect of the Integrated Management of Childhood Illness strategy on childhood mortality and nutrition in a rural area in Bangladesh: a cluster randomised trial. <i>Lancet, The</i> , <b>2009</b> , 374, 393-403                      | 4.0  | 178       |
| 106 | Methodological issues in diarrhoeal diseases epidemiology: definition of diarrhoeal episodes. <i>International Journal of Epidemiology</i> , <b>1991</b> , 20, 1057-63   | 7.8  | 161       |
| 105 | Progress and barriers for the control of diarrhoeal disease. <i>Lancet, The</i> , <b>2010</b> , 376, 63-7  | 4.0  | 156       |
| 104 | Rates, timing and causes of neonatal deaths in rural India: implications for neonatal health programmes. <i>Bulletin of the World Health Organization</i> , <b>2006</b> , 84, 706-13   | 8.2  | 146       |
| 103 | Estimates of burden and consequences of infants born small for gestational age in low and middle income countries with INTERGROWTH-21 standard: analysis of CHERG datasets. <i>BMJ, The</i> , <b>2017</b> , 358, j3677                 | 5.9  | 145       |
| 102 | Local understandings of vulnerability and protection during the neonatal period in Sylhet District, Bangladesh: a qualitative study. <i>Lancet, The</i> , <b>2005</b> , 366, 478-85  | 4.0  | 144       |
| 101 | The effect of cord cleansing with chlorhexidine on neonatal mortality in rural Bangladesh: a community-based, cluster-randomised trial. <i>Lancet, The</i> , <b>2012</b> , 379, 1022-8   | 4.0  | 133       |
| 100 | Depressive symptoms among rural Bangladeshi mothers: implications for infant development. <i>Journal of Child Psychology and Psychiatry and Allied Disciplines</i> , <b>2007</b> , 48, 764-72  | 7.9  | 114       |
| 99  | Risk of early-onset neonatal infection with maternal infection or colonization: a global systematic review and meta-analysis. <i>PLoS Medicine</i> , <b>2013</b> , 10, e1001502  | 11.6 | 110       |
| 98  | Simultaneous weekly supplementation of iron and zinc is associated with lower morbidity due to diarrhea and acute lower respiratory infection in Bangladeshi infants. <i>Journal of Nutrition</i> , <b>2003</b> , 133, 4150-7          | 4.1  | 101       |
| 97  | A randomized, placebo-controlled trial of the effect of zinc supplementation during pregnancy on pregnancy outcome in Bangladeshi urban poor. <i>American Journal of Clinical Nutrition</i> , <b>2000</b> , 71, 114-9                  | 7    | 99        |

|    |  |      |    |
|----|--|------|----|
| 96 | Maternal depressive symptoms and infant growth in rural Bangladesh. <i>American Journal of Clinical Nutrition</i> , <b>2009</b> , 89, 951S-957S  | 7    | 98 |
| 95 | Effectiveness of home-based management of newborn infections by community health workers in rural Bangladesh. <i>Pediatric Infectious Disease Journal</i> , <b>2009</b> , 28, 304-10   | 3.4  | 95 |
| 94 | Urinary tract infections in pregnancy in a rural population of Bangladesh: population-based prevalence, risk factors, etiology, and antibiotic resistance. <i>BMC Pregnancy and Childbirth</i> , <b>2019</b> , 20, 1   | 3.2  | 92 |
| 93 | Birth preparedness and complication readiness among slum women in Indore city, India. <i>Journal of Health, Population and Nutrition</i> , <b>2010</b> , 28, 383-91  | 2.5  | 87 |
| 92 | Factors affecting recruitment and retention of community health workers in a newborn care intervention in Bangladesh. <i>Human Resources for Health</i> , <b>2010</b> , 8, 12  | 4.6  | 87 |
| 91 | Effect of timing of first postnatal care home visit on neonatal mortality in Bangladesh: a observational cohort study. <i>BMJ, The</i> , <b>2009</b> , 339, b2826  | 5.9  | 85 |
| 90 | Evaluation of a cluster-randomized controlled trial of a package of community-based maternal and newborn interventions in Mirzapur, Bangladesh. <i>PLoS ONE</i> , <b>2010</b> , 5, e9696   | 3.7  | 79 |
| 89 | Malnutrition, cell-mediated immune deficiency, and diarrhea: a community-based longitudinal study in rural Bangladeshi children. <i>American Journal of Epidemiology</i> , <b>1993</b> , 137, 355-65   | 3.8  | 78 |
| 88 | Care seeking for neonatal illness in low- and middle-income countries: a systematic review. <i>PLoS Medicine</i> , <b>2012</b> , 9, e1001183   | 11.6 | 77 |
| 87 | Population-based incidence and etiology of community-acquired neonatal bacteremia in Mirzapur, Bangladesh: an observational study. <i>Journal of Infectious Diseases</i> , <b>2009</b> , 200, 906-15   | 7    | 70 |
| 86 | Safety and efficacy of alternative antibiotic regimens compared with 7 day injectable procaine benzylpenicillin and gentamicin for outpatient treatment of neonates and young infants with clinical signs of severe infection when referral is not possible: a randomised, open-label, equivalence trial. <i>The Lancet Global Health</i> , <b>2015</b> , 3, e279-87 | 13.6 | 62 |
| 85 | Validation of community health workers assessment of neonatal illness in rural Bangladesh. <i>Bulletin of the World Health Organization</i> , <b>2009</b> , 87, 12-9   | 8.2  | 60 |
| 84 | Newborn care in rural Uttar Pradesh. <i>Indian Journal of Pediatrics</i> , <b>2007</b> , 74, 241-7   | 3    | 58 |
| 83 | Comparison of antibiotic resistance and serotype composition of carriage and invasive pneumococci among Bangladeshi children: implications for treatment policy and vaccine formulation. <i>Journal of Clinical Microbiology</i> , <b>2003</b> , 41, 5582-7  | 9.7  | 58 |
| 82 | Determinants of postnatal care use at health facilities in rural Tanzania: multilevel analysis of a household survey. <i>BMC Pregnancy and Childbirth</i> , <b>2015</b> , 15, 282  | 3.2  | 52 |
| 81 | Community-based intervention packages for improving perinatal health in developing countries: a review of the evidence. <i>Seminars in Perinatology</i> , <b>2010</b> , 34, 462-76   | 3.3  | 52 |
| 80 | Newborn care practices at home and in health facilities in 4 regions of Ethiopia. <i>BMC Pediatrics</i> , <b>2013</b> , 13, 198  | 2.6  | 51 |
| 79 | Comparing modelled predictions of neonatal mortality impacts using LiST with observed results of community-based intervention trials in South Asia. <i>International Journal of Epidemiology</i> , <b>2010</b> , 39 Suppl 1, i11-20  | 7.8  | 44 |

|    |   |      |    |
|----|---|------|----|
| 78 | Care at first-level facilities for children with severe pneumonia in Bangladesh: a cohort study. <i>Lancet, The</i> , <b>2008</b> , 372, 822-30   | 4.0  | 44 |
| 77 | Incidence and risk factors of preterm birth in a rural Bangladeshi cohort. <i>BMC Pediatrics</i> , <b>2014</b> , 14, 112  | 2.6  | 40 |
| 76 | Community-based validation of assessment of newborn illnesses by trained community health workers in Sylhet district of Bangladesh. <i>Tropical Medicine and International Health</i> , <b>2009</b> , 14, 1448-56   | 2.3  | 37 |
| 75 | Household surveillance of severe neonatal illness by community health workers in Mirzapur, Bangladesh: coverage and compliance with referral. <i>Health Policy and Planning</i> , <b>2010</b> , 25, 112-24  | 3.4  | 36 |
| 74 | Epidemiological and clinical characteristics of acute and persistent diarrhoea in rural Bangladeshi children. <i>Acta Paediatrica, International Journal of Paediatrics</i> , <b>1992</b> , 381, 15-21  | 3.1  | 34 |
| 73 | Comparison of US Birth Weight References and the International Fetal and Newborn Growth Consortium for the 21st Century Standard. <i>JAMA Pediatrics</i> , <b>2015</b> , 169, e151438   | 8.3  | 33 |
| 72 | Patterns of maternal care seeking behaviours in rural Bangladesh. <i>Tropical Medicine and International Health</i> , <b>2007</b> , 12, 823-32  | 2.3  | 33 |
| 71 | Prevalence of early-onset neonatal infection among newborns of mothers with bacterial infection or colonization: a systematic review and meta-analysis. <i>BMC Infectious Diseases</i> , <b>2015</b> , 15, 118  | 4    | 32 |
| 70 | Economic evaluation of neonatal care packages in a cluster-randomized controlled trial in Sylhet, Bangladesh. <i>Bulletin of the World Health Organization</i> , <b>2013</b> , 91, 736-45   | 8.2  | 32 |
| 69 | Trends in use of referral hospital services for care of sick newborns in a community-based intervention in Tangail District, Bangladesh. <i>Journal of Health, Population and Nutrition</i> , <b>2006</b> , 24, 519-29 <sup>2.5</sup>                     |      | 32 |
| 68 | Treatment of infections in young infants in low- and middle-income countries: a systematic review and meta-analysis of frontline health worker diagnosis and antibiotic access. <i>PLoS Medicine</i> , <b>2014</b> , 11, e1001741                         | 11.6 | 31 |
| 67 | Validity of Newborn Clinical Assessment to Determine Gestational Age in Bangladesh. <i>Pediatrics</i> , <b>2016</b> , 138,  | 7.4  | 30 |
| 66 | The Effect of Integrating Family Planning with a Maternal and Newborn Health Program on Postpartum Contraceptive Use and Optimal Birth Spacing in Rural Bangladesh. <i>Studies in Family Planning</i> , <b>2015</b> , 46, 297-312                         | 3.4  | 30 |
| 65 | Animal husbandry practices in rural Bangladesh: potential risk factors for antimicrobial drug resistance and emerging diseases. <i>American Journal of Tropical Medicine and Hygiene</i> , <b>2013</b> , 89, 965-70                                       | 3.2  | 30 |
| 64 | Community-driven impact of a newborn-focused behavioral intervention on maternal health in Shivgarh, India. <i>International Journal of Gynecology and Obstetrics</i> , <b>2012</b> , 117, 48-55  | 4    | 29 |
| 63 | Contribution of community-based newborn health promotion to reducing inequities in healthy newborn care practices and knowledge: evidence of improvement from a three-district pilot program in Malawi. <i>BMC Public Health</i> , <b>2013</b> , 13, 1052 | 4.1  | 29 |
| 62 | Process evaluation of a community-based intervention promoting multiple maternal and neonatal care practices in rural Nepal. <i>BMC Pregnancy and Childbirth</i> , <b>2010</b> , 10, 31   | 3.2  | 27 |
| 61 | Effect of knowledge of community health workers on essential newborn health care: a study from rural India. <i>Health Policy and Planning</i> , <b>2012</b> , 27, 115-26  | 3.4  | 27 |

|    |   |      |    |
|----|---|------|----|
| 60 | Determinants and pattern of care seeking for preterm newborns in a rural Bangladeshi cohort. <i>BMC Health Services Research</i> , <b>2014</b> , 14, 417  | 2.9  | 25 |
| 59 | Impact of 4.0% chlorhexidine cleansing of the umbilical cord on mortality and omphalitis among newborns of Sylhet, Bangladesh: design of a community-based cluster randomized trial. <i>BMC Pediatrics</i> , <b>2009</b> , 9, 67  | 2.6  | 25 |
| 58 | Scientific rationale for study design of community-based simplified antibiotic therapy trials in newborns and young infants with clinically diagnosed severe infections or fast breathing in South Asia and sub-Saharan Africa. <i>Pediatric Infectious Disease Journal</i> , <b>2013</b> , 32 Suppl 1, S7-11 | 3.4  | 24 |
| 57 | Operations research to add postpartum family planning to maternal and neonatal health to improve birth spacing in Sylhet District, Bangladesh. <i>Global Health, Science and Practice</i> , <b>2013</b> , 1, 262-76   | 2.8  | 24 |
| 56 | Factors associated with delay in care-seeking for fatal neonatal illness in the Sylhet district of Bangladesh: results from a verbal and social autopsy study. <i>Journal of Global Health</i> , <b>2016</b> , 6, 010605  | 4.3  | 24 |
| 55 | Weekly iron supplementation does not block increases in serum zinc due to weekly zinc supplementation in Bangladeshi infants. <i>Journal of Nutrition</i> , <b>2005</b> , 135, 2187-91  | 4.1  | 23 |
| 54 | Vitamin D status of infants in northeastern rural Bangladesh: preliminary observations and a review of potential determinants. <i>Journal of Health, Population and Nutrition</i> , <b>2010</b> , 28, 458-69  | 2.5  | 21 |
| 53 | Validation of a clinical algorithm to identify neonates with severe illness during routine household visits in rural Bangladesh. <i>Archives of Disease in Childhood</i> , <b>2011</b> , 96, 1140-6   | 2.2  | 21 |
| 52 | Association between clinical type of diarrhoea and growth of children under 5 years in rural Bangladesh. <i>International Journal of Epidemiology</i> , <b>2000</b> , 29, 916-21  | 7.8  | 21 |
| 51 | Improving and sustaining quality of child health care through IMCI training and supervision: experience from rural Bangladesh. <i>Health Policy and Planning</i> , <b>2014</b> , 29, 753-62   | 3.4  | 20 |
| 50 | Screening and treatment of maternal genitourinary tract infections in early pregnancy to prevent preterm birth in rural Sylhet, Bangladesh: a cluster randomized trial. <i>BMC Pregnancy and Childbirth</i> , <b>2015</b> , 15, 326   | 3.2  | 17 |
| 49 | Safety and efficacy of simplified antibiotic regimens for outpatient treatment of serious infection in neonates and young infants 0-59 days of age in Bangladesh: design of a randomized controlled trial. <i>Pediatric Infectious Disease Journal</i> , <b>2013</b> , 32 Suppl 1, S12-8                      | 3.4  | 17 |
| 48 | Infection Surveillance Protocol for a Multicountry Population-based Study in South Asia to Determine the Incidence, Etiology and Risk Factors for Infections Among Young Infants of 0 to 59 Days Old. <i>Pediatric Infectious Disease Journal</i> , <b>2016</b> , 35, S9-15                                   | 3.4  | 17 |
| 47 | Effect of population-based antenatal screening and treatment of genitourinary tract infections on birth outcomes in Sylhet, Bangladesh (MIST): a cluster-randomised clinical trial. <i>The Lancet Global Health</i> , <b>2019</b> , 7, e148-e159  | 13.6 | 17 |
| 46 | Levels, timing, and etiology of stillbirths in Sylhet district of Bangladesh. <i>BMC Pregnancy and Childbirth</i> , <b>2011</b> , 11, 25  | 3.2  | 15 |
| 45 | Effect of a package of integrated demand- and supply-side interventions on facility delivery rates in rural Bangladesh: Implications for large-scale programs. <i>PLoS ONE</i> , <b>2017</b> , 12, e0186182   | 3.7  | 15 |
| 44 | Impact of integrating a postpartum family planning program into a community-based maternal and newborn health program on birth spacing and preterm birth in rural Bangladesh. <i>Journal of Global Health</i> , <b>2018</b> , 8, 020406   | 4.3  | 15 |
| 43 | Prenatal vitamin D supplementation and infant vitamin D status in Bangladesh. <i>Public Health Nutrition</i> , <b>2017</b> , 20, 1865-1873  | 3.3  | 14 |

|    |  |      |    |
|----|--|------|----|
| 42 | Population-based Incidence and Etiology of Community-acquired Neonatal Viral Infections in Bangladesh: A Community-based and Hospital-based Surveillance Study. <i>Pediatric Infectious Disease Journal</i> , <b>2015</b> , 34, 706-11   | 3.4  | 14 |
| 41 | Antepartum complications and perinatal mortality in rural Bangladesh. <i>BMC Pregnancy and Childbirth</i> , <b>2017</b> , 17, 81   | 3.2  | 13 |
| 40 | Ongoing trials of simplified antibiotic regimens for the treatment of serious infections in young infants in South Asia and sub-Saharan Africa: implications for policy. <i>Pediatric Infectious Disease Journal</i> , <b>2013</b> , 32 Suppl 1, S46-9   | 3.4  | 13 |
| 39 | Can facility delivery reduce the risk of intrapartum complications-related perinatal mortality? Findings from a cohort study. <i>Journal of Global Health</i> , <b>2018</b> , 8, 010408  | 4.3  | 12 |
| 38 | Performance of late pregnancy biometry for gestational age dating in low-income and middle-income countries: a prospective, multicountry, population-based cohort study from the WHO Alliance for Maternal and Newborn Health Improvement (AMANHI) Study Group. <i>The Lancet Global Health</i> , <b>2020</b> , 8, e545-e554 | 13.6 | 11 |
| 37 | Clean cord care practices and neonatal mortality: evidence from rural Uttar Pradesh, India. <i>Journal of Epidemiology and Community Health</i> , <b>2012</b> , 66, 755-8  | 5.1  | 11 |
| 36 | Incidence and risk factors of neonatal infections in a rural Bangladeshi population: a community-based prospective study. <i>Journal of Health, Population and Nutrition</i> , <b>2018</b> , 37, 6   | 2.5  | 10 |
| 35 | Equity improvements in maternal and newborn care indicators: results from the Bardiya district of Nepal. <i>Health Policy and Planning</i> , <b>2016</b> , 31, 405-14  | 3.4  | 10 |
| 34 | Levels and correlates of nutritional status of women of childbearing age in rural Bangladesh. <i>Public Health Nutrition</i> , <b>2018</b> , 21, 3037-3047   | 3.3  | 10 |
| 33 | Newborn care practices in rural Bangladesh: Implications for the adaptation of kangaroo mother care for community-based interventions. <i>Social Science and Medicine</i> , <b>2014</b> , 122, 21-30   | 5.1  | 10 |
| 32 | Sociocultural factors perpetuating the practices of early marriage and childbirth in Sylhet District, Bangladesh. <i>International Health</i> , <b>2015</b> , 7, 212-7   | 2.4  | 10 |
| 31 | Neonatal mortality risks among preterm births in a rural Bangladeshi cohort. <i>Paediatric and Perinatal Epidemiology</i> , <b>2014</b> , 28, 510-20   | 2.7  | 10 |
| 30 | Classification of Blood Culture Isolates Into Contaminants and Pathogens on the Basis of Clinical and Laboratory Data. <i>Pediatric Infectious Disease Journal</i> , <b>2016</b> , 35, S52-4   | 3.4  | 10 |
| 29 | Caregiver acceptability of the guidelines for managing young infants with possible serious bacterial infections (PSBI) in primary care facilities in rural Bangladesh. <i>PLoS ONE</i> , <b>2020</b> , 15, e0231490  | 3.7  | 10 |
| 28 | Community-based health workers achieve high coverage in neonatal intervention trials: a case study from Sylhet, Bangladesh. <i>Journal of Health, Population and Nutrition</i> , <b>2010</b> , 28, 610-8   | 2.5  | 9  |
| 27 | The effect of intrapartum antibiotics on early-onset neonatal sepsis in Dhaka, Bangladesh: a propensity score matched analysis. <i>BMC Pediatrics</i> , <b>2014</b> , 14, 104  | 2.6  | 8  |
| 26 | "You should go so that others can come"; the role of facilities in determining an early departure after childbirth in Morogoro Region, Tanzania. <i>BMC Pregnancy and Childbirth</i> , <b>2015</b> , 15, 328   | 3.2  | 8  |
| 25 | Pneumococcal Conjugate Vaccine impact assessment in Bangladesh. <i>Gates Open Research</i> , <b>2018</b> , 2, 21   | 2.4  | 8  |



|    |  |     |   |
|----|--|-----|---|
| 24 | Methods Employed in Monitoring and Evaluating Field and Laboratory Systems in the ANISA Study: Ensuring Quality. <i>Pediatric Infectious Disease Journal</i> , <b>2016</b> , 35, S39-44  | 3.4 | 8 |
| 23 | Early implementation of guidelines for managing young infants with possible serious bacterial infection in Bangladesh. <i>BMJ Global Health</i> , <b>2019</b> , 4, e001643   | 6.6 | 8 |
| 22 | Provider performance and facility readiness for managing infections in young infants in primary care facilities in rural Bangladesh. <i>PLoS ONE</i> , <b>2020</b> , 15, e0229988  | 3.7 | 8 |
| 21 | Strengthening the capabilities of families and communities to improve child health in low and middle income countries. <i>BMJ, The</i> , <b>2018</b> , 362, bmj.k2649  | 5.9 | 8 |
| 20 | Maternal short stature and under-weight status are independent risk factors for preterm birth and small for gestational age in rural Bangladesh. <i>European Journal of Clinical Nutrition</i> , <b>2019</b> , 73, 733-742   | 5.2 | 7 |
| 19 | Forecasting burden of long-term disability from neonatal conditions: results from the Projahnmo I trial, Sylhet, Bangladesh. <i>Health Policy and Planning</i> , <b>2013</b> , 28, 435-52  | 3.4 | 7 |
| 18 | Assessment of the proportion of neonates and children in low and middle income countries with access to a healthcare facility: A systematic review. <i>BMC Research Notes</i> , <b>2011</b> , 4, 536   | 2.3 | 7 |
| 17 | Implementation of the ANISA Protocol in Sylhet, Bangladesh: Challenges and Solutions. <i>Pediatric Infectious Disease Journal</i> , <b>2016</b> , 35, S55-9  | 3.4 | 7 |
| 16 | Clustering of neonatal deaths in Bangladesh: results from the Projahnmo studies. <i>Paediatric and Perinatal Epidemiology</i> , <b>2013</b> , 27, 165-71   | 2.7 | 6 |
| 15 | Patterns and Determinants of Care-Seeking for Antepartum and Intrapartum Complications in Rural Bangladesh: Results from a Cohort Study. <i>PLoS ONE</i> , <b>2016</b> , 11, e0167814  | 3.7 | 6 |
| 14 | Neonatal health program management in a resource-constrained setting in rural Uttar Pradesh, India. <i>International Journal of Health Planning and Management</i> , <b>2009</b> , 24, 173-84  | 2.2 | 5 |
| 13 | Utilization and perceptions of neonatal healthcare providers in rural Uttar Pradesh, India. <i>International Journal for Quality in Health Care</i> , <b>2011</b> , 23, 487-94   | 1.9 | 4 |
| 12 | Impact of community-based behaviour-change management on perceived neonatal morbidity: a cluster-randomized controlled trial in Shivgarh, Uttar Pradesh, India. <i>Journal of Tropical Pediatrics</i> , <b>2012</b> , 58, 286-91   | 1.2 | 4 |
| 11 | Effect of Birth Preparedness on Institutional Delivery in Semiurban Ethiopia: A Cross-Sectional Study. <i>Annals of Global Health</i> , <b>2019</b> , 85,  | 3.3 | 4 |
| 10 | Pregnancy intervals after stillbirth, neonatal death and spontaneous abortion and the risk of an adverse outcome in the next pregnancy in rural Bangladesh. <i>BMC Pregnancy and Childbirth</i> , <b>2019</b> , 19, 62   | 3.2 | 4 |
| 9  | Implementation research to support Bangladesh Ministry of Health and Family Welfare to implement its national guidelines for management of infections in young infants in two rural districts. <i>Journal of Health, Population and Nutrition</i> , <b>2019</b> , 38, 41 | 2.5 | 4 |
| 8  | Local understandings and current barriers to optimal birth intervals among recently delivered women in Sylhet District, Bangladesh. <i>International Health</i> , <b>2013</b> , 5, 266-72  | 2.4 | 3 |
| 7  | Population-based incidence and serotype distribution of invasive pneumococcal disease prior to introduction of conjugate pneumococcal vaccine in Bangladesh. <i>PLoS ONE</i> , <b>2020</b> , 15, e0228799  | 3.7 | 2 |

|   |  |     |   |
|---|--|-----|---|
| 6 | Vulnerability of newborns to environmental factors: findings from community based surveillance data in Bangladesh. <i>International Journal of Environmental Research and Public Health</i> , <b>2011</b> , 8, 3437-52                                 | 4.6 | 2 |
| 5 | Digital auscultation as a novel childhood pneumonia diagnostic tool for community clinics in Sylhet, Bangladesh: protocol for a cross-sectional study.. <i>BMJ Open</i> , <b>2022</b> , 12, e059630  | 3   | 2 |
| 4 | Management of fast breathing pneumonia in young infants aged 7 to 59 days by community level health workers: protocol for a multi-centre cluster randomized controlled trial. <i>International Journal of Clinical Trials</i> , <b>2020</b> , 7, 83-93 | 2.7 | 0 |
| 3 | Maternal serum lipidomics identifies lysophosphatidic acid as a predictor of small for gestational age neonates. <i>Molecular Omics</i> , <b>2021</b> , 17, 956-966  | 4.4 | 0 |
| 2 | Depressive symptoms among rural Bangladeshi mothers: implications for infant development. <i>Journal of Child Psychology and Psychiatry and Allied Disciplines</i> , <b>2007</b> , 070510003552005-???   | 7.9 |   |
| 1 | Validation of community health worker identification of maternal puerperal sepsis using a clinical diagnostic algorithm in Bangladesh and Pakistan.. <i>Journal of Global Health</i> , <b>2021</b> , 11, 04039   | 4.3 |   |