

Youwen Zhou

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11823425/publications.pdf>

Version: 2024-02-01

51
papers

3,116
citations

159585

30
h-index

182427

51
g-index

51
all docs

51
docs citations

51
times ranked

3927
citing authors

#	ARTICLE	IF	CITATIONS
1	Evidence-Based Clinical Practice Guidelines for Extramammary Paget Disease. <i>JAMA Oncology</i> , 2022, 8, 618.	7.1	46
2	Immunohistochemistry analysis reveals lysyl oxidase-like 3 as a novel prognostic marker for primary melanoma. <i>Melanoma Research</i> , 2021, 31, 173-177.	1.2	3
3	Vitiligo Skin Biomarkers Associated With Favorable Therapeutic Response. <i>Frontiers in Immunology</i> , 2021, 12, 613031.	4.8	7
4	Computer-Aided Discovery of Small Molecule Inhibitors of Thymocyte Selection-Associated High Mobility Group Box Protein (TOX) as Potential Therapeutics for Cutaneous T-Cell Lymphomas. <i>Molecules</i> , 2019, 24, 3459.	3.8	6
5	Whole-Exome Sequencing Reveals Frequent Mutations in Chromatin Remodeling Genes in Mammary and Extramammary Paget's Diseases. <i>Journal of Investigative Dermatology</i> , 2019, 139, 789-795.	0.7	35
6	Targeting Hypoxia-Induced Carbonic Anhydrase IX Enhances Immune-Checkpoint Blockade Locally and Systemically. <i>Cancer Immunology Research</i> , 2019, 7, 1064-1078.	3.4	104
7	Existing and Emerging Therapies for Cutaneous T-Cell Lymphoma. <i>Journal of Cutaneous Medicine and Surgery</i> , 2019, 23, 319-327.	1.2	5
8	IL-10 is overexpressed in human cutaneous T-cell lymphoma and is required for maximal tumor growth in a mouse model. <i>Leukemia and Lymphoma</i> , 2019, 60, 1244-1252.	1.3	14
9	Transcriptome analyses reveal FOXA1 dysregulation in mammary and extramammary Paget's disease. <i>Human Pathology</i> , 2018, 77, 152-158.	2.0	19
10	Prognostic significance of the expression of nuclear eukaryotic translation initiation factor 5A2 in human melanoma. <i>Oncology Letters</i> , 2016, 12, 3089-3100.	1.8	7
11	The prevalence of anxiety and depression in patients with or without hyperhidrosis (HH). <i>Journal of the American Academy of Dermatology</i> , 2016, 75, 1126-1133.	1.2	59
12	Hyperhidrosis Prevalence and Demographical Characteristics in Dermatology Outpatients in Shanghai and Vancouver. <i>PLoS ONE</i> , 2016, 11, e0153719.	2.5	40
13	Evidence of an oncogenic role of aberrant TOX activation in cutaneous T-cell lymphoma. <i>Blood</i> , 2015, 125, 1435-1443.	1.4	61
14	The Use of Transcriptional Profiling to Improve Personalized Diagnosis and Management of Cutaneous T-cell Lymphoma (CTCL). <i>Clinical Cancer Research</i> , 2015, 21, 2820-2829.	7.0	76
15	Ectopic expression of a novel CD22 splice-variant regulates survival and proliferation in malignant T cells from cutaneous T cell lymphoma (CTCL) patients. <i>Oncotarget</i> , 2015, 6, 14374-14384.	1.8	4
16	IL-15 and IL-17F are differentially regulated and expressed in mycosis fungoides (MF). <i>Cell Cycle</i> , 2014, 13, 1306-1312.	2.6	27
17	Analysis of STAT4 expression in cutaneous T-cell lymphoma (CTCL) patients and patient-derived cell lines. <i>Cell Cycle</i> , 2014, 13, 2975-2982.	2.6	62
18	CXCL10 Is Critical for the Progression and Maintenance of Depigmentation in a Mouse Model of Vitiligo. <i>Science Translational Medicine</i> , 2014, 6, 223ra23.	12.4	333

#	ARTICLE	IF	CITATIONS
19	Ectopic expression of embryonic stem cell and other developmental genes in cutaneous T-cell lymphoma. <i>Oncolmmunology</i> , 2014, 3, e970025.	4.6	38
20	SATB1 overexpression promotes malignant T-cell proliferation in cutaneous CD30+ lymphoproliferative disease by repressing p21. <i>Blood</i> , 2014, 123, 3452-3461.	1.4	31
21	Ectopic Expression of Cancer-Testis Antigens in Cutaneous T-cell Lymphoma Patients. <i>Clinical Cancer Research</i> , 2014, 20, 3799-3808.	7.0	40
22	Depletion of M2-Like Tumor-Associated Macrophages Delays Cutaneous T-Cell Lymphoma Development In Vivo. <i>Journal of Investigative Dermatology</i> , 2014, 134, 2814-2822.	0.7	102
23	Thymocyte selection-associated high mobility group box gene (TOX) is aberrantly over-expressed in mycosis fungoides and correlates with poor prognosis. <i>Oncotarget</i> , 2014, 5, 4418-4425.	1.8	55
24	Loss of BCL7A expression correlates with poor disease prognosis in patients with early-stage cutaneous T-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2013, 54, 653-654.	1.3	20
25	Elucidating the role of interleukin-17F in cutaneous T-cell lymphoma. <i>Blood</i> , 2013, 122, 943-950.	1.4	78
26	Potential Role of Neurogenic Inflammatory Factors in the Pathogenesis of Vitiligo. <i>Journal of Cutaneous Medicine and Surgery</i> , 2012, 16, 230-244.	1.2	19
27	Molecular Markers of Early-Stage Mycosis Fungoides. <i>Journal of Investigative Dermatology</i> , 2012, 132, 1698-1706.	0.7	88
28	Transcriptome Analysis Reveals Markers of Aberrantly Activated Innate Immunity in Vitiligo Lesional and Non-Lesional Skin. <i>PLoS ONE</i> , 2012, 7, e51040.	2.5	83
29	Deficiency of SATB1 expression in S α zary cells causes apoptosis resistance by regulating FasL/CD95L transcription. <i>Blood</i> , 2011, 117, 3826-3835.	1.4	49
30	Collagen Triple Helix Repeat Containing 1 Promotes Melanoma Cell Adhesion and Survival. <i>Journal of Cutaneous Medicine and Surgery</i> , 2011, 15, 103-110.	1.2	30
31	Genome-wide association study for vitiligo identifies susceptibility loci at 6q27 and the MHC. <i>Nature Genetics</i> , 2010, 42, 614-618.	21.4	174
32	Platelet-derived Growth Factor Receptor Alpha Gene Mutations in Vitiligo Vulgaris. <i>Acta Dermato-Venereologica</i> , 2010, 90, 131-135.	1.3	15
33	Alpha 1 antichymotrypsin is aberrantly expressed during melanoma progression and predicts poor survival for patients with metastatic melanoma. <i>Pigment Cell and Melanoma Research</i> , 2010, 23, 575-586.	3.3	18
34	Genetic Variation of Promoter Sequence Modulates XBP1 Expression and Genetic Risk for Vitiligo. <i>PLoS Genetics</i> , 2009, 5, e1000523.	3.5	77
35	Identification of tyrosine kinase, HCK, and tumor suppressor, BIN1, as potential mediators of AHI-1 oncogene in primary and transformed CTCL cells. <i>Blood</i> , 2009, 113, 4646-4655.	1.4	31
36	Endothelin signaling axis activates osteopontin expression through PI3 kinase pathway in A375 melanoma cells. <i>Journal of Dermatological Science</i> , 2008, 52, 130-132.	1.9	6

#	ARTICLE	IF	CITATIONS
37	Endothelin-3 Is Produced by Metastatic Melanoma Cells and Promotes Melanoma Cell Survival. <i>Journal of Cutaneous Medicine and Surgery</i> , 2008, 12, 64-70.	1.2	18
38	Efficacy of a Day-Care Program in the Treatment of Psoriasis. <i>Journal of Cutaneous Medicine and Surgery</i> , 2008, 12, 211-216.	1.2	1
39	Evidence for Two Susceptibility Loci on Chromosomes 22q12 and 6p21 in Chinese Generalized Vitiligo Families. <i>Journal of Investigative Dermatology</i> , 2007, 127, 2552-2557.	0.7	36
40	Expression of Endothelins and Their Receptors in Nonmelanoma Skin Cancers. <i>Journal of Cutaneous Medicine and Surgery</i> , 2006, 10, 269-276.	1.2	7
41	Aberrant Expression of Collagen Triple Helix Repeat Containing 1 in Human Solid Cancers. <i>Clinical Cancer Research</i> , 2006, 12, 3716-3722.	7.0	133
42	Osteopontin Expression Correlates with Melanoma Invasion. <i>Journal of Investigative Dermatology</i> , 2005, 124, 1044-1052.	0.7	122
43	Stable Suppression of a Novel Oncogene, AHI-1, in Human Cutaneous T-Cell Leukemia Cells Normalizes Its Transforming Activity In Vitro and In Vivo and Aberrant Expression of AHI-1 Is Also Present in Leukemic Sezary Cells from Patients with Sezary Syndrome.. <i>Blood</i> , 2005, 106, 2605-2605.	1.4	32
44	Mycophenolate Mofetil (CellCept®) for Psoriasis: A Two-Center, Prospective, Open-Label Clinical Trial. <i>Journal of Cutaneous Medicine and Surgery</i> , 2003, 7, 193-197.	1.2	38
45	Increased expression of integrin-linked kinase is correlated with melanoma progression and poor patient survival. <i>Clinical Cancer Research</i> , 2003, 9, 4409-14.	7.0	110
46	Aberrant expression of T-plastin in Sezary cells. <i>Cancer Research</i> , 2003, 63, 7122-7.	0.9	60
47	Effects of UVB irradiation on keratinocyte growth factor (KGF) and receptor (KGFR) expression in cultured human keratinocytes. <i>Experimental Dermatology</i> , 1996, 5, 138-144.	2.9	11
48	Expression of the UNC-5 guidance receptor in the touch neurons of <i>C. elegans</i> steers their axons dorsally. <i>Nature</i> , 1993, 364, 327-330.	27.8	229
49	UNC-5, a transmembrane protein with immunoglobulin and thrombospondin type 1 domains, guides cell and pioneer axon migrations in <i>C. elegans</i> . <i>Cell</i> , 1992, 71, 289-299.	28.9	389
50	Glomerular basement membrane degradation by endogenous cysteine proteinases in isolated rat glomeruli. <i>Kidney International</i> , 1990, 38, 395-401.	5.2	37
51	The role of aspartic and cysteine proteinases in albumin degradation by rat kidney cortical lysosomes. <i>Archives of Biochemistry and Biophysics</i> , 1987, 256, 687-691.	3.0	31