

Jason Shearer

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1182349/jason-shearer-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

72
papers

2,555
citations

33
h-index

48
g-index

75
ext. papers

2,777
ext. citations

8.9
avg, IF

5.06
L-index

#	Paper	IF	Citations
72	Controlled Protonation of [2Fe-2S] Leading to MitoNEET Analogues and Concurrent Cluster Modification. <i>Inorganic Chemistry</i> , 2021 , 60, 16074-16078	5.1	
71	The Oxo-Wall Remains Intact: A Tetrahedrally Distorted Co(IV)-Oxo Complex. <i>Journal of the American Chemical Society</i> , 2021 , 143, 16943-16959	16.4	0
70	Access to Metal Centers and Fluxional Hydride Coordination Integral for CO Insertion into [Fe(H)] Clusters. <i>Inorganic Chemistry</i> , 2021 , 60, 7228-7239	5.1	0
69	Dinitrogen Insertion and Cleavage by a Metal-Metal Bonded Tricobalt(I) Cluster. <i>Journal of the American Chemical Society</i> , 2021 , 143, 5649-5653	16.4	6
68	Scaffold-based [Fe]-hydrogenase model: H activation initiates Fe(0)-hydride extrusion and non-biomimetic hydride transfer. <i>Chemical Science</i> , 2021 , 12, 12838-12846	9.4	0
67	Structure and Unprecedented Reactivity of a Mononuclear Nonheme Cobalt(III) Iodosylbenzene Complex. <i>Angewandte Chemie</i> , 2020 , 132, 13683-13687	3.6	
66	Structure and Unprecedented Reactivity of a Mononuclear Nonheme Cobalt(III) Iodosylbenzene Complex. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 13581-13585	16.4	12
65	Reduction of CO by a masked two-coordinate cobalt(i) complex and characterization of a proposed oxodicobalt(ii) intermediate. <i>Chemical Science</i> , 2019 , 10, 918-929	9.4	23
64	pH Dependent Reversible Formation of a Binuclear Ni ₂ Metal-Center within a Peptide Scaffold. <i>Inorganics</i> , 2019 , 7, 90	2.9	0
63	A Biochemical Nickel(I) State Supports Nucleophilic Alkyl Addition: A Roadmap for Methyl Reactivity in Acetyl Coenzyme A Synthase. <i>Inorganic Chemistry</i> , 2019 , 58, 8969-8982	5.1	11
62	Chalcogen Impact on Covalency within Molecular [Cu(E)] Clusters (E = O, S, Se): A Synthetic, Spectroscopic, and Computational Study. <i>Inorganic Chemistry</i> , 2018 , 57, 11382-11392	5.1	7
61	Interactions of Metal-Based and Ligand-Based Electronic Spins in Neutral Tripyrrindione Dimers. <i>Inorganic Chemistry</i> , 2017 , 56, 6755-6762	5.1	16
60	Synthesis and reactivity of a mononuclear non-haem cobalt(IV)-oxo complex. <i>Nature Communications</i> , 2017 , 8, 14839	17.4	94
59	Sequence proximity between Cu(II) and Cu(I) binding sites of human copper transporter 1 model peptides defines reactivity with ascorbate and O ₂ . <i>Journal of Inorganic Biochemistry</i> , 2016 , 158, 70-76	4.2	25
58	A [3Fe-3S](3+) cluster with exclusively Sulfide donors. <i>Chemical Communications</i> , 2016 , 52, 1174-7	5.8	23
57	Adiabaticity of the proton-coupled electron-transfer step in the reduction of superoxide effected by nickel-containing superoxide dismutase metalloprotein-based mimics. <i>Journal of Physical Chemistry B</i> , 2015 , 119, 5453-61	3.4	4
56	Model Peptide Studies Reveal a Mixed Histidine-Methionine Cu(I) Binding Site at the N-Terminus of Human Copper Transporter 1. <i>Inorganic Chemistry</i> , 2015 , 54, 8544-51	5.1	33

55	A Redox-Active, Compact Molecule for Cross-Linking Amyloidogenic Peptides into Nontoxic, Off-Pathway Aggregates: In Vitro and In Vivo Efficacy and Molecular Mechanisms. <i>Journal of the American Chemical Society</i> , 2015 , 137, 14785-97	16.4	52
54	An Air- and Water-Tolerant Zinc Hydride Cluster That Reacts Selectively With CO ₂ . <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 7047-50	16.4	29
53	Tripyrrindione as a Redox-Active Ligand: Palladium(II) Coordination in Three Redox States. <i>Angewandte Chemie - International Edition</i> , 2015 , 54, 14894-7	16.4	31
52	Tripyrrindione as a Redox-Active Ligand: Palladium(II) Coordination in Three Redox States. <i>Angewandte Chemie</i> , 2015 , 127, 15107-15110	3.6	12
51	Subtle Modulation of Cu ₄ X ₄ L ₂ Phosphine Cluster Cores Leads to Changes in Luminescence. <i>Inorganic Chemistry</i> , 2015 , 54, 6245-56	5.1	43
50	A nickel phosphine complex as a fast and efficient hydrogen production catalyst. <i>Journal of the American Chemical Society</i> , 2015 , 137, 1109-15	16.4	116
49	Cysteinate protonation and water hydrogen bonding at the active-site of a nickel superoxide dismutase metalloprotein-based mimic: implications for the mechanism of superoxide reduction. <i>Journal of the American Chemical Society</i> , 2014 , 136, 16009-22	16.4	16
48	Isolation of a (dinitrogen)tricopper(I) complex. <i>Journal of the American Chemical Society</i> , 2014 , 136, 13502-5	16.4	59
47	Transformation of a mononitrosyl iron complex to a [2Fe-2S] cluster by a cysteine analogue. <i>Journal of the American Chemical Society</i> , 2014 , 136, 7229-32	16.4	21
46	Insight into the structure and mechanism of nickel-containing superoxide dismutase derived from peptide-based mimics. <i>Accounts of Chemical Research</i> , 2014 , 47, 2332-41	24.3	46
45	Crystallographic and computational studies of luminescent, binuclear gold(I) complexes, Au(I) ₂ (Ph ₂ P(CH ₂) _n PPh ₂) ₂ I ₂ (n = 3-6). <i>Inorganic Chemistry</i> , 2013 , 52, 823-31	5.1	17
44	Dioxygen and superoxide stability of metalloprotein based mimics of nickel containing superoxide dismutase: the influence of amine/amidate vs. bis-amidate ligation. <i>Journal of Inorganic Biochemistry</i> , 2013 , 129, 145-9	4.2	10
43	Novel alkoxide cluster topologies featuring rare seesaw geometry at transition metal centers. <i>Chemistry - A European Journal</i> , 2013 , 19, 12225-8	4.8	27
42	Use of a metalloprotein-based mimic provides evidence for a proton-coupled electron-transfer mechanism for superoxide reduction by nickel-containing superoxide dismutase. <i>Angewandte Chemie - International Edition</i> , 2013 , 52, 2569-72	16.4	21
41	Use of a Metalloprotein-Based Mimic Provides Evidence for a Proton-Coupled Electron-Transfer Mechanism for Superoxide Reduction by Nickel-Containing Superoxide Dismutase. <i>Angewandte Chemie</i> , 2013 , 125, 2629-2632	3.6	4
40	Characterization and dioxygen reactivity of a new series of coordinatively unsaturated thiolate-ligated manganese(II) complexes. <i>Inorganic Chemistry</i> , 2012 , 51, 6633-44	5.1	33
39	Influence of sequential thiolate oxidation on a nitrile hydratase mimic probed by multiedge X-ray absorption spectroscopy. <i>Inorganic Chemistry</i> , 2012 , 51, 6032-45	5.1	23
38	Modulation of luminescence by subtle anion-cation and anion-π interactions in a trigonal Au(I)⋯Cu(I) complex. <i>Inorganic Chemistry</i> , 2012 , 51, 1207-9	5.1	37

37	One octarepeat expansion to the human prion protein alters both the Zn ²⁺ and Cu ²⁺ coordination environments within the octarepeat domain. <i>Inorganic Chemistry</i> , 2011 , 50, 1173-5	5.1	10
36	Phenol nitration induced by an {Fe(NO) ₂ }(10) dinitrosyl iron complex. <i>Journal of the American Chemical Society</i> , 2011 , 133, 1184-7	16.4	53
35	Bisamidate and mixed amine/amidate NiN ₂ S ₂ complexes as models for nickel-containing acetyl coenzyme A synthase and superoxide dismutase: an experimental and computational study. <i>Inorganic Chemistry</i> , 2010 , 49, 5393-406	5.1	60
34	Use of metalloprotein based mimics demonstrates that the metalloprotein nitrile hydratase requires two oxidized cysteines for catalytic activity. <i>Inorganic Chemistry</i> , 2010 , 49, 9064-77	5.1	19
33	Cu K-edge X-ray absorption spectroscopy reveals differential copper coordination within amyloid- β oligomers compared to amyloid- β monomers. <i>Chemical Communications</i> , 2010 , 46, 9137-9	5.8	37
32	Metalloprotein based mimics with substituted histidines approximate a key hydrogen bonding network in the metalloenzyme nickel superoxide dismutase. <i>Inorganic Chemistry</i> , 2009 , 48, 10560-71	5.1	47
31	Luminescent copper(I) halide butterfly dimers coordinated to [Au(CH ₃ imCH ₂ py) ₂]BF ₄ and [Au(CH ₃ imCH ₂ quin) ₂]BF ₄ . <i>Inorganic Chemistry</i> , 2009 , 48, 11362-75	5.1	37
30	The amyloid-beta peptide of Alzheimer's disease binds Cu(I) in a linear bis-his coordination environment: insight into a possible neuroprotective mechanism for the amyloid-beta peptide. <i>Journal of the American Chemical Society</i> , 2008 , 130, 17826-35	16.4	155
29	Influence of sequential guanidinium methylation on the energetics of the guanidinium...guanine dimer and guanidinium...guanine...cytosine trimer: implications for the control of protein...DNA interactions by arginine methyltransferases. <i>Journal of Physical Chemistry B</i> , 2008 , 112, 16995-7002	3.4	2
28	Probing variable amine/amide ligation in Ni(II)N ₂ S ₂ complexes using sulfur K-edge and nickel L-edge X-ray absorption spectroscopies: implications for the active site of nickel superoxide dismutase. <i>Inorganic Chemistry</i> , 2008 , 47, 2649-60	5.1	41
27	Properties of square-pyramidal alkyl-thiolate Fe(III) complexes, including an analogue of the unmodified form of nitrile hydratase. <i>Inorganic Chemistry</i> , 2008 , 47, 11228-36	5.1	25
26	Both Met(109) and Met(112) are utilized for Cu(II) coordination by the amyloidogenic fragment of the human prion protein at physiological pH. <i>Journal of Inorganic Biochemistry</i> , 2008 , 102, 2103-13	4.2	27
25	The copper(II) adduct of the unstructured region of the amyloidogenic fragment derived from the human prion protein is redox-active at physiological pH. <i>Inorganic Chemistry</i> , 2007 , 46, 710-9	5.1	46
24	Probing variable axial ligation in nickel superoxide dismutase utilizing metalloprotein-based models: insight into the superoxide disproportionation mechanism. <i>Journal of the American Chemical Society</i> , 2007 , 129, 14605-18	16.4	58
23	Ni K-edge XAS suggests that coordination of Ni(II) to the unstructured amyloidogenic region of the human prion protein produces a Ni(2) bis- μ -hydroxo dimer. <i>Journal of Inorganic Biochemistry</i> , 2007 , 101, 370-3	4.2	9
22	Periodic trends within a series of five-coordinate thiolate-ligated [MII(SMe ₂ N ₄ (tren))] ⁺ (M = Mn, Fe, Co, Ni, Cu, Zn) complexes, including a rare example of a stable CuII-thiolate. <i>Inorganic Chemistry</i> , 2007 , 46, 9267-77	5.1	38
21	Manganese complexes of 1,3,5-triaza-7-phosphadadamantane (PTA): the first nitrogen-bound transition-metal complex of PTA. <i>Inorganic Chemistry</i> , 2006 , 45, 3481-3	5.1	49
20	The influence of amine/amide versus bisamide coordination in nickel superoxide dismutase. <i>Inorganic Chemistry</i> , 2006 , 45, 10552-66	5.1	47

19	[Me ₄ N](Ni(II)(BEAAM)): a synthetic model for nickel superoxide dismutase that contains Ni in a mixed amine/amide coordination environment. <i>Inorganic Chemistry</i> , 2006 , 45, 9637-9	5.1	54
18	A nickel superoxide dismutase maquette that reproduces the spectroscopic and functional properties of the metalloenzyme. <i>Inorganic Chemistry</i> , 2006 , 45, 2358-60	5.1	71
17	Synthesis, characterization, and crystal structure of a quadruply bonded dimolybdenum(II) complex containing the water-soluble phosphine 1,3,5-triaza-7-phosphaadamantane (PTA). <i>Inorganica Chimica Acta</i> , 2006 , 359, 283-288	2.7	16
16	Substrate oxidation by copper-dioxygen adducts: mechanistic considerations. <i>Journal of the American Chemical Society</i> , 2005 , 127, 5469-83	16.4	92
15	Steric and electronic control over the reactivity of a thiolate-ligated Fe(II) complex with dioxygen and superoxide: reversible μ -oxo dimer formation. <i>Inorganic Chemistry</i> , 2004 , 43, 7682-90	5.1	38
14	Heme/Cu/O ₂ reactivity: change in Fe(III)-(O ₂ ²⁻)-Cu(II) unit peroxo binding geometry effected by tridentate copper chelation. <i>Journal of the American Chemical Society</i> , 2004 , 126, 12716-7	16.4	34
13	Tuning copper-dioxygen reactivity and exogenous substrate oxidations via alterations in ligand electronics. <i>Journal of the American Chemical Society</i> , 2003 , 125, 634-5	16.4	84
12	Distinguishing rate-limiting electron versus H-atom transfers in Cu ₂ O ₂ -mediated oxidative N-dealkylations: application of inter- versus intramolecular kinetic isotope effects. <i>Journal of the American Chemical Society</i> , 2003 , 125, 12670-1	16.4	62
11	How does cyanide inhibit superoxide reductase? Insight from synthetic Fe(III)N ₄ S model complexes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 3671-6	11.5	33
10	Preparation and properties of [Ni(II)(BEES)(Cl)](BPh ₄): a Ni(II) complex in a mixed nitrogen/thioether coordination environment. <i>Inorganica Chimica Acta</i> , 2002 , 336, 61-64	2.7	1
9	The first example of a nitrile hydratase model complex that reversibly binds nitriles. <i>Journal of the American Chemical Society</i> , 2002 , 124, 11417-28	16.4	49
8	Synthetic models for the cysteinylated non-heme iron enzyme superoxide reductase: observation and structural characterization by XAS of an Fe(III)-OOH intermediate. <i>Journal of the American Chemical Society</i> , 2002 , 124, 11709-17	16.4	78
7	Enhancing reactivity via structural distortion. <i>Inorganic Chemistry</i> , 2002 , 41, 3128-36	5.1	25
6	Modeling the reactivity of superoxide reducing metalloenzymes with a nitrogen and sulfur coordinated iron complex. <i>Inorganic Chemistry</i> , 2001 , 40, 5483-4	5.1	33
5	Why is there an "inert" metal center in the active site of nitrile hydratase? Reactivity and ligand dissociation from a five-coordinate Co(III) nitrile hydratase model. <i>Journal of the American Chemical Society</i> , 2001 , 123, 463-8	16.4	59
4	How Do Oxidized Thiolate Ligands Affect the Electronic and Reactivity Properties of a Nitrile Hydratase Model Compound?. <i>Journal of the American Chemical Society</i> , 2000 , 122, 8299-8300	16.4	56
3	A Co(III) complex in a mixed sulfur/nitrogen ligand environment: modeling the substrate- and product-bound forms of the metalloenzyme thiocyanate hydrolase. <i>Inorganic Chemistry</i> , 2000 , 39, 4998-9	5.1	14
2	A Ni(Salen)-Biotin Conjugate for Rapid Isolation of Accessible DNA. <i>Journal of the American Chemical Society</i> , 2000 , 122, 9046-9047	16.4	22

1 Alkylation of Nucleic Acids by a Model Quinone Methide. *Journal of the American Chemical Society*,
1999, 121, 6773-6779

16.4 123