

Katsuhiko Kohara

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11833213/publications.pdf>

Version: 2024-02-01

169
papers

9,363
citations

57681

46
h-index

53065

89
g-index

178
all docs

178
docs citations

178
times ranked

14059
citing authors

#	ARTICLE	IF	CITATIONS
1	Association Between Body Mass Index and Functional Dyspepsia in Young Japanese People. <i>Journal of Neurogastroenterology and Motility</i> , 2022, 28, 276-282.	0.8	6
2	Multi-ancestry genetic study of type 2 diabetes highlights the power of diverse populations for discovery and translation. <i>Nature Genetics</i> , 2022, 54, 560-572.	9.4	250
3	Association Between Eating Behavior, Frequency of Meals, and Functional Dyspepsia in Young Japanese Population. <i>Journal of Neurogastroenterology and Motility</i> , 2022, 28, 418-423.	0.8	3
4	Creatinine-to-cystatin C ratio as a marker of skeletal muscle mass in older adults: J-SHIPP study. <i>Clinical Nutrition</i> , 2020, 39, 1857-1862.	2.3	40
5	Identification of type 2 diabetes loci in 433,540 East Asian individuals. <i>Nature</i> , 2020, 582, 240-245.	13.7	282
6	The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2019). <i>Hypertension Research</i> , 2019, 42, 1235-1481.	1.5	1,047
7	Distribution of cerebral microbleeds in the East and West. <i>Neurology</i> , 2019, 92, e1086-e1097.	1.5	53
8	Hyperglycemia in non-obese patients with type 2 diabetes is associated with low muscle mass: The Multicenter Study for Clarifying Evidence for Sarcopenia in Patients with Diabetes Mellitus. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1471-1479.	1.1	91
9	Steno-Stiffness Approach for Cardiovascular Disease Risk Assessment in Primary Prevention. <i>Hypertension</i> , 2019, 73, 508-513.	1.3	9
10	Clinical significance of an elevated ankle-brachial index differs depending on the amount of appendicular muscle mass: the J-SHIPP and Nagahama studies. <i>Hypertension Research</i> , 2018, 41, 354-362.	1.5	7
11	Simultaneously Measured Interarm Blood Pressure Difference and Stroke. <i>Hypertension</i> , 2018, 71, 1030-1038.	1.3	22
12	A double-blind, placebo-controlled, randomised clinical study of the effect of pork collagen peptide supplementation on atherosclerosis in healthy older individuals. <i>Bioscience, Biotechnology and Biochemistry</i> , 2018, 82, 893-895.	0.6	12
13	Correlation between the 24-h urinary angiotensinogen or aldosterone level and muscle mass: Japan shimanami health promoting program study. <i>Hypertension Research</i> , 2018, 41, 326-333.	1.5	4
14	Interethnic analyses of blood pressure loci in populations of East Asian and European descent. <i>Nature Communications</i> , 2018, 9, 5052.	5.8	75
15	Ankle-brachial index measured by oscillometry is predictive for cardiovascular disease and premature death in the Japanese population: An individual participant data meta-analysis. <i>Atherosclerosis</i> , 2018, 275, 141-148.	0.4	34
16	High central blood pressure is associated with incident cardiovascular events in treated hypertensives: the ABC-J II Study. <i>Hypertension Research</i> , 2018, 41, 947-956.	1.5	14
17	Habitual hot water bathing protects cardiovascular function in middle-aged to elderly Japanese subjects. <i>Scientific Reports</i> , 2018, 8, 8687.	1.6	9
18	Brachial-Ankle Pulse Wave Velocity and the Risk Prediction of Cardiovascular Disease. <i>Hypertension</i> , 2017, 69, 1045-1052.	1.3	382

#	ARTICLE	IF	CITATIONS
19	Cross-sectional study of equal producer status and cognitive impairment in older adults. <i>Geriatrics and Gerontology International</i> , 2017, 17, 2103-2108.	0.7	15
20	Muscle mass decline, arterial stiffness, white matter hyperintensity, and cognitive impairment: Japan Shimanami Health Promoting Program study. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2017, 8, 557-566.	2.9	67
21	Dual Effects of a RETN Single Nucleotide Polymorphism (SNP) at 420 on Plasma Resistin: Genotype and DNA Methylation. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 884-892.	1.8	12
22	Office-based simple frailty score and central blood pressure predict mild cognitive impairment in an apparently healthy Japanese population: J-SHIPP study. <i>Scientific Reports</i> , 2017, 7, 46419.	1.6	11
23	Proposed Cutoff Value of Brachial-Ankle Pulse Wave Velocity for the Management of Hypertension. <i>Circulation Journal</i> , 2017, 81, 1540-1542.	0.7	36
24	Impact of common genetic determinants of Hemoglobin A1c on type 2 diabetes risk and diagnosis in ancestrally diverse populations: A transethnic genome-wide meta-analysis. <i>PLoS Medicine</i> , 2017, 14, e1002383.	3.9	341
25	B-type natriuretic peptide is a determinant of the nocturnal increase in blood pressure independently of arterial hypertrophy and hypoxia. <i>Journal of Hypertension</i> , 2016, 34, 2393-2401.	0.3	12
26	Genome-wide association study of plasma resistin levels identified rs1423096 and rs10401670 as possible functional variants in the Japanese population. <i>Physiological Genomics</i> , 2016, 48, 874-881.	1.0	11
27	Mendelian randomization analysis in three Japanese populations supports a causal role of alcohol consumption in lowering low-density lipid cholesterol levels and particle numbers. <i>Atherosclerosis</i> , 2016, 254, 242-248.	0.4	27
28	Skin Autofluorescence Examination as a Diagnostic Tool for Mild Cognitive Impairment in Healthy People. <i>Journal of Alzheimer's Disease</i> , 2016, 55, 1481-1487.	1.2	9
29	Orthostatic hypertension as a predisposing factor for masked hypertension: the J-SHIPP study. <i>Hypertension Research</i> , 2016, 39, 664-669.	1.5	31
30	Association of office-based frailty score with hypertensive end organ damage in the J-SHIPP cross-sectional study. <i>International Journal of Cardiology</i> , 2016, 216, 25-31.	0.8	16
31	Diabetic mice exhibited a peculiar alteration in body composition with exaggerated ectopic fat deposition after muscle injury due to anomalous cell differentiation. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2016, 7, 213-224.	2.9	28
32	Synergistic association of changes in serum uric acid and triglycerides with changes in insulin resistance after walking exercise in community-dwelling older women. <i>Endocrine Research</i> , 2016, 41, 116-123.	0.6	13
33	Usefulness of the second derivative of the finger photoplethysmogram for assessment of end-organ damage: the J-SHIPP study. <i>Hypertension Research</i> , 2016, 39, 552-556.	1.5	12
34	Determinants of change in insulin resistance response to Nordic walking in community-dwelling elderly women. <i>Journal of Clinical Gerontology and Geriatrics</i> , 2015, 6, 100-105.	0.7	2
35	Brachial-ankle pulse wave velocity is a predictor of walking distance in community-dwelling adults. <i>Aging Clinical and Experimental Research</i> , 2015, 27, 187-193.	1.4	3
36	Portable indices for sarcopenia are associated with pressure wave reflection and central pulse pressure. <i>Journal of Hypertension</i> , 2015, 33, 314-322.	0.3	24

#	ARTICLE	IF	CITATIONS
37	Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. <i>Nature Genetics</i> , 2015, 47, 1282-1293.	9.4	294
38	Association of Postural Instability With Asymptomatic Cerebrovascular Damage and Cognitive Decline. <i>Stroke</i> , 2015, 46, 16-22.	1.0	35
39	CDH13 Genotype-Dependent Association of High-Molecular Weight Adiponectin With All-Cause Mortality: The J-SHIPP Study. <i>Diabetes Care</i> , 2014, 37, 396-401.	4.3	22
40	Change in arterial stiffness associated with monthly bisphosphonate treatment in women with postmenopausal osteoporosis. <i>Menopause</i> , 2014, 21, 962-966.	0.8	12
41	Mechanical Stresses, Arterial Stiffness, and Brain Small Vessel Diseases. <i>Stroke</i> , 2014, 45, 3287-3292.	1.0	16
42	Sarcopenic obesity in aging population: current status and future directions for research. <i>Endocrine</i> , 2014, 45, 15-25.	1.1	144
43	Effect of weight loss on central systolic blood pressure in elderly community-dwelling persons. <i>Hypertension Research</i> , 2014, 37, 933-938.	1.5	18
44	The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2014). <i>Hypertension Research</i> , 2014, 37, 253-253.	1.5	962
45	Sarcopenic obesity and arterial stiffness, pressure wave reflection and central pulse pressure: The J-SHIPP study. <i>International Journal of Cardiology</i> , 2014, 174, 214-217.	0.8	23
46	Clinical characteristics of high plasma adiponectin and high plasma leptin as risk factors for arterial stiffness and related end-organ damage. <i>Atherosclerosis</i> , 2014, 235, 424-429.	0.4	18
47	Muscle Mass, Visceral Fat, and Plasma Levels of B-Type Natriuretic Peptide in Healthy Individuals (from) <i>Tj ETQq1 1 0,784314,rgBT /Over</i>	0.7	16
48	Postprandial hypotension as a risk marker for asymptomatic lacunar infarction. <i>Journal of Hypertension</i> , 2014, 32, 1084-1090.	0.3	44
49	Usefulness of combining serum uric acid and high-sensitivity C-reactive protein for risk stratification of patients with metabolic syndrome in community-dwelling women. <i>Endocrine</i> , 2013, 44, 132-139.	1.1	23
50	Hematological parameters are associated with metabolic syndrome in Japanese community-dwelling persons. <i>Endocrine</i> , 2013, 43, 334-341.	1.1	35
51	Genetic Variation in <i>CDH13</i> Is Associated With Lower Plasma Adiponectin Levels but Greater Adiponectin Sensitivity in East Asian Populations. <i>Diabetes</i> , 2013, 62, 4277-4283.	0.3	48
52	Deep cerebral microbleeds are negatively associated with HDL-C in elderly first-time ischemic stroke patients. <i>Journal of the Neurological Sciences</i> , 2013, 325, 137-141.	0.3	6
53	Association of Longer QT Interval With Arterial Waveform and Lower Pulse Pressure Amplification: The Nagahama Study. <i>American Journal of Hypertension</i> , 2013, 26, 973-980.	1.0	6
54	Positive natural selection of <i>TRIB2</i> , a novel gene that influences visceral fat accumulation, in East Asia. <i>Human Genetics</i> , 2013, 132, 201-217.	1.8	19

#	ARTICLE	IF	CITATIONS
55	A slightly high-normal glucose level is associated with increased arterial stiffness in Japanese community-dwelling persons with pre-diabetes. <i>Vascular Medicine</i> , 2013, 18, 251-256.	0.8	8
56	An association between body mass index and high-sensitivity C-reactive protein concentrations is influenced by age in community-dwelling persons. <i>Annals of Clinical Biochemistry</i> , 2013, 50, 457-464.	0.8	23
57	Association of hematological parameters with insulin resistance, insulin sensitivity, and asymptomatic cerebrovascular damage: The J-SHIP and Toon Health Study. <i>Clinical Hemorheology and Microcirculation</i> , 2013, 55, 297-311.	0.9	7
58	Plasma Resistin Is Associated With Single Nucleotide Polymorphisms of a Possible Resistin Receptor, the Decorin Gene, in the General Japanese Population. <i>Diabetes</i> , 2013, 62, 649-652.	0.3	24
59	Visit-To-Visit Variability in Systolic Blood Pressure Is a Novel Risk Factor for the Growth of Intracranial Aneurysms. <i>Cerebrovascular Diseases</i> , 2013, 36, 401-406.	0.8	18
60	Synergistic influence of age and serum uric acid on blood pressure among community-dwelling Japanese women. <i>Hypertension Research</i> , 2013, 36, 634-638.	1.5	13
61	A Slightly Low Hemoglobin Level Is Beneficially Associated with Arterial Stiffness in Japanese Community-Dwelling Women. <i>Clinical and Experimental Hypertension</i> , 2012, 34, 92-98.	0.5	42
62	Î³-Glutamyl Transferase and High-Molecular-Weight Adiponectin Levels Are Synergistically Associated with Metabolic Syndrome and Insulin Resistance in Community-Dwelling Persons. <i>Metabolic Syndrome and Related Disorders</i> , 2012, 10, 83-91.	0.5	6
63	Mice Lacking Hypertension Candidate Gene ATP2B1 in Vascular Smooth Muscle Cells Show Significant Blood Pressure Elevation. <i>Hypertension</i> , 2012, 59, 854-860.	1.3	79
64	Central Blood Pressure and End-organ Damage. <i>Current Hypertension Reviews</i> , 2012, 8, 100-107.	0.5	4
65	Low-dose rosuvastatin improves the functional and morphological markers of atherosclerosis in asymptomatic postmenopausal women with dyslipidemia. <i>Menopause</i> , 2012, 19, 1294-1299.	0.8	19
66	Postprandial hypertension, an overlooked risk marker for arteriosclerosis. <i>Atherosclerosis</i> , 2012, 224, 500-505.	0.4	20
67	Plasma Resistin Levels Are Associated with Insulin Resistance in Older Japanese Men from a Rural Village. <i>Metabolic Syndrome and Related Disorders</i> , 2012, 10, 380-386.	0.5	3
68	Arterial stiffness in sarcopenic visceral obesity in the elderly: J-SHIP study. <i>International Journal of Cardiology</i> , 2012, 158, 146-148.	0.8	54
69	Alanine aminotransferase/aspartate aminotransferase ratio is the best surrogate marker for insulin resistance in non-obese Japanese adults. <i>Cardiovascular Diabetology</i> , 2012, 11, 117.	2.7	60
70	Atherosclerotic Indices for the Prediction of Cognitive Impairment in a Middle-aged to Elderly General Population: Shimanami Health Promoting Program Study. <i>Journal of the American Geriatrics Society</i> , 2012, 60, 1996-1997.	1.3	1
71	Hunting for genes for hypertension: the Millennium Genome Project for Hypertension. <i>Hypertension Research</i> , 2012, 35, 567-573.	1.5	14
72	Potential Utility of Soluble p3-Aldoactin Plasma Levels as a Biomarker for Sporadic Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2012, 31, 421-428.	1.2	11

#	ARTICLE	IF	CITATIONS
73	Perceived age of facial features is a significant diagnosis criterion for age-related carotid atherosclerosis in Japanese subjects: J-SHIPP study. <i>Geriatrics and Gerontology International</i> , 2012, 12, 733-740.	0.7	22
74	Leptin in Sarcopenic Visceral Obesity: Possible Link between Adipocytes and Myocytes. <i>PLoS ONE</i> , 2011, 6, e24633.	1.1	74
75	Relatively lower central aortic pressure in patients with impaired insulin sensitivity and resistance. <i>Journal of Hypertension</i> , 2011, 29, 1948-1954.	0.3	21
76	Genotype risk score of common susceptible variants for prediction of type 2 diabetes mellitus in Japanese: the Shimanami Health Promoting Program (J-SHIPP study). <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 1634-1640.	1.5	25
77	A single-nucleotide polymorphism in the human THADA gene is associated with circulating resistin in the general Japanese population. <i>Diabetology International</i> , 2011, 2, 190-196.	0.7	2
78	Association between fasting plasma glucose and high-sensitivity C-reactive protein: gender differences in a Japanese community-dwelling population. <i>Cardiovascular Diabetology</i> , 2011, 10, 51.	2.7	19
79	Serum Gamma-Glutamyl Transferase within Its Normal Concentration Range Is Related to the Presence of Impaired Fasting Glucose and Diabetes among Japanese Community-Dwelling Persons. <i>Endocrine Research</i> , 2011, 36, 64-73.	0.6	10
80	Serum High Molecular Weight Adiponectin Correlates with Arterial Stiffness in Community-Dwelling Persons. <i>Endocrine Research</i> , 2011, 36, 53-63.	0.6	10
81	Association of Serum High Molecular Weight Adeponectin and Blood Pressure among Non-Diabetic Community-Dwelling Men. <i>Clinical and Experimental Hypertension</i> , 2011, 33, 336-344.	0.5	7
82	Hemoglobin is Associated with Serum High Molecular Weight Adiponectin in Japanese Community-Dwelling Persons. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011, 18, 182-189.	0.9	17
83	Associations between short one-leg standing time and speed of sound of calcaneal bone in a general population: The Shimanami Health Promoting Program (J-SHIPP) study. <i>Geriatrics and Gerontology International</i> , 2010, 10, 138-144.	0.7	7
84	High-sensitivity c-reactive protein and gamma-glutamyl transferase levels are synergistically associated with metabolic syndrome in community-dwelling persons. <i>Cardiovascular Diabetology</i> , 2010, 9, 87.	2.7	33
85	Quadriceps sarcopenia and visceral obesity are risk factors for postural instability in the middle-aged to elderly population. <i>Geriatrics and Gerontology International</i> , 2010, 10, 233-243.	0.7	70
86	Insulin Resistance and Prevalence of Prehypertension and Hypertension Among Community-Dwelling Persons. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010, 17, 148-155.	0.9	36
87	Smoking Status is Associated with Serum High Molecular Adiponectin Levels in Community-Dwelling Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010, 17, 423-430.	0.9	30
88	A at Single Nucleotide Polymorphism-358 Is Required for G at -420 to Confer the Highest Plasma Resistin in the General Japanese Population. <i>PLoS ONE</i> , 2010, 5, e9718.	1.1	37
89	Common Variants in the ATP2B1 Gene Are Associated With Susceptibility to Hypertension. <i>Hypertension</i> , 2010, 56, 973-980.	1.3	96
90	Abdominal Fat, Adipose-Derived Hormones and Mild Cognitive Impairment: The J-SHIPP Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2010, 30, 432-439.	0.7	56

#	ARTICLE	IF	CITATIONS
91	Cross-Sectional Characterization of all Classes of Antihypertensives in Terms of Central Blood Pressure in Japanese Hypertensive Patients. <i>American Journal of Hypertension</i> , 2010, 23, 260-268.	1.0	49
92	Association of Central Systolic Blood Pressure With Intracerebral Small Vessel Disease in Japanese. <i>American Journal of Hypertension</i> , 2010, 23, 889-894.	1.0	53
93	Accumulation of common polymorphisms is associated with development of hypertension: a 12-year follow-up from the Ohasama study. <i>Hypertension Research</i> , 2010, 33, 129-134.	1.5	37
94	Association of Four Genetic Loci with Uric Acid Levels and Reduced Renal Function: The J-SHIPP Suita Study. <i>American Journal of Nephrology</i> , 2010, 32, 279-286.	1.4	35
95	Abnormal nocturnal blood pressure profile is associated with mild cognitive impairment in the elderly: the J-SHIPP study. <i>Hypertension Research</i> , 2010, 33, 32-36.	1.5	59
96	Arterial stiffness is associated with low thigh muscle mass in middle-aged to elderly men. <i>Atherosclerosis</i> , 2010, 212, 327-332.	0.4	225
97	Postural Instability Is Associated with Brain Atrophy and Cognitive Impairment in the Elderly: The J-SHIPP Study. <i>Dementia and Geriatric Cognitive Disorders</i> , 2010, 29, 379-387.	0.7	61
98	The GCKR rs780094 polymorphism is associated with susceptibility of type 2 diabetes, reduced fasting plasma glucose levels, increased triglycerides levels and lower HOMA-IR in Japanese population. <i>Journal of Human Genetics</i> , 2010, 55, 600-604.	1.1	57
99	Alcohol Drinking Status is Associated with Serum High Molecular Weight Adiponectin in Community-Dwelling Japanese Men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010, 17, 953-962.	0.9	17
100	Serum High Molecular Weight Adiponectin is Associated with Mild Renal Dysfunction in Japanese Adults. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010, 17, 1141-1148.	0.9	10
101	Serum Gamma-Glutamyl Transferase Levels are Associated with Metabolic Syndrome in Community-Dwelling Individuals. <i>Journal of Atherosclerosis and Thrombosis</i> , 2009, 16, 355-362.	0.9	32
102	Replication Study of Candidate Genes Associated With Type 2 Diabetes Based On Genome-Wide Screening. <i>Diabetes</i> , 2009, 58, 493-498.	0.3	136
103	Central blood pressure, arterial stiffness and the heart in hypertensive patients. <i>Hypertension Research</i> , 2009, 32, 1056-1058.	1.5	7
104	Silent cerebral microbleeds associated with arterial stiffness in an apparently healthy subject. <i>Hypertension Research</i> , 2009, 32, 255-260.	1.5	55
105	Association of monocyte chemoattractant protein 1 gene polymorphism with susceptibility to nonfamilial idiopathic dilated cardiomyopathy. <i>Journal of Cardiology</i> , 2009, 54, 66-70.	0.8	6
106	PPAR β Pro12Ala Pro/Pro and resistin SNP β 420 G/G genotypes are synergistically associated with plasma resistin in the Japanese general population. <i>Clinical Endocrinology</i> , 2009, 71, 341-345.	1.2	9
107	Composition of lower extremity in relation to a high ankle-brachial index. <i>Journal of Hypertension</i> , 2009, 27, 167-173.	0.3	31
108	Alcohol Consumption Is Associated with Decreased Insulin Resistance Independent of Body Mass Index in Japanese Community-Dwelling Men. <i>Tohoku Journal of Experimental Medicine</i> , 2009, 218, 331-337.	0.5	21

#	ARTICLE	IF	CITATIONS
109	Asymptomatic Cerebral Microbleeds Seen in Healthy Subjects Have a Strong Association With Asymptomatic Lacunar Infarction. <i>Circulation Journal</i> , 2009, 73, 530-533.	0.7	39
110	No Association Between <i>INSIG2</i> Gene rs7566605 Polymorphism and Being Overweight in Japanese Population. <i>Obesity</i> , 2008, 16, 211-215.	1.5	21
111	An Association between Body Mass Index and Estimated Glomerular Filtration Rate. <i>Hypertension Research</i> , 2008, 31, 1559-1564.	1.5	45
112	Effects of Obesity and Smoking on Mental Stress-Induced Blood Pressure and Augmentation Index Responses in Normotensive Young Males: The J-SHIPP Study. <i>Hypertension Research</i> , 2008, 31, 1219-1224.	1.5	20
113	Reduced High-Molecular-Weight Adiponectin and Elevated High-Sensitivity C-Reactive Protein Are Synergistic Risk Factors for Metabolic Syndrome in a Large-Scale Middle-Aged to Elderly Population: the Shimanami Health Promoting Program Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 715-722.	1.8	50
114	Identification of Hypertension-Susceptibility Genes and Pathways by a Systemic Multiple Candidate Gene Approach: The Millennium Genome Project for Hypertension. <i>Hypertension Research</i> , 2008, 31, 203-212.	1.5	57
115	Blood Pressure Is the Main Determinant of the Reflection Wave in Patients with Type 2 Diabetes. <i>Hypertension Research</i> , 2008, 31, 493-499.	1.5	17
116	High Prevalence of Prehypertension Is Associated with the Increased Body Mass Index in Community-Dwelling Japanese. <i>Tohoku Journal of Experimental Medicine</i> , 2008, 216, 353-361.	0.5	22
117	An Association between Decreased Estimated Glomerular Filtration Rate and Arterial Stiffness. <i>Internal Medicine</i> , 2008, 47, 593-598.	0.3	55
118	An Association between Metabolic Syndrome and the Estimated Glomerular Filtration Rate. <i>Internal Medicine</i> , 2008, 47, 1399-1406.	0.3	29
119	Association between Serum Gamma-Glutamyl Transferase Level and Prehypertension among Community-Dwelling Men. <i>Tohoku Journal of Experimental Medicine</i> , 2008, 216, 213-221.	0.5	16
120	Plasma Resistin, Associated With Single Nucleotide Polymorphism -420, Is Correlated With Insulin Resistance, Lower HDL Cholesterol, and High-Sensitivity C-Reactive Protein in the Japanese General Population. <i>Diabetes Care</i> , 2007, 30, 1501-1506.	4.3	129
121	Effect of Acute and Long-Term Aerobic Exercise on Arterial Stiffness in the Elderly. <i>Hypertension Research</i> , 2007, 30, 895-902.	1.5	51
122	Migraine Is Associated with Enhanced Arterial Stiffness. <i>Hypertension Research</i> , 2007, 30, 577-583.	1.5	26
123	High-density association study and nomination of susceptibility genes for hypertension in the Japanese National Project. <i>Human Molecular Genetics</i> , 2007, 17, 617-627.	1.4	53
124	Case of pneumococcal meningoenkephalitis with periodic synchronous discharge patterns on electroencephalogram. <i>Geriatrics and Gerontology International</i> , 2007, 7, 401-405.	0.7	1
125	Interaction between Serotonin 2A Receptor and Endothelin-1 Variants in Association with Hypertension in Japanese. <i>Hypertension Research</i> , 2006, 29, 227-232.	1.5	18
126	Association of Dopamine .BETA.-Hydroxylase Polymorphism with Hypertension through Interaction with Fasting Plasma Glucose in Japanese. <i>Hypertension Research</i> , 2005, 28, 215-221.	1.5	10

#	ARTICLE	IF	CITATIONS
127	Hypotension Associated with Prone Body Position: A Possible Overlooked Postural Hypotension. Hypertension Research, 2005, 28, 741-746.	1.5	34
128	DELAYED REFLECTION PRESSURE WAVE: NEW ASPECTS FOR POSTPRANDIAL HYPOTENSION. Journal of the American Geriatrics Society, 2005, 53, 1833-1834.	1.3	1
129	Effect of time standing up on orthostatic blood pressure change in the elderly: The J-SHIP Study. Geriatrics and Gerontology International, 2005, 5, 254-258.	0.7	1
130	An association of 5,10-Methylenetetrahydrofolate Reductase (MTHFR) gene polymorphism and ischemic stroke. Journal of Stroke and Cerebrovascular Diseases, 2005, 14, 67-74.	0.7	30
131	Radial augmentation index: A useful and easily obtainable parameter for vascular aging. American Journal of Hypertension, 2005, 18, 11-14.	1.0	220
132	Orthostatic Systolic Hypotension and the Reflection Pressure Wave. Hypertension Research, 2005, 28, 537-543.	1.5	28
133	Home Blood Pressure Is a Better Predictor of Carotid Atherosclerosis than Office Blood Pressure in Community-Dwelling Subjects. Hypertension Research, 2004, 27, 633-639.	1.5	52
134	Genetic Predisposition to Neurological Symptoms in Lacunar Infarction. Cerebrovascular Diseases, 2004, 17, 273-279.	0.8	27
135	Microalbuminuria and Arterial Stiffness in a General Population: the Shimanami Health Promoting Program (J-SHIP) Study. Hypertension Research, 2004, 27, 471-477.	1.5	85
136	Comprehensive studies of carotid atherosclerosis. Geriatrics and Gerontology International, 2003, 3, 12-14.	0.7	0
137	Association of Endothelin-1 Gene Variant With Hypertension. Hypertension, 2003, 41, 163-167.	1.3	102
138	MTHFR Gene Polymorphism as a Risk Factor for Silent Brain Infarcts and White Matter Lesions in the Japanese General Population. Stroke, 2003, 34, 1130-1135.	1.0	103
139	Association of Angiotensin II Type 2 Receptor Gene Variant with Hypertension. Hypertension Research, 2003, 26, 547-552.	1.5	50
140	Association of the GNAS1 Gene Variant with Hypertension Is Dependent on Alcohol Consumption.. Hypertension Research, 2003, 26, 439-444.	1.5	27
141	Polymorphism of the Monocyte Chemoattractant Protein (MCP-1) Gene Is Associated with the Plasma Level of MCP-1 But Not with Carotid Intima-Media Thickness. Hypertension Research, 2003, 26, 677-683.	1.5	62
142	Genome-Wide Linkage Disequilibrium Mapping of Hypertension in Japan. Hypertension Research, 2003, 26, 533-540.	1.5	10
143	Soluble Fas ligand and atherosclerosis in hypertensive patients. Journal of Hypertension, 2002, 20, 895-898.	0.3	34
144	Polymorphisms of genes encoding components of the sympathetic nervous system but not the renin-angiotensin system as risk factors for orthostatic hypotension. Journal of Hypertension, 2002, 20, 651-656.	0.3	39

#	ARTICLE	IF	CITATIONS
145	Questionnaire Survey on the Japanese Guidelines for Treatment of Hypertension in the Elderly: 1999 Revised Version. <i>Hypertension Research</i> , 2002, 25, 69-75.	1.5	10
146	An Interaction between Systolic Blood Pressure and Angiotensin-Converting Enzyme Gene Polymorphism on Carotid Atherosclerosis.. <i>Hypertension Research</i> , 2002, 25, 875-880.	1.5	19
147	Chlamydia pneumoniae Seropositivity Is Associated With Increased Plasma Levels of Soluble Cellular Adhesion Molecules in Community-Dwelling Subjects. <i>Stroke</i> , 2002, 33, 1474-1479.	1.0	18
148	Supine hypertension with transient papilledema. <i>Geriatrics and Gerontology International</i> , 2002, 2, 148-152.	0.7	0
149	Genotype-Specific Association between Circulating Soluble Cellular Adhesion Molecules and Carotid Intima-Media Thickness in Community Residents: J-SHIP Study. <i>Hypertension Research</i> , 2002, 25, 31-39.	1.5	16
150	Plasma Hepatocyte Growth Factor and the Relationship between Risk Factors and Carotid Atherosclerosis.. <i>Hypertension Research</i> , 2002, 25, 661-667.	1.5	17
151	Risk factor-gene interaction in carotid atherosclerosis: effect of genes encoding the renin-angiotensin system. <i>American Journal of Hypertension</i> , 2001, 14, A77.	1.0	0
152	Association between carotid arterial remodeling and plasma concentration of circulating hepatocyte growth factor. <i>Journal of Hypertension</i> , 2001, 19, 1975-1979.	0.3	25
153	CAROTID WALL SHEAR STRESS AND AGING; <i>To the Editor</i>.: <i>Journal of the American Geriatrics Society</i> , 2000, 48, 1349-1350.	1.3	3
154	ORTHOSTATIC HYPERTENSION: ANOTHER ORTHOSTATIC DISORDER TO BE AWARE OF. <i>Journal of the American Geriatrics Society</i> , 2000, 48, 1538-1539.	1.3	19
155	Association Between Risk Factors for Atherosclerosis and Mechanical Forces in Carotid Artery. <i>Stroke</i> , 2000, 31, 2319-2324.	1.0	144
156	Deletion Polymorphism of ACE Gene Is Associated with Higher Blood Pressure after Hospitalization in Normotensive Subjects.. <i>Hypertension Research</i> , 2000, 23, 201-205.	1.5	14
157	Low Wall Shear Stress Contributes to Atherosclerosis of the Carotid Artery in Hypertensive Patients.. <i>Hypertension Research</i> , 1999, 22, 203-207.	1.5	46
158	Guidelines for Hypertension in the Elderly. 1999 Revised Version.. <i>Hypertension Research</i> , 1999, 22, 231-259.	1.5	23
159	Postprandial Hypotension Is Associated With Asymptomatic Cerebrovascular Damage in Essential Hypertensive Patients. <i>Hypertension</i> , 1999, 33, 565-568.	1.3	111
160	Relation of left ventricular hypertrophy and geometry to asymptomatic cerebrovascular damage in essential hypertension. <i>American Journal of Cardiology</i> , 1999, 83, 367-370.	0.7	63
161	CONTRIBUTION OF REFLECTION OF PRESSURE WAVE ON CENTRAL SYSTOLIC BLOOD PRESSURE IN OLDER HYPERTENSIVE PATIENTS. <i>Journal of the American Geriatrics Society</i> , 1999, 47, 499-499.	1.3	5
162	Postprandial hypotension: evaluation by ambulatory blood pressure monitoring. <i>American Journal of Hypertension</i> , 1998, 11, 1358-1363.	1.0	39

#	ARTICLE	IF	CITATIONS
163	Molecular structure and function of rat platelet-derived growth factor β^2 -receptor gene promoter. Journal of Hypertension, 1998, 16, 437-445.	0.3	7
164	Molecular Structure and Function of Rat CCAAT-Enhancer Binding Protein-Delta Gene Promoter. Biochemical and Biophysical Research Communications, 1997, 231, 30-36.	1.0	12
165	Left Ventricular Hypertrophy Precedes Other Target-Organ Damage in Primary Aldosteronism. Hypertension, 1997, 29, 723-727.	1.3	81
166	Autonomic Nervous Function in Non-dipper Essential Hypertensive Subjects. Hypertension, 1995, 26, 808-814.	1.3	217
167	Angiotensin(1â^7) in the spontaneously hypertensive rat. Peptides, 1993, 14, 883-891.	1.2	141
168	Peripheral vs. Central Blockade of the Renin-Angiotensin System in Spontaneously Hypertensive Rats: Comparison of Novel AT1 Receptor Antagonist TCV-116 with Angiotensin Converting Enzyme Inhibitor Delapril.. Hypertension Research, 1993, 16, 239-246.	1.5	22
169	Reassessment of plasma angiotensins measurement: Effects of protease inhibitors and sample handling procedures. Peptides, 1991, 12, 1135-1141.	1.2	83