
Valerie M Corrigall

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11860733/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The Human Endoplasmic Reticulum Molecular Chaperone BiP Is an Autoantigen for Rheumatoid Arthritis and Prevents the Induction of Experimental Arthritis. Journal of Immunology, 2001, 166, 1492-1498.	0.8	171
2	PATHOGENESIS OF RHEUMATOID ARTHRITIS. Rheumatic Disease Clinics of North America, 2001, 27, 317-334.	1.9	96
3	Inhibition of antigen-presenting cell function and stimulation of human peripheral blood mononuclear cells to express an antiinflammatory cytokine profile by the stress protein BiP: Relevance to the treatment of inflammatory arthritis. Arthritis and Rheumatism, 2004, 50, 1164-1171.	6.7	93
4	Treatment of murine collagen-induced arthritis by the stress protein BiP via interleukin-4–producing regulatory T cells: A novel function for an ancient protein. Arthritis and Rheumatism, 2006, 54, 854-863.	6.7	75
5	Evidence for the continuous recruitment and activation of T cells into the joints of patients with rheumatoid arthritis. European Journal of Immunology, 1994, 24, 2706-2713.	2.9	72
6	Glucose-regulated protein 78 (Grp78/BiP) is secreted by human oviduct epithelial cells and the recombinant protein modulates sperm–zona pellucida binding. Fertility and Sterility, 2010, 93, 1574-1584.	1.0	65
7	Stress cytokines: pivotal proteins in immune regulatory networks. Current Opinion in Immunology, 2004, 16, 531-534.	5.5	60
8	Autoantigens and immune pathways in rheumatoid arthritis. Critical Reviews in Immunology, 2002, 22, 281-93.	0.5	44
9	Binding immunoglobulin proteinâ€treated peripheral blood monocyteâ€derived dendritic cells are refractory to maturation and induce regulatory Tâ€cell development. Immunology, 2009, 128, 218-226.	4.4	40
10	Safety and patient response as indicated by biomarker changes to binding immunoglobulin protein in the phase I/IIA RAGULA clinical trial in rheumatoid arthritis. Rheumatology, 2016, 55, 1993-2000.	1.9	36
11	Immunoglobulin heavy-chain-binding protein (BiP): a stress protein that has the potential to be a novel therapy for rheumatoid arthritis. Biochemical Society Transactions, 2014, 42, 1752-1755.	3.4	34
12	Lack of CD80 expression by fibroblast-like synoviocytes leading to anergy in T lymphocytes. Arthritis and Rheumatism, 2000, 43, 1606-1615.	6.7	33
13	Pro-resolution immunological networks: binding immunoglobulin protein and other resolution-associated molecular patterns. Rheumatology, 2012, 51, 780-788.	1.9	25
14	A New-Age for Biologic Therapies: Long-Term Drug-Free Therapy with BiP?. Frontiers in Immunology, 2012, 3, 17.	4.8	24
15	Binding immunoglobulin protein resolves rheumatoid synovitis: a xenogeneic study using rheumatoid arthritis synovial membrane transplants in SCID mice. Arthritis Research and Therapy, 2011, 13, R149.	3.5	19
16	BiP, An Antiâ€Inflammatory ER Protein, is a Potential New Therapy for the Treatment of Rheumatoid Arthritis. Novartis Foundation Symposium, 2008, 291, 212-220.	1.1	15
17	Binding Immunoglobulin Protein (<scp>BIP</scp>) Inhibits <scp>TNF</scp> â€Î±â€"Induced Osteoclast Differentiation and Systemic Bone Loss in an Erosive Arthritis Model. ACR Open Rheumatology, 2019, 1, 382-393.	2.1	10

BiP, a Negative Regulator Involved in Rheumatoid Arthritis. , 2005, , 234-248.

2

IF

ARTICLE #