
Ans Van Ginckel

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1188646/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Greater magnitude tibiofemoral contact forces are associated with reduced prevalence of osteochondral pathologies 2–3Âyears following anterior cruciate ligament reconstruction. Knee Surgery, Sports Traumatology, Arthroscopy, 2019, 27, 707-715.	4.2	16
2	Diet-induced weight loss alone or combined with exercise in overweight or obese people with knee osteoarthritis: A systematic review and meta-analysis. Seminars in Arthritis and Rheumatism, 2019, 48, 765-777.	3.4	71
3	Effects of long-term exercise therapy on knee joint structure in people with knee osteoarthritis: A systematic review and meta-analysis. Seminars in Arthritis and Rheumatism, 2019, 48, 941-949.	3.4	29
4	Frontal plane hip joint loading according to pain severity in people with hip osteoarthritis. Journal of Orthopaedic Research, 2018, 36, 1637-1644.	2.3	8
5	Factors Influencing Cane Use for the Management of Knee Osteoarthritis: A Crossâ€Sectional Survey. Arthritis Care and Research, 2018, 70, 1455-1460.	3.4	4
6	Presence of comorbidities and prognosis of clinical symptoms in knee and/or hip osteoarthritis: A systematic review and meta-analysis. Seminars in Arthritis and Rheumatism, 2018, 47, 805-813.	3.4	124
7	The effects of aerobic exercise on eGFR, blood pressure and VO2peak in patients with chronic kidney disease stages 3-4: A systematic review and meta-analysis. PLoS ONE, 2018, 13, e0203662.	2.5	59
8	Cartilage morphology at 2–3Âyears following anterior cruciate ligament reconstruction with or without concomitant meniscal pathology. Knee Surgery, Sports Traumatology, Arthroscopy, 2017, 25, 426-436.	4.2	20
9	Interventions to increase adherence to therapeutic exercise in older adults with low back pain and/or hip/knee osteoarthritis: a systematic review and meta-analysis. British Journal of Sports Medicine, 2017, 51, 791-799.	6.7	130
10	Impact of Cane Use on Bone Marrow Lesion Volume in People With Medial Knee Osteoarthritis (CUBA) Tj ETQq	0 0 0 rgBT 2.4	/Overlock 10 4
11	Hip Strength as an Intrinsic Risk Factor for Lateral Ankle Sprains in Youth Soccer Players: A 3-Season Prospective Study. American Journal of Sports Medicine, 2017, 45, 410-416.	4.2	59
12	In vivo deformation of thin cartilage layers: Feasibility and applicability of T2* mapping. Journal of Orthopaedic Research, 2016, 34, 771-778.	2.3	6
13	T2* mapping of subtalar cartilage: Precision and association between anatomical variants and cartilage composition. Journal of Orthopaedic Research, 2016, 34, 1969-1976.	2.3	12
14	Patient Knowledge and Beliefs About Knee Osteoarthritis After Anterior Cruciate Ligament Injury and Reconstruction. Arthritis Care and Research, 2016, 68, 1180-1185.	3.4	13
15	Cartilage Status in Relation to Return to Sports After Anterior Cruciate Ligament Reconstruction. American Journal of Sports Medicine, 2013, 41, 550-559.	4.2	63
16	Acute Cartilage Loading Responses After an In Vivo Squatting Exercise in People With Doubtful to Mild Knee Osteoarthritis: A Case-Control Study. Physical Therapy, 2013, 93, 1049-1060.	2.4	12

17	Human ankle cartilage deformation after different in vivo impact conditions. Knee Surgery, Sports Traumatology, Arthroscopy, 2011, 19, 137-143.	4.2	15

18Intrinsic gait-related risk factors for Achilles tendinopathy in novice runners: A prospective study.1.4109Cait and Posture, 2009, 29, 387-391.1.4