

# Annette Becker

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11892470/publications.pdf>

Version: 2024-02-01

12  
papers

5,168  
citations

758635

12  
h-index

1199166

12  
g-index

12  
all docs

12  
docs citations

12  
times ranked

9734  
citing authors

#	ARTICLE	IF	CITATIONS
1	Pancreatic cancer exosomes initiate pre-metastatic niche formation in the liver. <i>Nature Cell Biology</i> , 2015, 17, 816-826.	4.6	2,064
2	Extracellular Vesicles in Cancer: Cell-to-Cell Mediators of Metastasis. <i>Cancer Cell</i> , 2016, 30, 836-848.	7.7	1,401
3	Double-stranded DNA in exosomes: a novel biomarker in cancer detection. <i>Cell Research</i> , 2014, 24, 766-769.	5.7	1,282
4	MeCP2 interacts with HP1 and modulates its heterochromatin association during myogenic differentiation. <i>Nucleic Acids Research</i> , 2007, 35, 5402-5408.	6.5	137
5	MeCP2 Rett mutations affect large scale chromatin organization. <i>Human Molecular Genetics</i> , 2011, 20, 4187-4195.	1.4	72
6	Targeted manipulation of heterochromatin rescues MeCP2 Rett mutants and re-establishes higher order chromatin organization. <i>Nucleic Acids Research</i> , 2012, 40, e176-e176.	6.5	44
7	Involvement of p53 in the cytotoxic activity of the NAMPT inhibitor FK866 in myeloid leukemic cells. <i>International Journal of Cancer</i> , 2013, 132, 766-774.	2.3	40
8	Poly(ADP-ribosyl)ation of Methyl CpG Binding Domain Protein 2 Regulates Chromatin Structure. <i>Journal of Biological Chemistry</i> , 2016, 291, 4873-4881.	1.6	28
9	Direct Homo- and Hetero-Interactions of MeCP2 and MBD2. <i>PLoS ONE</i> , 2013, 8, e53730.	1.1	28
10	Inhibition of NAMPT pathway by FK866 activates the function of p53 in HEK293T cells. <i>Biochemical and Biophysical Research Communications</i> , 2012, 424, 371-377.	1.0	27
11	Processive DNA synthesis is associated with localized decompaction of constitutive heterochromatin at the sites of DNA replication and repair. <i>Nucleus</i> , 2019, 10, 231-253.	0.6	25
12	Generation and Characterization of Rat and Mouse Monoclonal Antibodies Specific for MeCP2 and Their Use in X-Inactivation Studies. <i>PLoS ONE</i> , 2011, 6, e26499.	1.1	20