Shi Yue

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11901057/publications.pdf

Version: 2024-02-01

		471509	839539
18	1,107	17	18
papers	citations	h-index	g-index
18	18	18	1708
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	KEAP1-NRF2 complex in ischemia-induced hepatocellular damage of mouse liver transplants. Journal of Hepatology, 2013, 59, 1200-1207.	3.7	132
2	Sphingosine kinase/sphingosine 1-phosphate (S1P)/S1P receptor axis is involved in liver fibrosis-associated angiogenesis. Journal of Hepatology, 2013, 59, 114-123.	3.7	102
3	Involvement of Sphingosine 1-Phosphate (SIP)/S1P3 Signaling in Cholestasis-Induced Liver Fibrosis. American Journal of Pathology, 2009, 175, 1464-1472.	3.8	97
4	The myeloid heat shock transcription factor 1/βâ€catenin axis regulates NLR family, pyrin domainâ€containing 3 inflammasome activation in mouse liver ischemia/reperfusion injury. Hepatology, 2016, 64, 1683-1698.	7.3	84
5	Myeloid PTEN Deficiency Protects Livers from Ischemia Reperfusion Injury by Facilitating M2 Macrophage Differentiation. Journal of Immunology, 2014, 192, 5343-5353.	0.8	74
6	Glycogen synthase kinase $3\hat{l}^2$ promotes liver innate immune activation by restraining AMP-activated protein kinase activation. Journal of Hepatology, 2018, 69, 99-109.	3.7	64
7	\hat{l}^2 -catenin regulates innate and adaptive immunity in mouse liver ischemia-reperfusion injury. Hepatology, 2013, 57, 1203-1214.	7.3	60
8	Heme oxygenase-1 regulates sirtuin-1–autophagy pathway in liver transplantation: From mouse to human. American Journal of Transplantation, 2018, 18, 1110-1121.	4.7	60
9	Prolonged Ischemia Triggers Necrotic Depletion of Tissue-Resident Macrophages To Facilitate Inflammatory Immune Activation in Liver Ischemia Reperfusion Injury. Journal of Immunology, 2017, 198, 3588-3595.	0.8	58
10	Essential roles of sphingosine 1â€phosphate receptor types 1 and 3 in human hepatic stellate cells motility and activation. Journal of Cellular Physiology, 2011, 226, 2370-2377.	4.1	56
11	Rapamycin Protection of Livers From Ischemia and Reperfusion Injury Is Dependent on Both Autophagy Induction and Mammalian Target of Rapamycin Complex 2-Akt Activation. Transplantation, 2015, 99, 48-55.	1.0	53
12	Myeloid Notch1 deficiency activates the RhoA/ROCK pathway and aggravates hepatocellular damage in mouse ischemic livers. Hepatology, 2018, 67, 1041-1055.	7.3	52
13	15-deoxy-Î" ^{12,14} -prostaglandin J ₂ reduces recruitment of bone marrow-derived monocyte/macrophages in chronic liver injury in mice. Hepatology, 2012, 56, 350-360.	7.3	48
14	Adoptive Transfer of Heme Oxygenase-1 (HO-1)-Modified Macrophages Rescues the Nuclear Factor Erythroid 2-Related Factor (Nrf2) Antiinflammatory Phenotype in Liver Ischemia/Reperfusion Injury. Molecular Medicine, 2014, 20, 448-455.	4.4	45
15	The Dichotomy of Endoplasmic Reticulum Stress Response in Liver Ischemia-Reperfusion Injury. Transplantation, 2016, 100, 365-372.	1.0	40
16	Nuclear Factor Erythroid 2–Related Factor 2 Regulates Toll-Like Receptor 4 Innate Responses in Mouse Liver Ischemia-Reperfusion Injury Through Akt-Forkhead box Protein O1 Signaling Network. Transplantation, 2014, 98, 721-728.	1.0	35
17	Blockade of Notch signaling promotes acetaminophen-induced liver injury. Immunologic Research, 2017, 65, 739-749.	2.9	29
18	Phosphatase and tensin homolog‑β atenin signaling modulates regulatory T cells and inflammatory responses in mouse liver ischemia/reperfusion injury. Liver Transplantation, 2017, 23, 813-825.	2.4	18