Martin W G Brinkhof

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11912408/publications.pdf

Version: 2024-02-01

471509 839539 3,358 18 17 18 citations h-index g-index papers 18 18 18 3189 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Mortality of HIV-1-infected patients in the first year of antiretroviral therapy: comparison between low-income and high-income countries. Lancet, The, 2006, 367, 817-824.	13.7	1,030
2	Mortality of Patients Lost to Follow-Up in Antiretroviral Treatment Programmes in Resource-Limited Settings: Systematic Review and Meta-Analysis. PLoS ONE, 2009, 4, e5790.	2.5	409
3	Eearly loss of HIV-infected patients on potent antiretroviral therapy programmes in lower-income countries. Bulletin of the World Health Organization, 2008, 86, 559-567.	3.3	275
4	Gender and the Use of Antiretroviral Treatment in Resource-Constrained Settings: Findings from a Multicenter Collaboration. Journal of Women's Health, 2008, 17, 47-55.	3.3	178
5	Universal Definition of Loss to Follow-Up in HIV Treatment Programs: A Statistical Analysis of 111 Facilities in Africa, Asia, and Latin America. PLoS Medicine, 2011, 8, e1001111.	8.4	167
6	Antiretroviral therapy in resourceâ€limited settings 1996 to 2006: patient characteristics, treatment regimens and monitoring in subâ€Saharan Africa, Asia and Latin America. Tropical Medicine and International Health, 2008, 13, 870-879.	2.3	162
7	Mortality of HIV-Infected Patients Starting Antiretroviral Therapy in Sub-Saharan Africa: Comparison with HIV-Unrelated Mortality. PLoS Medicine, 2009, 6, e1000066.	8.4	161
8	Long-term immunologic response to antiretroviral therapy in low-income countries: a collaborative analysis of prospective studies. Aids, 2008, 22, 2291-2302.	2.2	152
9	Electronic medical record systems, data quality and loss to follow-up: survey of antiretroviral therapy programmes in resource-limited settings. Bulletin of the World Health Organization, 2008, 86, 939-947.	3.3	139
10	Switching to second-line antiretroviral therapy in resource-limited settings: comparison of programmes with and without viral load monitoring. Aids, 2009, 23, 1867-1874.	2.2	136
11	Correcting Mortality for Loss to Follow-Up: A Nomogram Applied to Antiretroviral Treatment Programmes in Sub-Saharan Africa. PLoS Medicine, 2011, 8, e1000390.	8.4	136
12	Public-Health and Individual Approaches to Antiretroviral Therapy: Township South Africa and Switzerland Compared. PLoS Medicine, 2008, 5, e148.	8.4	113
13	Early Mortality and Loss to Follow-up in HIV-Infected Children Starting Antiretroviral Therapy in Southern Africa. Journal of Acquired Immune Deficiency Syndromes (1999), 2010, 54, 524-532.	2.1	88
14	Adjusting Mortality for Loss to Follow-Up: Analysis of Five ART Programmes in Sub-Saharan Africa. PLoS ONE, 2010, 5, e14149.	2.5	85
15	Mortality after failure of antiretroviral therapy in subâ€Saharan Africa. Tropical Medicine and International Health, 2010, 15, 251-258.	2.3	71
16	Derivation of parameters used in Spectrum for eligibility for antiretroviral therapy and survival on antiretroviral therapy. Sexually Transmitted Infections, 2010, 86, ii28-ii34.	1.9	24
17	Kinetics of maternally acquired anti-hepatitis A antibodies: Prediction of waning based on maternal or cord blood antibody levels. Vaccine, 2013, 31, 1490-1495.	3.8	22
18	Evaluation of p24-based Antiretroviral Treatment Monitoring in Pediatric HIV-1 Infection. Journal of Acquired Immune Deficiency Syndromes (1999), 2006, 41, 557-562.	2.1	10