

David Antonucci

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/11923257/david-antoniucci-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

203
papers

12,615
citations

54
h-index

109
g-index

207
ext. papers

13,958
ext. citations

5.3
avg, IF

5.55
L-index

| # | Paper | IF | Citations |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 203 | Long-Term Mortality Comparison of Patients With Acute Myocardial Infarction Complicated by Cardiogenic Shock and Treated With Culprit-Only or Multivessel Percutaneous Coronary Intervention. <i>Cardiovascular Revascularization Medicine</i> , 2021 , 22, 10-15 | 1.6 | 1 |
| 202 | Five-year clinical outcome of multicenter randomized trial comparing amphilius - with paclitaxel-eluting stents in de novo native coronary artery lesions. <i>International Journal of Cardiology</i> , 2020 , 301, 50-55 | 3.2 | 6 |
| 201 | Impact of complete percutaneous revascularization in elderly patients with chronic total occlusion. <i>Catheterization and Cardiovascular Interventions</i> , 2020 , 95, 145-153 | 2.7 | 6 |
| 200 | Ticagrelor or Prasugrel in Patients with Acute Coronary Syndromes. <i>New England Journal of Medicine</i> , 2019 , 381, 1524-1534 | 59.2 | 327 |
| 199 | One-Year Follow-Up Results From the Observational, Multicenter, Prospective, and Controlled Registry: The WALTZ All-Comers Study. <i>Clinical Medicine Insights: Cardiology</i> , 2019 , 13, 1179546819854039 ² | 3.2 | 1 |
| 198 | Patterns and Impact of Dual Antiplatelet Cessation on Cardiovascular Risk After Percutaneous Coronary Intervention in Patients With Acute Coronary Syndromes. <i>American Journal of Cardiology</i> , 2019 , 123, 709-716 | 3 | 3 |
| 197 | Lowering risk score profile during PCI in multiple vessel disease is associated with low adverse events: The ERACI risk score. <i>Cardiovascular Revascularization Medicine</i> , 2018 , 19, 792-794 | 1.6 | 4 |
| 196 | Relationship between CHADS-VASc score, coronary artery disease severity, residual platelet reactivity and long-term clinical outcomes in patients with acute coronary syndrome. <i>International Journal of Cardiology</i> , 2018 , 262, 9-13 | 3.2 | 14 |
| 195 | On-Treatment Platelet Reactivity is a Predictor of Adverse Events in Peripheral Artery Disease Patients Undergoing Percutaneous Angioplasty. <i>European Journal of Vascular and Endovascular Surgery</i> , 2018 , 56, 545-552 | 2.3 | 4 |
| 194 | Second vs. First generation drug eluting stents in multiple vessel disease and left main stenosis: Two-year follow-up of the observational, prospective, controlled, and multicenter ERACI IV registry. <i>Catheterization and Cardiovascular Interventions</i> , 2017 , 89, 37-46 | 2.7 | 14 |
| 193 | Early changes of left ventricular filling pattern after reperfused ST-elevation myocardial infarction and doxycycline therapy: Insights from the TIPTOP trial. <i>International Journal of Cardiology</i> , 2017 , 240, 43-48 | 3.2 | 7 |
| 192 | Angiographic and clinical outcome after crush of everolimus-eluting stent for distal unprotected left main disease. <i>Catheterization and Cardiovascular Interventions</i> , 2017 , 90, 72-77 | 2.7 | 3 |
| 191 | APpropriateness Assessment in Antiplatelet TherapY (APATHY) registry: Insight from current clinical practice. <i>International Journal of Cardiology</i> , 2017 , 244, 13-16 | 3.2 | 5 |
| 190 | A New Risk Score to Predict Long-Term Cardiac Mortality in Patients With Acute Myocardial Infarction Complicated by Cardiogenic Shock and Treated With Primary Percutaneous Intervention. <i>American Journal of Cardiology</i> , 2017 , 119, 351-354 | 3 | 11 |
| 189 | Clinical events beyond one year after an acute coronary syndrome: insights from the RECLOSE 2-ACS study. <i>EuroIntervention</i> , 2017 , 12, 2018-2024 | 3.1 | 9 |
| 188 | Rheolytic thrombectomy in acute myocardial infarction: Effect on microvascular obstruction, infarct size, and left ventricular remodeling. <i>Catheterization and Cardiovascular Interventions</i> , 2016 , 87, E1-8 | 2.7 | 3 |
| 187 | Predictors of restenosis following contemporary subintimal tracking and reentry technique: The importance of final TIMI flow grade. <i>Catheterization and Cardiovascular Interventions</i> , 2016 , 87, 884-92 | 2.7 | 24 |

| | | | |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----|
| 186 | High on-aspirin platelet reactivity predicts cardiac death in acute coronary syndrome patients undergoing PCI. <i>European Journal of Internal Medicine</i> , 2016 , 30, 49-54 | 3.9 | 16 |
| 185 | Reply: Personalized Antiplatelet Therapy: The Odyssey Continues. <i>JACC: Cardiovascular Interventions</i> , 2016 , 9, 106-107 | 5 | |
| 184 | Bleeding events and maintenance dose of prasugrel: BLESS pilot study. <i>Open Heart</i> , 2016 , 3, e000460 | 3 | 4 |
| 183 | Prevalence of thrombophilic disorders in takotsubo patients: the (ThROmbophilia in TAKotsubo cardiomyopathy) TROTA study. <i>Clinical Research in Cardiology</i> , 2016 , 105, 717-26 | 6.1 | 12 |
| 182 | Angiographic and Clinical Outcomes After Everolimus-Eluting Stenting for Unprotected Left Main Disease and High Anatomic Coronary Complexity. <i>JACC: Cardiovascular Interventions</i> , 2016 , 9, 1001-7 | 5 | 12 |
| 181 | Residual thrombin potential predicts cardiovascular death in acute coronary syndrome patients undergoing percutaneous coronary intervention. <i>Thrombosis Research</i> , 2016 , 147, 52-57 | 8.2 | 11 |
| 180 | Reply: Optimal Medical Therapy in the Prognosis of Coronary Artery Disease. <i>JACC: Cardiovascular Interventions</i> , 2016 , 9, 1974-5 | 5 | |
| 179 | Morphine use and myocardial reperfusion in patients with acute myocardial infarction treated with primary PCI. <i>International Journal of Cardiology</i> , 2016 , 221, 567-71 | 3.2 | 32 |
| 178 | Morphine is associated with a delayed activity of oral antiplatelet agents in patients with ST-elevation acute myocardial infarction undergoing primary percutaneous coronary intervention. <i>Circulation: Cardiovascular Interventions</i> , 2015 , 8, | 6 | 125 |
| 177 | Residual platelet reactivity to predict long-term clinical outcomes after clopidogrel loading in patients with acute coronary syndromes: comparison of different cutoff values by light transmission aggregometry from the responsiveness to clopidogrel and stent thrombosis 2-acute coronary syndrome (RECLOSE-2 ACS) study. <i>Journal of Thrombosis and Thrombolysis</i> , 2015 , 19, 76-82 | 5.1 | 23 |
| 176 | Prognostic impact of high residual platelet reactivity after chronic total occlusion percutaneous coronary intervention in patients with diabetes mellitus. <i>International Journal of Cardiology</i> , 2015 , 201, 561-7 | 3.2 | 6 |
| 175 | Modifying angiographic syntax score according to PCI strategy: lessons learnt from ERACI IV Study. <i>Cardiovascular Revascularization Medicine</i> , 2015 , 16, 418-20 | 1.6 | 10 |
| 174 | Impact on Left Ventricular Function and Remodeling and on 1-Year Outcome in Patients With Left Bundle Branch Block After Transcatheter Aortic Valve Implantation. <i>American Journal of Cardiology</i> , 2015 , 116, 125-31 | 3 | 47 |
| 173 | Prasugrel in Clopidogrel Nonresponders Undergoing Percutaneous Coronary Intervention: The RECLOSE-3 Study (REsponsiveness to CLOpidogrel and StEnt Thrombosis). <i>JACC: Cardiovascular Interventions</i> , 2015 , 8, 1563-70 | 5 | 22 |
| 172 | Matrix metalloproteinases and their tissue inhibitor after reperfused ST-elevation myocardial infarction treated with doxycycline. Insights from the TIPTOP trial. <i>International Journal of Cardiology</i> , 2015 , 197, 147-53 | 3.2 | 20 |
| 171 | Effect of diabetes on scintigraphic infarct size in STEMI patients undergoing primary angioplasty. <i>Diabetes/Metabolism Research and Reviews</i> , 2015 , 31, 322-8 | 7.5 | 6 |
| 170 | Comorbidities frequency in Takotsubo syndrome: an international collaborative systematic review including 1109 patients. <i>American Journal of Medicine</i> , 2015 , 128, 654.e11-9 | 2.4 | 110 |
| 169 | Variable underlying morphology of culprit plaques associated with ST-elevation myocardial infarction: an optical coherence tomography analysis from the SMART trial. <i>European Heart Journal Cardiovascular Imaging</i> , 2015 , 16, 1381-9 | 4.1 | 35 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|-----|
| 168 | Ticagrelor crushed tablets administration in STEMI patients: the MOJITO study. <i>Journal of the American College of Cardiology</i> , 2015 , 65, 511-2 | 15.1 | 138 |
| 167 | Managing Intracoronary Thrombus During PCI 2015 , 175-185 | | |
| 166 | Randomized comparison of ticagrelor versus prasugrel in patients with acute coronary syndrome and planned invasive strategy--design and rationale of the iNtracoronary Stenting and Antithrombotic Regimen: Rapid Early Action for Coronary Treatment (ISAR-REACT) 5 trial. <i>Journal of Cardiovascular Translational Research</i> , 2014 , 7, 91-100 | 3.3 | 73 |
| 165 | Preprocedural TIMI flow and infarct size in STEMI undergoing primary angioplasty. <i>Journal of Thrombosis and Thrombolysis</i> , 2014 , 38, 81-6 | 5.1 | 7 |
| 164 | Relationship between changes in platelet reactivity and ischemic events following percutaneous coronary intervention: a meta-regression analysis of 30 randomized trials. <i>Atherosclerosis</i> , 2014 , 234, 176-84 | 3.1 | 15 |
| 163 | Comparison of double (360 mg) ticagrelor loading dose with standard (60 mg) prasugrel loading dose in ST-elevation myocardial infarction patients: the Rapid Activity of Platelet Inhibitor Drugs (RAPID) primary PCI 2 study. <i>American Heart Journal</i> , 2014 , 167, 909-14 | 4.9 | 43 |
| 162 | Early short-term doxycycline therapy in patients with acute myocardial infarction and left ventricular dysfunction to prevent the ominous progression to adverse remodelling: the TIPTOP trial. <i>European Heart Journal</i> , 2014 , 35, 184-91 | 9.5 | 77 |
| 161 | ORAL iMmunosuppressive therapy to prevent in-Stent rEstenosiS (RAMSES) cooperation: a patient-level meta-analysis of randomized trials. <i>Atherosclerosis</i> , 2014 , 237, 410-7 | 3.1 | 11 |
| 160 | Switching from ticagrelor to prasugrel: a warning. <i>International Journal of Cardiology</i> , 2014 , 176, 1089-90,2 | 3.2 | 2 |
| 159 | Effects of a timely therapy with doxycycline on the left ventricular remodeling according to the pre-procedural TIMI flow grade in patients with ST-elevation acute myocardial infarction. <i>Basic Research in Cardiology</i> , 2014 , 109, 412 | 11.8 | 11 |
| 158 | Switching from high-dose clopidogrel to prasugrel in ACS patients undergoing PCI: a single-center experience. <i>Journal of Thrombosis and Thrombolysis</i> , 2014 , 38, 388-94 | 5.1 | 13 |
| 157 | Switching from clopidogrel to prasugrel in patients having coronary stent implantation. <i>Journal of Thrombosis and Thrombolysis</i> , 2014 , 38, 395-401 | 5.1 | 7 |
| 156 | Residual platelet reactivity and outcomes with 5 mg prasugrel therapy in elderly patients undergoing percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2014 , 176, 874-7 | 3.2 | 7 |
| 155 | Concerning antiplatelet therapy. Reply. <i>American Journal of Cardiology</i> , 2014 , 113, 2087 | 3 | 1 |
| 154 | Comparison of cost-effectiveness of oral rapamycin plus bare-metal stents versus first generation of drug-eluting stents (from the Randomized Oral Rapamycin in Argentina [ORAR] 3 trial). <i>American Journal of Cardiology</i> , 2014 , 113, 815-21 | 3 | 9 |
| 153 | Long-term follow-up of elective chronic total coronary occlusion angioplasty. <i>Journal of the American College of Cardiology</i> , 2014 , 64, 2709 | 15.1 | 3 |
| 152 | Impact of chronic total occlusion revascularization in patients with acute myocardial infarction treated by primary percutaneous coronary intervention. <i>American Journal of Cardiology</i> , 2014 , 114, 1794-800 | 3.800 | 19 |
| 151 | Smoking and infarct size among STEMI patients undergoing primary angioplasty. <i>Atherosclerosis</i> , 2014 , 233, 145-8 | 3.1 | 4 |

| | | | |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 150 | Impact of multivessel disease on infarct size among STEMI patients undergoing primary angioplasty. <i>Atherosclerosis</i> , 2014 , 234, 244-8 | 3.1 | 6 |
| 149 | Cre8Unique Technology in Challenging Daily Practice: Proceedings of a satellite symposium held at EuroPCR on 20th - 23rd May 2014 in Paris. <i>Interventional Cardiology Review</i> , 2014 , 9, 180-183 | 4.2 | 0 |
| 148 | Cessation of dual antiplatelet treatment and cardiac events after percutaneous coronary intervention (PARIS): 2 year results from a prospective observational study. <i>Lancet, The</i> , 2013 , 382, 1714-22 | 4.0 | 405 |
| 147 | Predictors of reocclusion after successful drug-eluting stent-supported percutaneous coronary intervention of chronic total occlusion. <i>Journal of the American College of Cardiology</i> , 2013 , 61, 545-50 | 15.1 | 137 |
| 146 | Percutaneous coronary intervention for multiple chronic total occlusions. <i>American Journal of Cardiology</i> , 2013 , 112, 1849-53 | 3 | 15 |
| 145 | Preinfarction angina does not affect infarct size in STEMI patients undergoing primary angioplasty. <i>Atherosclerosis</i> , 2013 , 226, 153-6 | 3.1 | 9 |
| 144 | Time-to-treatment and infarct size in STEMI patients undergoing primary angioplasty. <i>International Journal of Cardiology</i> , 2013 , 167, 1508-13 | 3.2 | 14 |
| 143 | Prognostic value of myocardial injury following transcatheter aortic valve implantation. <i>American Journal of Cardiology</i> , 2013 , 111, 1475-81 | 3 | 38 |
| 142 | Impact of coronary artery disease in elderly patients undergoing transcatheter aortic valve implantation: insight from the Italian CoreValve Registry. <i>International Journal of Cardiology</i> , 2013 , 167, 943-50 | 3.2 | 58 |
| 141 | Reply: Mechanisms affecting platelet response to antiplatelet therapy in patients with ST-segment elevation myocardial infarction. <i>Journal of the American College of Cardiology</i> , 2013 , 62, 1637 | 15.1 | |
| 140 | Reply: To PMID 23273395. <i>Journal of the American College of Cardiology</i> , 2013 , 61, 2570-1 | 15.1 | |
| 139 | Comparison of the degree of platelet aggregation inhibition with prasugrel versus clopidogrel and clinical outcomes in patients with unprotected left main disease treated with everolimus-eluting stents. <i>American Journal of Cardiology</i> , 2013 , 112, 1843-8 | 3 | 7 |
| 138 | Percutaneous recanalization of chronic total occlusions: wherein lies the body of proof?. <i>American Heart Journal</i> , 2013 , 165, 133-42 | 4.9 | 20 |
| 137 | Comparison of prasugrel and ticagrelor loading doses in ST-segment elevation myocardial infarction patients: RAPID (Rapid Activity of Platelet Inhibitor Drugs) primary PCI study. <i>Journal of the American College of Cardiology</i> , 2013 , 61, 1601-6 | 15.1 | 347 |
| 136 | Relation of gender to infarct size in patients with ST-segment elevation myocardial infarction undergoing primary angioplasty. <i>American Journal of Cardiology</i> , 2013 , 111, 936-40 | 3 | 19 |
| 135 | Comparison of variables in men versus women undergoing transcatheter aortic valve implantation for severe aortic stenosis (from Italian Multicenter CoreValve registry). <i>American Journal of Cardiology</i> , 2013 , 111, 88-93 | 3 | 51 |
| 134 | Detection of infarct size safety threshold for left ventricular ejection fraction impairment in acute myocardial infarction successfully treated with primary percutaneous coronary intervention. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2013 , 40, 542-7 | 8.8 | 2 |
| 133 | Impact of hypertension on infarct size in ST elevation myocardial infarction patients undergoing primary angioplasty. <i>Journal of Hypertension</i> , 2013 , 31, 2433-7 | 1.9 | 4 |

| | | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 132 | ARCTIC: Additional proof against antiplatelet adjusted therapy. <i>Global Cardiology Science & Practice</i> , 2013 , 2013, 130-2 | 0.7 | 1 |
| 131 | Comparison of manual thrombus aspiration with rheolytic thrombectomy in acute myocardial infarction. <i>Circulation: Cardiovascular Interventions</i> , 2013 , 6, 224-30 | 6 | 34 |
| 130 | Prognostic value of reverse left ventricular remodeling after primary angioplasty for STEMI. <i>Atherosclerosis</i> , 2012 , 222, 123-8 | 3.1 | 17 |
| 129 | High on-treatment platelet reactivity by ADP and increased risk of MACE in good clopidogrel metabolizers. <i>Platelets</i> , 2012 , 23, 586-93 | 3.6 | 14 |
| 128 | A multicenter randomized trial comparing amphilimus- with paclitaxel-eluting stents in de novo native coronary artery lesions. <i>Journal of the American College of Cardiology</i> , 2012 , 59, 1371-6 | 15.1 | 85 |
| 127 | Clinical and angiographic outcomes of patients treated with everolimus-eluting stents or first-generation Paclitaxel-eluting stents for unprotected left main disease. <i>Journal of the American College of Cardiology</i> , 2012 , 60, 1217-22 | 15.1 | 35 |
| 126 | Randomized comparison of cost-saving and effectiveness of oral rapamycin plus bare-metal stents with drug-eluting stents: three-year outcome from the randomized oral rapamycin in Argentina (ORAR) III trial. <i>Catheterization and Cardiovascular Interventions</i> , 2012 , 80, 385-94 | 2.7 | 10 |
| 125 | Residual platelet reactivity, bleedings, and adherence to treatment in patients having coronary stent implantation treated with prasugrel. <i>American Journal of Cardiology</i> , 2012 , 109, 214-8 | 3 | 65 |
| 124 | Transcatheter aortic valve implantation: 3-year outcomes of self-expanding CoreValve prosthesis. <i>European Heart Journal</i> , 2012 , 33, 969-76 | 9.5 | 226 |
| 123 | The impact of right coronary artery chronic total occlusion on clinical outcome of patients undergoing percutaneous coronary intervention for unprotected left main disease. <i>Journal of the American College of Cardiology</i> , 2011 , 58, 125-30 | 15.1 | 25 |
| 122 | Incidence and predictors of early and late mortality after transcatheter aortic valve implantation in 663 patients with severe aortic stenosis. <i>Circulation</i> , 2011 , 123, 299-308 | 16.7 | 902 |
| 121 | Natural history of tako-tsubo cardiomyopathy. <i>Chest</i> , 2011 , 139, 887-892 | 5.3 | 110 |
| 120 | Comparison of everolimus-eluting stent with paclitaxel-eluting stent in long chronic total occlusions. <i>American Journal of Cardiology</i> , 2011 , 107, 1768-71 | 3 | 27 |
| 119 | Anxiety trait in patients with stress-induced cardiomyopathy: a case-control study. <i>Clinical Research in Cardiology</i> , 2011 , 100, 523-9 | 6.1 | 42 |
| 118 | High residual platelet reactivity after clopidogrel loading and long-term cardiovascular events among patients with acute coronary syndromes undergoing PCI. <i>JAMA - Journal of the American Medical Association</i> , 2011 , 306, 1215-23 | 27.4 | 304 |
| 117 | Tako-Tsubo Cardiomyopathy: Response. <i>Chest</i> , 2011 , 140, 1101-1102 | 5.3 | 3 |
| 116 | Predictor of stent thrombosis in patients treated with turbostratic carbon-coated stent implantation for acute myocardial infarction. <i>Journal of Interventional Cardiology</i> , 2010 , 23, 554-9 | 1.8 | 3 |
| 115 | High on-treatment platelet reactivity by more than one agonist predicts 12-month follow-up cardiovascular death and non-fatal myocardial infarction in acute coronary syndrome patients receiving coronary stenting. <i>Thrombosis and Haemostasis</i> , 2010 , 104, 279-86 | 7 | 40 |

| | | | |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 114 | Comparison of AngioJet rheolytic thrombectomy before direct infarct artery stenting with direct stenting alone in patients with acute myocardial infarction. The JETSTENT trial. <i>Journal of the American College of Cardiology</i> , 2010 , 56, 1298-306 | 15.1 | 136 |
| 113 | Left ventricular remodeling after primary percutaneous coronary intervention. <i>American Heart Journal</i> , 2010 , 160, S11-5 | 4.9 | 13 |
| 112 | Abnormal response to mental stress in patients with Takotsubo cardiomyopathy detected by gated single photon emission computed tomography. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2010 , 37, 765-72 | 8.8 | 9 |
| 111 | Heart rate as an independent prognostic risk factor in patients with acute myocardial infarction undergoing primary percutaneous coronary intervention. <i>Atherosclerosis</i> , 2010 , 211, 255-9 | 3.1 | 32 |
| 110 | Evaluation of the influence of age and gender on the relationships between infarct size, infarct severity, and left ventricular ejection fraction in patients successfully treated with primary percutaneous coronary intervention. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 444-9 | 2.1 | 4 |
| 109 | Diagnostic accuracy of 64-slice computed tomography coronary angiography for the detection of in-stent restenosis: a meta-analysis. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 470-8 | 2.1 | 46 |
| 108 | Safety of immediate reversal of anticoagulation by protamine to reduce bleeding complications after infarct artery stenting for acute myocardial infarction and adjunctive abciximab therapy. <i>Journal of Thrombosis and Thrombolysis</i> , 2010 , 30, 446-51 | 5.1 | 11 |
| 107 | Safety and benefits of protamine administration to revert anticoagulation soon after coronary angioplasty. A meta-analysis. <i>Journal of Thrombosis and Thrombolysis</i> , 2010 , 30, 452-8 | 5.1 | 13 |
| 106 | Comparison of bivalirudin and unfractionated heparin plus protamine in patients with coronary heart disease undergoing percutaneous coronary intervention (from the Antithrombotic Regimens aNd Outcome [ARNO] trial). <i>American Journal of Cardiology</i> , 2010 , 105, 1053-9 | 3 | 32 |
| 105 | Usefulness of primary angioplasty in nonagenarians with acute myocardial infarction. <i>American Journal of Cardiology</i> , 2010 , 106, 770-3 | 3 | 24 |
| 104 | Impact of bivalirudin therapy in high-risk patients with acute myocardial infarction: 1-year results from the HORIZONS-AMI (Harmonizing Outcomes with RevascularizatiON and Stents in Acute Myocardial Infarction) trial. <i>JACC: Cardiovascular Interventions</i> , 2010 , 3, 796-802 | 5 | 18 |
| 103 | Long-term follow-up (four years) of unprotected left main coronary artery disease treated with paclitaxel-eluting stents (from the TRUE Registry). <i>EuroIntervention</i> , 2010 , 5, 906-916 | 3.1 | 13 |
| 102 | Optical coherence tomography in unprotected left main coronary artery stenting. <i>EuroIntervention</i> , 2010 , 6, 94-99 | 3.1 | 31 |
| 101 | Thrombectomy Devices 2010 , 144-151 | | |
| 100 | Individual patient-data meta-analysis comparing clinical outcome in patients with ST-elevation myocardial infarction treated with percutaneous coronary intervention with or without prior thrombectomy. ATTEMPT study: a pooled Analysis of Trials on ThrombEctomy in acute Myocardial Infarction based on individual Patient data. <i>Vascular Health and Risk Management</i> , 2009 , 5, 243-7 | 4.4 | 15 |
| 99 | Cardiovascular death and nonfatal myocardial infarction in acute coronary syndrome patients receiving coronary stenting are predicted by residual platelet reactivity to ADP detected by a point-of-care assay: a 12-month follow-up. <i>Circulation</i> , 2009 , 119, 237-42 | 16.7 | 459 |
| 98 | High residual platelet reactivity after clopidogrel loading and long-term clinical outcome after drug-eluting stenting for unprotected left main coronary disease. <i>Circulation</i> , 2009 , 120, 2214-21 | 16.7 | 99 |
| 97 | Relation of cytochrome P450 2C19 loss-of-function polymorphism to occurrence of drug-eluting coronary stent thrombosis. <i>American Journal of Cardiology</i> , 2009 , 103, 806-11 | 3 | 191 |

| | | | |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 96 | Effectiveness of primary percutaneous coronary interventions for stent thrombosis. <i>American Journal of Cardiology</i> , 2009 , 103, 913-6 | 3 | 21 |
| 95 | Comparison of effects of primary coronary angioplasty on left ventricular remodeling and heart failure in patients or =70 years with acute myocardial infarction. <i>American Journal of Cardiology</i> , 2009 , 104, 926-31 | 3 | 7 |
| 94 | Left bundle branch block as an electrocardiographic pattern at presentation of patients with Tako-tsubo cardiomyopathy. <i>Journal of Cardiovascular Medicine</i> , 2009 , 10, 100-3 | 1.9 | 14 |
| 93 | Clinical impact of thrombectomy in acute ST-elevation myocardial infarction: an individual patient-data pooled analysis of 11 trials. <i>European Heart Journal</i> , 2009 , 30, 2193-203 | 9.5 | 201 |
| 92 | Percutaneous coronary intervention with oral sirolimus and bare metal stents has comparable safety and efficacy to treatment with drug eluting stents, but with significant cost saving: long-term follow-up results from the randomised, controlled ORAR III (Oral Rapamycin in ARgentina) study. <i>EuroIntervention</i> , 2009 , 5, 255-64 | 3.1 | 19 |
| 91 | Clinical and angiographic follow-up of small vessel lesions treated with paclitaxel-eluting stents (from the TRUE Registry). <i>American Journal of Cardiology</i> , 2008 , 102, 1002-8 | 3 | 28 |
| 90 | Incidence and clinical impact of dual nonresponsiveness to aspirin and clopidogrel in patients with drug-eluting stents. <i>Journal of the American College of Cardiology</i> , 2008 , 52, 734-9 | 15.1 | 171 |
| 89 | Transient left ventricular apical ballooning syndrome after inadvertent epidural administration of potassium chloride. <i>International Journal of Cardiology</i> , 2008 , 124, e14-5 | 3.2 | 20 |
| 88 | Coronary stenting versus balloon angioplasty for acute myocardial infarction: a meta-regression analysis of randomized trials. <i>International Journal of Cardiology</i> , 2008 , 126, 37-44 | 3.2 | 94 |
| 87 | Left ventricular shape and function in primary coronary angioplasty. <i>International Journal of Cardiology</i> , 2008 , 125, 364-75 | 3.2 | 2 |
| 86 | A collaborative systematic review and meta-analysis on 1278 patients undergoing percutaneous drug-eluting stenting for unprotected left main coronary artery disease. <i>American Heart Journal</i> , 2008 , 155, 274-83 | 4.9 | 152 |
| 85 | Clinical implications of early mitral regurgitation in patients with reperfused acute myocardial infarction. <i>Journal of Cardiac Failure</i> , 2008 , 14, 48-54 | 3.3 | 16 |
| 84 | Impact of complete revascularization with percutaneous coronary intervention on survival in patients with at least one chronic total occlusion. <i>European Heart Journal</i> , 2008 , 29, 2336-42 | 9.5 | 172 |
| 83 | Estimate of myocardial salvage in late presentation acute myocardial infarction by comparing functional and perfusion abnormalities in predischage gated SPECT. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2008 , 35, 906-11 | 8.8 | 8 |
| 82 | Is an occluded artery better than an open one?. <i>JACC: Cardiovascular Interventions</i> , 2008 , 1, 521-3 | 5 | 1 |
| 81 | Relationship of sustained brain natriuretic peptide release after reperfused acute myocardial infarction with gated SPECT infarct measurements and its connection with collagen turnover and left ventricular remodeling. <i>Journal of Nuclear Cardiology</i> , 2008 , 15, 644-54 | 2.1 | 3 |
| 80 | Thrombotic events in high risk patients are predicted by evaluating different pathways of platelet function. <i>Thrombosis and Haemostasis</i> , 2008 , 100, 1136-45 | 7 | 31 |
| 79 | Significance of additional ST segment elevation in patients with no reflow after angioplasty for acute myocardial infarction. <i>Journal of the American Society of Echocardiography</i> , 2007 , 20, 262-9 | 5.8 | 1 |

| | | | |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 78 | Impact of platelet reactivity after clopidogrel administration on drug-eluting stent thrombosis. <i>Journal of the American College of Cardiology</i> , 2007 , 49, 2312-7 | 15.1 | 545 |
| 77 | Predictors of restenosis after treatment of bifurcational lesions with paclitaxel eluting stents: a multicenter prospective registry of 150 consecutive patients. <i>Catheterization and Cardiovascular Interventions</i> , 2007 , 69, 416-24 | 2.7 | 34 |
| 76 | Incidence, clinical findings, and outcome of women with left ventricular apical ballooning syndrome. <i>American Journal of Cardiology</i> , 2007 , 99, 182-5 | 3 | 133 |
| 75 | Relation between plasma brain natriuretic peptide, serum indexes of collagen type I turnover, and left ventricular remodeling after reperfused acute myocardial infarction. <i>American Journal of Cardiology</i> , 2007 , 99, 651-6 | 3 | 23 |
| 74 | Impact of time to treatment on myocardial reperfusion and infarct size with primary percutaneous coronary intervention for acute myocardial infarction (from the EMERALD Trial). <i>American Journal of Cardiology</i> , 2007 , 99, 1680-6 | 3 | 74 |
| 73 | Prevalence, predictors, time course, and long-term clinical implications of left ventricular functional recovery after mechanical reperfusion for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2007 , 100, 1718-22 | 3 | 47 |
| 72 | Usefulness of 64-slice multidetector computed tomography for detecting drug eluting in-stent restenosis. <i>American Journal of Cardiology</i> , 2007 , 100, 1754-8 | 3 | 33 |
| 71 | Abciximab in primary coronary stenting of ST-elevation myocardial infarction: a European meta-analysis on individual patients data with long-term follow-up. <i>European Heart Journal</i> , 2007 , 28, 443-9 | 9.5 | 190 |
| 70 | Adjunctive mechanical devices to prevent distal embolization in patients undergoing mechanical revascularization for acute myocardial infarction: a meta-analysis of randomized trials. <i>American Heart Journal</i> , 2007 , 153, 343-53 | 4.9 | 117 |
| 69 | Testing prospectively the effectiveness and safety of paclitaxel-eluting stents in over 1000 very high-risk patients: design, baseline characteristics, procedural data and in-hospital outcomes of the multicenter Taxus in Real-life Usage Evaluation (TRUE) Study. <i>International Journal of Cardiology</i> , 2007 , 117, 349-54 | 3.2 | 32 |
| 68 | Residual platelet reactivity is associated with clinical and laboratory characteristics in patients with ischemic heart disease undergoing PCI on dual antiplatelet therapy. <i>Atherosclerosis</i> , 2007 , 195, e217-23 | 3.1 | 35 |
| 67 | Appraising the effectiveness and safety of paclitaxel-eluting stents in over 1,000 very high-risk patients: overall results of the Taxus in Real-life Usage Evaluation (TRUE) registry. <i>EuroIntervention</i> , 2007 , 3, 333-9 | 3.1 | 5 |
| 66 | ST-segment analysis to predict infarct size and functional outcome in acute myocardial infarction treated with primary coronary intervention and adjunctive abciximab therapy. <i>American Journal of Cardiology</i> , 2006 , 97, 48-54 | 3 | 25 |
| 65 | Drug-eluting stent-supported percutaneous coronary intervention for chronic total coronary occlusion. <i>Catheterization and Cardiovascular Interventions</i> , 2006 , 67, 344-8 | 2.7 | 55 |
| 64 | Long-term prognostic implications of nonoptimal primary angioplasty for acute myocardial infarction. <i>Catheterization and Cardiovascular Interventions</i> , 2006 , 68, 50-5 | 2.7 | 20 |
| 63 | Drug-eluting stent supported percutaneous coronary intervention for unprotected left main disease. <i>Catheterization and Cardiovascular Interventions</i> , 2006 , 68, 225-30 | 2.7 | 30 |
| 62 | Heart failure and left ventricular remodeling after reperfused acute myocardial infarction in patients with hypertension. <i>Hypertension</i> , 2006 , 47, 706-10 | 8.5 | 27 |
| 61 | Relationship between patient's risk profile and benefits in mortality from adjunctive abciximab to mechanical revascularization for ST-segment elevation myocardial infarction: a meta-regression analysis of randomized trials. <i>Journal of the American College of Cardiology</i> , 2006 , 47, 685-6 | 15.1 | 44 |

| | | | |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 60 | Ability of mechanical reperfusion to salvage myocardium in patients with acute myocardial infarction presenting beyond 12 hours after onset of symptoms. <i>American Heart Journal</i> , 2006 , 152, 1133-8 | 4.9 | 23 |
| 59 | Routine percutaneous coronary intervention in elderly patients with cardiogenic shock complicating acute myocardial infarction. <i>American Heart Journal</i> , 2006 , 152, 903-8 | 4.9 | 19 |
| 58 | Comparison of ticlopidine vs. clopidogrel in addition to aspirin after paclitaxel-eluting stent implantation: insights from the TRUE (Taxusin Real-life Usage Evaluation) Study. <i>International Journal of Cardiology</i> , 2006 , 108, 406-7 | 3.2 | 20 |
| 57 | Myocardial infarction redefined: impact on case-load and outcome of patients with suspected acute coronary syndrome and nondiagnostic ECG at presentation. <i>International Journal of Cardiology</i> , 2006 , 111, 195-201 | 3.2 | 1 |
| 56 | How to prevent and manage complications during primary percutaneous cardiovascular interventions 2006 , 181-198 | | |
| 55 | Patient selection bias in primary percutaneous coronary intervention trials: a critical issue. <i>Country Review Ukraine</i> , 2005 , 7, 121-126 | | 1 |
| 54 | Assessment of patients with low-risk chest pain in the emergency department: Head-to-head comparison of exercise stress echocardiography and exercise myocardial SPECT. <i>American Heart Journal</i> , 2005 , 149, 894-901 | 4.9 | 51 |
| 53 | A randomized trial comparing clopidogrel versus ticlopidine therapy in patients undergoing infarct artery stenting for acute myocardial infarction with abciximab as adjunctive therapy. <i>American Heart Journal</i> , 2005 , 150, 220 | 4.9 | 17 |
| 52 | Distal microcirculatory protection during percutaneous coronary intervention in acute ST-segment elevation myocardial infarction: a randomized controlled trial. <i>JAMA - Journal of the American Medical Association</i> , 2005 , 293, 1063-72 | 27.4 | 438 |
| 51 | Comparison of the usefulness of Doppler-derived deceleration time versus plasma brain natriuretic peptide to predict left ventricular remodeling after mechanical revascularization in patients with ST-elevation acute myocardial infarction and left ventricular systolic dysfunction. <i>American Journal of Cardiology</i> , 2005 , 95, 228-34 | 3 | 13 |
| 50 | Mechanical reperfusion in patients with acute myocardial infarction presenting more than 12 hours from symptom onset: a randomized controlled trial. <i>JAMA - Journal of the American Medical Association</i> , 2005 , 293, 2865-72 | 27.4 | 183 |
| 49 | Abciximab as adjunctive therapy to reperfusion in acute ST-segment elevation myocardial infarction: a meta-analysis of randomized trials. <i>JAMA - Journal of the American Medical Association</i> , 2005 , 293, 1759-65 | 27.4 | 457 |
| 48 | Gated SPECT evaluation of outcome after abciximab-supported primary infarct artery stenting for acute myocardial infarction: the scintigraphic data of the abciximab and carbostent evaluation (ACE) randomized trial. <i>Journal of Nuclear Medicine</i> , 2005 , 46, 722-7 | 8.9 | 10 |
| 47 | Abciximab-supported infarct artery stent implantation for acute myocardial infarction and long-term survival: a prospective, multicenter, randomized trial comparing infarct artery stenting plus abciximab with stenting alone. <i>Circulation</i> , 2004 , 109, 1704-6 | 16.7 | 126 |
| 46 | Left ventricular remodeling and heart failure in diabetic patients treated with primary angioplasty for acute myocardial infarction. <i>Circulation</i> , 2004 , 110, 1974-9 | 16.7 | 60 |
| 45 | Impact of microvascular dysfunction on left ventricular remodeling and long-term clinical outcome after primary coronary angioplasty for acute myocardial infarction. <i>Circulation</i> , 2004 , 109, 1121-6 | 16.7 | 339 |
| 44 | Comparison of rheolytic thrombectomy before direct infarct artery stenting versus direct stenting alone in patients undergoing percutaneous coronary intervention for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2004 , 93, 1033-5 | 3 | 136 |
| 43 | Impact of insulin-requiring diabetes mellitus on effectiveness of reperfusion and outcome of patients undergoing primary percutaneous coronary intervention for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2004 , 93, 1170-2 | 3 | 24 |

| | | | |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 42 | Relationship of infarct size and severity versus left ventricular ejection fraction and volumes obtained from 99mTc-sestamibi gated single-photon emission computed tomography in patients treated with primary percutaneous coronary intervention. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2004 , 31, 969-74 | 8.8 | 32 |
| 41 | Direct Left Main Trunk Bifurcation Stenting in Acute Myocardial Infarction (AMI) 2004 , 7-10 | | |
| 40 | Angiographic findings, time course of regional and global left ventricular function, and clinical outcome in diabetic patients with acute myocardial infarction treated with primary percutaneous transluminal coronary angioplasty. <i>American Journal of Cardiology</i> , 2003 , 91, 544-9 | 3 | 26 |
| 39 | Comparison of impact of emergency percutaneous revascularization on outcome of patients > or =75 to those American Journal of Cardiology, 2003 , 91, 1458-61, A6 | 3 | 33 |
| 38 | Does gender affect the clinical outcome of patients with acute myocardial infarction complicated by cardiogenic shock who undergo percutaneous coronary intervention?. <i>Catheterization and Cardiovascular Interventions</i> , 2003 , 59, 423-8 | 2.7 | 14 |
| 37 | A randomized trial comparing primary infarct artery stenting with or without abciximab in acute myocardial infarction. <i>Journal of the American College of Cardiology</i> , 2003 , 42, 1879-85 | 15.1 | 214 |
| 36 | Relation between preintervention angiographic evidence of coronary collateral circulation and clinical and angiographic outcomes after primary angioplasty or stenting for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2002 , 89, 121-5 | 3 | 98 |
| 35 | Relation of time to treatment and mortality in patients with acute myocardial infarction undergoing primary coronary angioplasty. <i>American Journal of Cardiology</i> , 2002 , 89, 1248-52 | 3 | 126 |
| 34 | Abciximab therapy improves survival in patients with acute myocardial infarction complicated by early cardiogenic shock undergoing coronary artery stent implantation. <i>American Journal of Cardiology</i> , 2002 , 90, 353-7 | 3 | 59 |
| 33 | Percutaneous reperfusion of left main coronary disease complicated by acute myocardial infarction. <i>Catheterization and Cardiovascular Interventions</i> , 2002 , 56, 31-4 | 2.7 | 28 |
| 32 | Left ventricular remodeling after primary coronary angioplasty: patterns of left ventricular dilation and long-term prognostic implications. <i>Circulation</i> , 2002 , 106, 2351-7 | 16.7 | 438 |
| 31 | Abciximab therapy improves 1-month survival rate in unselected patients with acute myocardial infarction undergoing routine infarct artery stent implantation. <i>American Heart Journal</i> , 2002 , 144, 315-22 ⁹ | 4.9 | 8 |
| 30 | Effectiveness of a multidisciplinary chest pain unit for the assessment of coronary syndromes and risk stratification in the Florence area. <i>American Heart Journal</i> , 2002 , 144, 630-635 | 4.9 | 32 |
| 29 | Early detection of myocardial ischaemia in the emergency department by rest or exercise (99m)Tc tracer myocardial SPET in patients with chest pain and non-diagnostic ECG. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2001 , 28, 1806-10 | 8.8 | 33 |
| 28 | Clinical and angiographic outcomes following elective implantation of the Carbostent in patients at high risk of restenosis and target vessel failure. <i>Catheterization and Cardiovascular Interventions</i> , 2001 , 54, 420-6 | 2.7 | 23 |
| 27 | Sex-based differences in clinical and angiographic outcomes after primary angioplasty or stenting for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2001 , 87, 289-93 | 3 | 73 |
| 26 | Prognostic implications of restrictive left ventricular filling in reperfused anterior acute myocardial infarction. <i>Journal of the American College of Cardiology</i> , 2001 , 37, 793-9 | 15.1 | 82 |
| 25 | Direct infarct artery stenting without predilation and no-reflow in patients with acute myocardial infarction. <i>American Heart Journal</i> , 2001 , 142, 684-90 | 4.9 | 66 |

| | | | |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 24 | Stenting for in-stent restenosis. <i>Catheterization and Cardiovascular Interventions</i> , 2000 , 49, 376-81 | 2.7 | 18 |
| 23 | Primary stenting in nonselected patients with acute myocardial infarction: the Multilink Duet in Acute Myocardial Infarction (MIAMI) trial. <i>Catheterization and Cardiovascular Interventions</i> , 2000 , 51, 273-9 | 7 | 7 |
| 22 | Clinical and angiographic outcome after coronary arterial stenting with the carbostent. <i>American Journal of Cardiology</i> , 2000 , 85, 821-5 | 3 | 79 |
| 21 | Cost-effective analysis of primary infarct-artery stenting versus optimal primary angioplasty (the Florence Randomized Elective Stenting in Acute Coronary Occlusions [FRESCO] trial). <i>American Journal of Cardiology</i> , 2000 , 85, 1247-9 | 3 | 5 |
| 20 | Early dobutamine echocardiography predicts improvement in regional and global left ventricular function after reperfused acute myocardial infarction without residual stenosis of the infarct-related artery. <i>American Heart Journal</i> , 2000 , 139, 153-63 | 4.9 | 12 |
| 19 | A randomized study of intravenous magnesium in acute myocardial infarction treated with direct coronary angioplasty. <i>American Heart Journal</i> , 2000 , 140, 891-7 | 4.9 | 19 |
| 18 | Doppler-derived mitral deceleration time: an early strong predictor of left ventricular remodeling after reperfused anterior acute myocardial infarction. <i>Circulation</i> , 1999 , 99, 230-6 | 16.7 | 111 |
| 17 | Primary coronary infarct artery stenting in acute myocardial infarction. <i>American Journal of Cardiology</i> , 1999 , 84, 505-10 | 3 | 15 |
| 16 | Systematic primary angioplasty in octogenarian and older patients. <i>American Heart Journal</i> , 1999 , 138, 670-4 | 4.9 | 30 |
| 15 | Current role of stenting in acute myocardial infarction. <i>American Heart Journal</i> , 1999 , 138, S147-52 | 4.9 | 4 |
| 14 | Relation between ST-segment changes and myocardial perfusion evaluated by myocardial contrast echocardiography in patients with acute myocardial infarction treated with direct angioplasty. <i>American Journal of Cardiology</i> , 1998 , 82, 932-7 | 3 | 188 |
| 13 | Systematic direct angioplasty and stent-supported direct angioplasty therapy for cardiogenic shock complicating acute myocardial infarction: in-hospital and long-term survival. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 294-300 | 15.1 | 103 |
| 12 | A clinical trial comparing primary stenting of the infarct-related artery with optimal primary angioplasty for acute myocardial infarction: results from the Florence Randomized Elective Stenting in Acute Coronary Occlusions (FRESCO) trial. <i>Journal of the American College of Cardiology</i> , 1998 , 31, 1234-9 | 15.1 | 244 |
| 11 | Restenosis after coronary stenting in current clinical practice. <i>American Heart Journal</i> , 1998 , 135, 510-8 | 4.9 | 74 |
| 10 | Rapid reduction of ST-segment elevation after successful direct angioplasty in acute myocardial infarction. <i>American Journal of Cardiology</i> , 1997 , 80, 685-9 | 3 | 51 |
| 9 | Bailout coronary stenting without anticoagulation or intravascular ultrasound guidance: acute and six-month angiographic results in a series of 120 consecutive patients. <i>Catheterization and Cardiovascular Diagnosis</i> , 1997 , 41, 14-9 | | 3 |
| 8 | Influence of infarct-zone viability on left ventricular remodeling after acute myocardial infarction. <i>Circulation</i> , 1997 , 96, 3353-9 | 16.7 | 94 |
| 7 | Myocardial contrast echocardiography versus dobutamine echocardiography for predicting functional recovery after acute myocardial infarction treated with primary coronary angioplasty. <i>Journal of the American College of Cardiology</i> , 1996 , 28, 1677-83 | 15.1 | 124 |

| | | | |
|---|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 6 | Predictive value of sequential testing in screening for silent myocardial ischemia in asymptomatic middle-aged men (the ECCIS Project). <i>Cardiology</i> , 1996 , 87, 240-3 | 1.6 | 3 |
| 5 | Direct angioplasty and stenting of the infarct-related artery in acute myocardial infarction. <i>American Journal of Cardiology</i> , 1996 , 78, 568-71 | 3 | 28 |
| 4 | Bailout Palmaz-Schatz coronary stenting in 39 patients with occlusive dissection complicating conventional angioplasty. <i>Catheterization and Cardiovascular Diagnosis</i> , 1995 , 35, 204-9 | | 10 |
| 3 | Coronary angiographic findings in asymptomatic men with suspected silent myocardial ischemia (the ECCIS Project). <i>American Journal of Cardiology</i> , 1994 , 73, 960-2 | 3 | 1 |
| 2 | Coronary risk factors and silent ischemic heart disease. The ECCIS Project. <i>International Journal of Cardiology</i> , 1994 , 45, 35-43 | 3.2 | 3 |
| 1 | Epidemiology of silent myocardial ischemia in asymptomatic middle-aged men (the ECCIS Project). <i>American Journal of Cardiology</i> , 1993 , 72, 1383-8 | 3 | 44 |