

Mahmut Gumus

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11926634/publications.pdf>

Version: 2024-02-01

15
papers

142
citations

1307594

7
h-index

1199594

12
g-index

15
all docs

15
docs citations

15
times ranked

229
citing authors

#	ARTICLE	IF	CITATIONS
1	Prognostic factors of patients who received chemotherapy after cranial irradiation for non-small cell lung cancer with brain metastases: A retrospective analysis of multicenter study (Anatolian) Tj ETQq1 1 0.784304rgBT /Gverlock 10	1.2	7
2	Pretreatment PET/CT Standardized Uptake Values Play a Role in Predicting Response to Treatment and Survival in Patients with Small Cell Lung Cancer. <i>Oncology Research and Treatment</i> , 2016, 39, 130-134.	1.2	7
3	Adjuvant systemic chemotherapy with or without bevacizumab in patients with resected pulmonary metastases from colorectal cancer. <i>Thoracic Cancer</i> , 2014, 5, 398-404.	1.9	6
4	The prognostic importance of changing serum M30 and M65 values after chemotherapy in patients with advanced-stage non-small-cell lung cancer. <i>Medical Oncology</i> , 2013, 30, 551.	2.5	9
5	Adjuvant Systemic Chemotherapy with or without Bevacizumab in Patients with Resected Liver Metastases from Colorectal Cancer. <i>Oncology</i> , 2013, 84, 14-21.	1.9	22
6	Prognostic Factors for Recurrence-Free Survival in Patients with HER2-Positive Early-Stage Breast Cancer Treated with Adjuvant Trastuzumab. <i>Onkologie</i> , 2013, 36, 554-558.	0.8	2
7	Nine-week trastuzumab treatment versus 52-week trastuzumab treatment for HER2-positive early-stage breast cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2012, 138, 2145-2151.	2.5	10
8	The prognostic significance of the increase in the serum M30 and M65 values after chemotherapy and relationship between these values and clinicopathological factors in patients with advanced gastric cancer. <i>Tumor Biology</i> , 2012, 33, 2201-2208.	1.8	10
9	Pulmonary BALT Lymphoma Successfully Treated with Eight Cycles Weekly Rituximab: Report of First Case and F-18 FDG PET/CT Images. <i>Journal of Korean Medical Science</i> , 2011, 26, 574.	2.5	7
10	Is there any impact of plasma M30 and M65 levels on progression-free survival of patients with advanced gastric cancer?. <i>Cancer Chemotherapy and Pharmacology</i> , 2011, 68, 309-316.	2.3	17
11	The value of COX-2 expression in the prognostic parameters of invasive ductal carcinoma of the breast. <i>Medical Oncology</i> , 2011, 28, 703-708.	2.5	11
12	Eight-cycle rituximab therapy resulted in complete remission in primary cutaneous marginal zone lymphoma. <i>Leukemia Research</i> , 2010, 34, e160-e163.	0.8	8
13	Conjunctival malt lymphoma successfully treated with single agent rituximab therapy. <i>Leukemia Research</i> , 2009, 33, e10-e13.	0.8	16
14	Extended rituximab schedules may result in increased efficacy in pulmonary malt lymphoma. <i>Leukemia Research</i> , 2009, 33, e154-e156.	0.8	6
15	What is the optimal number of lymph nodes to be dissected in colorectal cancer surgery?. <i>Tumori</i> , 2005, 91, 168-72.	1.1	6