
Paul E Gottschall

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11948206/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Neuronal cells derived from human induced pluripotent stem cells as a functional tool of melanocortin system. Neuropeptides, 2017, 65, 10-20.	2.2	4
2	Hippocampal administration of chondroitinase ABC increases plaque-adjacent synaptic marker and diminishes amyloid burden in aged APPswe/PS1dE9 mice. Acta Neuropathologica Communications, 2015, 3, 54.	5.2	38
3	ADAMTS expression and function in central nervous system injury and disorders. Matrix Biology, 2015, 44-46, 70-76.	3.6	73
4	Altered Synaptic Marker Abundance in the Hippocampal Stratum Oriens of Ts65Dn Mice is Associated with Exuberant Expression of Versican. ASN Neuro, 2012, 4, AN20110037.	2.7	10
5	Selective Decline of Synaptic Protein Levels in the Frontal Cortex of Female Mice Deficient in the Extracellular Metalloproteinase ADAMTS1. PLoS ONE, 2012, 7, e47226.	2.5	20
6	Panel of synaptic protein ELISAs for evaluating neurological phenotype. Experimental Brain Research, 2010, 201, 885-893.	1.5	22
7	Abnormal postâ€ŧranslational and extracellular processing of brevican in plaqueâ€bearing mice overâ€expressing APPsw. Journal of Neurochemistry, 2010, 113, 784-795.	3.9	33
8	Trafficking CD11b-Positive Blood Cells Deliver Therapeutic Genes to the Brain of Amyloid-Depositing Transgenic Mice. Journal of Neuroscience, 2010, 30, 9651-9658.	3.6	116
9	Deglycosylated Anti-Aβ Antibody Dose–Response Effects on Pathology and Memory in APP Transgenic Mice. Journal of NeuroImmune Pharmacology, 2008, 3, 187-197.	4.1	22
10	Versican and brevican are expressed with distinct pathology in neonatal hypoxicâ€ i schemic injury. Journal of Neuroscience Research, 2008, 86, 1106-1114.	2.9	29
11	Discordant localization of WFA reactivity and brevican/ADAMTS-derived fragment in rodent brain. BMC Neuroscience, 2008, 9, 14.	1.9	48
12	Delayed administration of a matrix metalloproteinase inhibitor limits progressive brain injury after hypoxia-ischemia in the neonatal rat. Journal of Neuroinflammation, 2008, 5, 34.	7.2	56
13	Multimodal signaling by the ADAMTSs (a disintegrin and metalloproteinase with thrombospondin) Tj ETQq1 1 0	.784314 rg 4.1	gBT_/Overlock
14	Adeno-associated Viral (AAV) Serotype 5 Vector Mediated Gene Delivery of Endothelin-converting Enzyme Reduces Aβ Deposits in APP + PS1 Transgenic Mice. Molecular Therapy, 2008, 16, 1580-1586.	8.2	64
15	Intracranial administration of deglycosylated C-terminal-specific anti-Abeta antibody efficiently clears amyloid plaques without activating microglia in amyloid-depositing transgenic mice. Journal of Neuroinflammation, 2006, 3, 11.	7.2	42
16	Deglycosylated Anti-Amyloid-beta Antibodies Eliminate Cognitive Deficits and Reduce Parenchymal Amyloid with Minimal Vascular Consequences in Aged Amyloid Precursor Protein Transgenic Mice. Journal of Neuroscience, 2006, 26, 5340-5346.	3.6	156
17	Altered production and proteolytic processing of brevican by transforming growth factor β in cultured astrocytes. Journal of Neurochemistry, 2005, 93, 1533-1541.	3.9	41
18	Evidence for proteolytic cleavage of brevican by the ADAMTSs in the dentate gyrus after excitotoxic lesion of the mouse entorhinal cortex. BMC Neuroscience, 2005, 6, 52.	1.9	50

PAUL E GOTTSCHALL

#	Article	lF	CITATIONS
19	The effect of hypoxic–ischemic brain injury in perinatal rats on the abundance and proteolysis of brevican and NG2. Experimental Neurology, 2005, 193, 149-162.	4.1	27
20	ADAMTS4 (aggrecanase-1) cleaves human brain versican V2 at Glu405-Gln406 to generate glial hyaluronate binding protein. Biochemical Journal, 2004, 377, 787-795.	3.7	95
21	β-Amyloid induces the production of active, matrix-degrading proteases in cultured rat astrocytes. Brain Research, 2003, 970, 205-213.	2.2	83
22	Activation of the Proteolytic Activity of ADAMTS4 (Aggrecanase-1) by C-terminal Truncation. Journal of Biological Chemistry, 2002, 277, 11034-11041.	3.4	160
23	Number of A <i>β</i> Inoculations in APP+PS1 Transgenic Mice Influences Antibody Titers, Microglial Activation, and Congophilic Plaque Levels. DNA and Cell Biology, 2001, 20, 731-736.	1.9	90
24	Aβ peptide vaccination prevents memory loss in an animal model of Alzheimer's disease. Nature, 2000, 408, 982-985.	27.8	1,506
25	Regional and age-related expression of gelatinases in the brains of young and old rats after treatment with kainic acid. Neuroscience Letters, 2000, 295, 9-12.	2.1	39
26	Activated isoforms of MMP-2 are induced in U87 human glioma cells in response to β-amyloid peptide. Journal of Neuroscience Research, 1999, 55, 44-53.	2.9	31
27	Activated isoforms of MMPâ€⊋ are induced in U87 human glioma cells in response to βâ€amyloid peptide. Journal of Neuroscience Research, 1999, 55, 44-53.	2.9	1
28	Regional and differential expression of gelatinases in rat brain after systemic kainic acid or bicuculline administration. European Journal of Neuroscience, 1998, 10, 3358-3368.	2.6	105
29	Zymographic measurement of gelatinase activity in brain tissue after detergent extraction and affinity-support purification. Journal of Neuroscience Methods, 1997, 76, 15-20.	2.5	108
30	Effects of phenobarbital and interleukin-6 on cytochrome P4502B1 and 2B2 in cultured rat hepatocytes. Biochemical Pharmacology, 1996, 51, 701-706.	4.4	8
31	β-amyloid induction of gelatinase B secretion in cultured microglia. NeuroReport, 1996, 7, 3077-3080.	1.2	34
32	Regulation of Matrix Metalloproteinase Expression in Astrocytes, Microglia and Neurons. NeuroImmunoModulation, 1996, 3, 69-75.	1.8	201
33	Increased Production of Matrix Metalloproteinases in Enriched Astrocyte and Mixed Hippocampal Cultures Treated with βâ€Amyloid Peptides. Journal of Neurochemistry, 1996, 66, 1641-1647.	3.9	140
34	Differential effect of cytokines on the phenobarbital or 3-methylcholanthrene induction of P450 mediated monooxygenase activity in cultured rat hepatocytes. Biochemical Pharmacology, 1995, 49, 97-104.	4.4	63
35	Cytokines Regulate Gelatinase A and B (Matrix Metalloproteinase 2 and 9) Activity in Cultured Rat Astrocytes. Journal of Neurochemistry, 1995, 64, 1513-1520.	3.9	174
36	Regulation of interleukin-6 (IL-6) secretion in primary cultured rat astrocytes: synergism of interleukin-1 (IL-1) and pituitary adenylate cyclase activating polypeptide (PACAP). Brain Research, 1994, 637, 197-203.	2.2	133

PAUL E GOTTSCHALL

#	Article	IF	CITATIONS
37	Cytochemical characterization of anterior pituitary target cells for the neuropeptide, pituitary adenylate cyclase activating polypeptide (PACAP), using biotinylated ligands. Peptides, 1993, 14, 59-65.	2.4	80
38	Methodological Evaluation of Sites and Mechanisms of Action Involved in Neuroendocrine Effects of Cytokines. Methods in Neurosciences, 1993, , 269-293.	0.5	0
39	Increased Circulating Interleukin-1 and Interleukin-6 after Intracerebroventricular Injection of Lipopolysaccharide. Neuroendocrinology, 1992, 56, 935-938.	2.5	50
40	Increased sensitivity of glioblastoma cells to interleukin 1 after long-term incubation with dexamethasone. Molecular and Cellular Neurosciences, 1992, 3, 49-55.	2.2	9
41	Interleukin-1 $\hat{1}^2$ activation of the central nervous system. , 1992, , 27-49.		12
42	Specific binding sites for pituitary adenylate cyclase activating polypeptide (PACAP) in rat cultured astrocytes: Molecular identification and interaction with vasoactive intestinal peptide (VIP). Peptides, 1991, 12, 617-621.	2.4	60
43	Hypothalamic binding sites for pituitary adenylate cyclase activating polypeptide: characterization and molecular identification 1. FASEB Journal, 1991, 5, 194-199.	0.5	71
44	Neuropeptide Regulation of Interleukin-6 Production from the Pituitary: Stimulation by Pituitary Adenylate Cyclase Activating Polypeptide and Calcitonin Gene-Related Peptide*. Endocrinology, 1991, 129, 1797-1804.	2.8	158
45	Inhibition of Mitogen-Stimulated Proliferation of Murine Splenocytes by a Novel Neuropeptide, Pituitary Adenylate Cyclase Activating Polypeptide: A Comparative Study with Vasoactive Intestinal Peptide*. Endocrinology, 1991, 128, 728-734.	2.8	65
46	Two High Affinity Binding Sites for Pituitary Adenylate Cyclase-Activating Polypeptide Have Different Tissue Distributions*. Endocrinology, 1991, 128, 3055-3065.	2.8	313
47	Effects of Bacterial Endotoxin (Lipopolysaccharide) on FSH-Induced Granulosa Cell Activities. , 1991, , 170-177.		2
48	Characterization and Distribution of Binding Sites for the Hypothalamic Peptide, Pituitary Adenylate Cyclase-Activating Polypeptide*. Endocrinology, 1990, 127, 272-277.	2.8	311
49	Demonstration of specific binding sites for pituitary adenylate cyclase activating polypeptide (PACAP) in rat astrocytes. Biochemical and Biophysical Research Communications, 1990, 168, 1027-1033.	2.1	126
50	Molecular identification of receptor for pituitary adenylate cyclase activating polypeptide. Biochemical and Biophysical Research Communications, 1990, 171, 838-844.	2.1	72
51	INTERLEUKIN-1 <i>BETA</i> INCREASES PROSTAGLANDIN E ₂ IN RAT ASTROCYTE CULTURES: MODULATORY EFFECT OF NEUROPEPTIDES. Endocrinology, 1989, 124, 3125-3127.	2.8	178
52	Interleukin-1 suppresses follicle-stimulating hormone-induced estradiol secretion from cultured ovarian granulosa cells. Journal of Reproductive Immunology, 1989, 15, 281-290.	1.9	47
53	Interleukin-1 beta is more potent than interleukin-1 alpha in supressing follicle-stimulating hormone-induced differentiation of ovarian granulosa cells. Biochemical and Biophysical Research Communications, 1989, 163, 764-770.	2.1	46
54	Identification of a high-affinity receptor for interleukin-1 beta in rat brain. Biochemical and Biophysical Research Communications, 1988, 156, 61-67.	2.1	168

PAUL E GOTTSCHALL

#	Article	IF	CITATIONS
55	Discordance in the Effects of Interleukin-1 on Rat Granulosa Cell Differentiation Induced by Follicle-Stimulating Hormone or Activators of Adenylate Cyclase1. Biology of Reproduction, 1988, 39, 1074-1085.	2.7	53
56	Adrenocorticotropin Release Induced by Intracerebroventricular Injection of Recombinant Human Interleukin-1 in Rats: Possible Involvement of Prostaglandin*. Endocrinology, 1988, 122, 1773-1779.	2.8	230
57	Interleukin 1: an inhibitor of luteinizing hormone receptor formation in cultured rat granulosa cells. FASEB Journal, 1988, 2, 2492-2496.	0.5	86
58	INTERLEUKIN-1 STIMULATES ACTH RELEASE BY AN INDIRECT ACTION WHICH REQUIRES ENDOGENOUS CORTICOTROPIN RELEASING FACTOR. Endocrinology, 1987, 121, 1580-1582.	2.8	365
59	Interleukin-1 inhibits follicle stimulating hormone-induced differentiation in rat granulosa cells in vitro. Biochemical and Biophysical Research Communications, 1987, 149, 502-509.	2.1	118
60	Stimulation of ACTH release by human interleukin-1β, but not by interleukin-1α, in conscious, freely-moving rats. Biochemical and Biophysical Research Communications, 1987, 146, 1286-1290.	2.1	128
61	Growth Hormone Secretory Patterns in Young, Middle-Aged and Old Female Rats. Neuroendocrinology, 1987, 46, 137-142.	2.5	85
62	Possible recognition of the GnRH receptor by an antiserum against a peptide encoded by nucleotide sequence complementary to mRNA of a GnRH precursor peptide. Peptides, 1986, 7, 1137-1145.	2.4	50
63	Increased secretion of somatostatin-28 from hypothalamic neurons of aged rats in vitro. Brain Research, 1986, 380, 229-234.	2.2	66
64	Evidence for a Permanent Decline in Tuberoinfundibular Dopaminergic Neuronal Function after Chronic Estrogen Treatment Is Terminated in Fischer 344 Rats. Neuroendocrinology, 1986, 44, 211-216.	2.5	13
65	Reduced Tuberoinfundibular Dopaminergic Neuronal Function in Rats with in situ Prolactin-Secreting Pituitary Tumors. Neuroendocrinology, 1984, 38, 498-503.	2.5	34
66	L-Dopa Restores Amplitude of Growth Hormone Pulses in Old Male Rats to That Observed in Young Male Rats. Neuroendocrinology, 1982, 34, 163-168.	2.5	51